

PART II

POLICIES AND PROCEDURES

for

Early and Periodic Screening, Diagnostic & Treatment (EPSDT)

HEALTH CHECK PROGRAM (COS 600)



GEORGIA DEPARTMENT OF COMMUNITY HEALTH

DIVISION OF MEDICAID

Revised: April 1, 2024

Policy Revisions Record

Part II Policies and Procedures Manual for EPSDT Health Check Services

2024

REVISION DATE	SECTION & PAGE	REVISION DESCRIPTION	REVISION TYPE A = Added D = Deleted M = Modified	CITATION (Revision required by Regulation, Legislation, etc.)
04/2024	904. p. IX-26, p. IX-29 Table C, p. IX-31 Appendix C-1, p. C-3	Effective 1/1/2024: <ul style="list-style-type: none"> • Updated Health Check policy to allow interperiodic visits 99204, 99205, 99215. • Applicable E/M codes 99202-99205, 99211-99215 may be reported along with the preventive visit (99381-99385, 99391-99395). 	M	DCH
04/2024	904. Table A-1, p. IX-28 Table B, p. IX-30 Table C-1, p. IX-31 1003, 24, p. X-10 Appendix C-1, p. C-1	Rate tables updated to reflect current rates only. The historical list of all the attested rates/HB changes and the non-attested rates will remain housed in Appendix S.	M	DCH
04/2024	Appendix C-1, p. C-4 Appendix C-2 p. C-3, p. C-5	RSV updates: <u>Beyfortus</u> - Added 96380 & 96381, vaccine admin codes for Beyfortus, (<i>eff. 10/6/2023</i>) which replaces the previously instructed generic CPT code 96372. 96372 should no longer be reported for the administration of Beyfortus (<i>eff. 3/1/2024</i>) <u>Abrysvo</u> - Added vaccine product Abrysvo (90678) (<i>eff. 01/01/2024</i>)	A D A	DCH/ CMS DCH DCH/ CMS
04/2024	Appendix J, p. J-1	Effective April 1, 2024, Health Check providers are allowed reimbursement for the application of topical fluoride varnish. Providers should report cpt code 99188 with the EP modifier and POS 99.	A	DCH

CONTENTS

INTRODUCTION

The Early and Periodic Screening Components of EPSDT
The Diagnostic and Treatment Components of EPSDT
The Medically Fragile Child

CHAPTER 600	SPECIAL CONDITIONS OF PARTICIPATION.....	VI
	601 Enrollment	
	602 Special Conditions of Participation	
CHAPTER 700	SPECIAL ELIGIBILITY CONDITIONS	VII
CHAPTER 800	PRIOR APPROVAL/AUTHORIZATION.....	VIII
CHAPTER 900	SCOPE OF SERVICES	IX
	901 General	
	902 AAP Periodicity Schedule and Minimum Standards for Screening Components	
	903 Required Equipment and Required Location Where Services are to be Provided	
	904 Periodic, Catch-Up and Interperiodic Screening	
	905 Immunizations	
	906 Diagnosis, Treatment, and Referral	
	907 Lead Risk Assessment and Screening	
	908 Oral Health and Dental Services	
	909 Other Related Medicaid Programs	
	910 Summary of Non-Covered Services	
	911 Health Check Profile (Appointment Tracking System)	
	912 EPSDT HIPAA Referral Codes	
	913 Access to Mental Health Services	
	914 Services for Foster Care Children	
CHAPTER 1000	BASIS FOR REIMBURSEMENT.....	X
	1001 Fee For Service Reimbursement Methodology	
	1002 Vaccine for Children	
	1003 Billing Tips	

APPENDIX A	Guidelines in Screening and Reporting Elevated Blood Lead Levels
APPENDIX B	Guidelines in Screening and Reporting TB Disease and Infection
APPENDIX C-1	Vaccine Administration Codes
APPENDIX C-2	Vaccine Procedure and Diagnosis Codes
APPENDIX C-3	TB Skin Test Procedure Codes
APPENDIX C-4	Blood Lead Level Testing Procedure Codes
APPENDIX C-5	Covid-19 Vaccines & Administration Codes
APPENDIX D	Children’s Intervention Services
APPENDIX E	Medicaid Non-Emergency Medical Transportation (NEMT)
APPENDIX F	Childhood Obesity – Weight Assessment and Counseling for Nutrition and Physical Activity for Children/Adolescents
APPENDIX G	Health Check Required Equipment Form
APPENDIX H	Georgia Families
APPENDIX I	Georgia Families 360°
APPENDIX J	Preventive Oral Health: Fluoride Varnish
APPENDIX K	EPSDT HIPAA Referral Code Examples
APPENDIX L	Health Insurance Claim Form (CMS-1500)
APPENDIX M	Resources for Children in Georgia
APPENDIX N	General Claims Submission Policy for Ordering, Prescribing, or Referring (OPR) Providers
APPENDIX O	Screening Tools (CRAFFT, PHQ-2, PHQ-9)
APPENDIX P	2016-2023 Policy Revisions Record
APPENDIX Q	AAP Bright Futures Periodicity Schedule - summary of changes – 2023, 2022, 2021, 2019, 2017
APPENDIX R	<i>(intentionally blank)</i>
APPENDIX S	EPSDT Health Check Reimbursement Rates

INTRODUCTION

The Early and Periodic Screening, Diagnostic, and Treatment (EPSDT) benefit includes a comprehensive array of preventive, diagnostic, and treatment services for Medicaid eligible infants, children and adolescents under age 21, as specified in Section 1905(r) of the Social Security Act (the Act). The EPSDT benefit is also available to PeachCare for Kids® members up to 19 years of age. The EPSDT benefit is designed to assure that children receive early detection and care, so that health problems are averted or diagnosed and treated as early as possible. The goal of the EPSDT benefit is to assure that individual children get the health care they need when they need it. The EPSDT benefit also covers medically necessary diagnostic services. When a screening examination indicates the need for further evaluation of a child's health, the child should be appropriately referred for diagnosis without delay. States are required to arrange for and cover under the EPSDT benefit any Medicaid covered service listed in Section 1905(a) of the Act if that treatment or service is determined to be medically necessary to correct or ameliorate defects and physical and mental illnesses or conditions. This includes physician, nurse practitioner and hospital services; physical, speech/language, and occupational therapies; home health services, including medical equipment, supplies, and appliances; treatment for mental health and substance use disorders; treatment for vision, hearing and dental diseases and disorders, and much more. This broad coverage requirement results in a comprehensive, high-quality health benefit for children under age 21 enrolled in Medicaid.

Reference: EPSDT – A Guide for States: Coverage in the Medicaid Benefit for Children and Adolescents available at https://www.medicaid.gov/medicaid/benefits/downloads/epsdt_coverage_guide.pdf

The Health Check Program provides reimbursement for preventive health services, interperiodic visits, developmental screenings, brief emotional/behavioral assessments, hearing and vision screenings, and immunizations under the EPSDT benefit. This manual provides information pertaining to the required screening components of the EPSDT program which should be performed in accordance with the American Academy of Pediatrics (AAP) Bright Futures recommendations for preventive health/well-child check-ups. Diagnostic and Treatment (DT) services available under the EPSDT benefit are reimbursed under other program areas within the Georgia Medicaid and PeachCare for Kids® programs.

The Early and Periodic Screening Components of EPSDT

All screening components for the preventive exam should be provided as outlined in this manual. The required screening components include: a comprehensive unclothed physical examination (unclothed means to the extent necessary to conduct a full, age-appropriate examination); a comprehensive health and developmental history; developmental appraisal (including mental, emotional and behavioral health components); anticipatory guidance and health education; measurements; dental/oral health assessment; vision and hearing tests; certain laboratory procedures; lead risk assessments, and immunizations. Immunizations as needed should be given at the time of the preventive health visit as appropriate. All the age-appropriate components per the periodicity schedule and this manual must be completed and documented for each screening as appropriate. All preventive/well-child services must be provided under the EPSDT benefit following the policies and procedures as outlined in this manual.

Rev. 10/14

Rev. 07/11
Rev. 07/14
Rev. 10/14
Rev. 04/16

Rev. 10/09
Rev. 07/14
Rev. 10/14

The Diagnostic and Treatment Components of EPSDT

The Diagnostic and Treatment policies, procedures, and billing for the EPSDT benefit are found under other Georgia Medicaid programs. However, the Health Check provider may address a medical condition at the time of the preventive health visit or during an interperiodic visit. Under these circumstances, the Health Check provider bills for the office visit associated with the medical condition.

Rev. 01/10
Rev. 07/14

Provider Manuals relevant to the EPSDT benefit include, but may not be limited to:

- Advanced Nurse Practitioner Services
- Autism Spectrum Disorders (ASD) Services
- Children’s Intervention Services (CIS)
- Children’s Intervention School Services (CISS)
- Dental Services
- Diagnostic Screening and Preventive Services (DSPS)
- Durable Medical Equipment (DME) Services
- Federally Qualified Health Center (FQHC) Services
- Georgia Pediatric Program (GAPP)
- Hospice Services
- Hospital Services
- Medicaid/PeachCare for Kids® Provider Billing Manuals
- Nurse Midwifery Services
- Orthotic and Prosthetic Services
- Pharmacy Services
- Physician Assistant Services
- Physician Services
- Rural Health Clinic (RHC) Services
- Vision Care Services

Rev. 04/14
Rev. 10/14
Rev. 01/15
Rev. 01/22

Provider Manuals are available for downloading. Contact Gainwell Technologies at 1-800-766-4456 or visit the website at www.mmis.georgia.gov for more information.

Rev. 04/11
Rev. 07/17
Rev. 01/21

The Medically Fragile Child

Many medically fragile children are under the care of medical specialists. The child’s primary care provider may request Diagnostic and Treatment services for the child by documenting the medical necessity for the proposed medical service. The practitioner must state the medical reason for the requested service as it relates to the child’s medical condition or diagnosis. These providers should refer to their applicable Medicaid programs’ policies and procedures manual for policy guidelines. See Appendix M for resources available to children in Georgia and Appendix D for an explanation of services provided under the *Children’s Intervention Services* program.

Rev. 07/14
Rev. 10/14

PART II - CHAPTER 600

Special Conditions of Participation

601. Enrollment

Rev. 07/11
Rev. 04/12
Rev. 04/14
Rev. 10/14

Physicians (pediatricians, family practitioners, general practitioners, internists, and OB/GYN specialists), Advanced Nurse Practitioners (pediatric, OB/GYN, family medicine or adult medicine), Certified Nurse Midwives, Local Education Agencies (LEAs) [*school districts*], hospitals, Local Boards of Health (county health departments), Rural Health Centers (RHCs) or Federally Qualified Health Centers (FQHCs) may enroll in the Health Check Program to provide EPSDT services. Physician sponsored advanced nurse practitioners (pediatric, OB/GYN, family medicine or adult medicine) and physician's assistants may also enroll in the Health Check program but must maintain current written protocols and physician sponsorship. These non-physician providers must submit an official letter from their physician sponsor as proof of physician sponsorship at the time of enrollment and at the time their physician sponsorship changes. Providers who wish to provide Diagnostic and Treatment services should enroll in their respective Medicaid program, such as Physician Services, Advanced Nurse Practitioner Services, etc. Physicians, nurse practitioners and nurse midwives may enroll in the Health Check Program to provide EPSDT services and their respective programs to provide diagnostic and treatment services by completing only one provider data form.

Local Education Agencies, hospitals, Local Boards of Health, RHCs or FQHCs must enroll as an entity, as opposed to each provider that will be providing services enrolling individually. The enrolling entity must ensure that only staff members who meet the qualifications listed in Section 602 of this manual are providing services.

Application Process

Rev. 01/10
Rev. 10/14
Rev. 07/17

Providers who wish to enroll in the Health Check program are required to:

- Meet the Conditions of Participation in Medicaid's Part I Policies and Procedures for Medicaid and PeachCare for Kids® Manual (Part I Manual) and the special conditions listed in Section 602;
- Read the EPSDT Benefit Policy Manual prior to signing enrollment forms and;
- Complete and sign the Health Check Required Equipment form in Appendix G

In addition, it is strongly encouraged that providers submit an application for enrollment into the Vaccines For Children (VFC) Program – see Section 905.3 for more information.

Rev. 01/21

The Department of Community Health contracts with Gainwell Technologies to provide an electronic health care administration system for its contracted providers. The Gainwell Technologies field representatives are responsible for assisting Medicaid and PeachCare for Kids[®] providers with claims adjudication, the web portal and technical support. Contact Gainwell Technologies at 1-800-766-4456 for more information.

602. Special Conditions of Participation

Rev. 04/09
Rev. 01/10
Rev. 10/10
Rev. 07/11
Rev. 04/14
Rev. 07/14
Rev. 07/15
Rev. 01/16

In addition to the general Conditions of Participation contained in Part I Policies and Procedures for Medicaid and PeachCare for Kids[®], providers in the Health Check program must meet the following requirements:

- A. Physicians, including those employed or contracted by an LEA, must be currently licensed to practice medicine. (Refer to the Physician Services Manual)
- B. Nurse Practitioners, including those employed or contracted by an LEA, must maintain a current registered nurse license for the State of Georgia and current specialty certification by the appropriate certifying agent of the American Nurses Association. (Refer to the Advanced Nurse Practitioners Manual)
- C. Nurse Midwives, including those employed or contracted by an LEA, must maintain a current registered nurse license and current certification as a nurse-midwife by the American College of Nurse-Midwives (ACNW). A copy of the national certification must be on file with the Division of Medicaid. (Refer to the Nurse Midwifery Manual)
- D. Physician-sponsored providers, including those employed or contracted by an LEA, must be currently licensed to practice and must submit a copy of their license with the application. They must also maintain current written protocols, physician sponsorship and submit an official letter from their physician sponsor as proof of physician sponsorship. These providers include:
 - Certified pediatric, OB/GYN, family, general or adult nurse practitioners. A recent graduate of a Nurse Practitioner Program may enroll as a Nurse Practitioner once he/she passes the Specialty Certification exam.
 - Physician assistants must be licensed by the Georgia Board of Medical Examiners and be associated with one or more sponsoring physician(s) on file with the Composite State Board of Medical Examiners. (Refer to the Physician Assistant Services Manual)
 - Public Health registered nurses, affiliated with a Georgia local board of health, who have successfully completed the required training for expanded role nurses.

602.1 Health Check providers must provide immunizations. It is recommended the provider enroll in the VFC program and submit a VFC Provider Enrollment Letter with their Health Check Provider Enrollment Application. This is encouraged because the vaccine administration fee is the only reimbursement a provider will receive for administering vaccines otherwise available through the VFC program. (The VFC vaccines may only be used by certain populations. See Section 905.3.) For members nineteen (19) years of age through twenty (20) years of age, VFC stock is not available. Providers must use their own stock of vaccines for these Medicaid eligible members and the Division will reimburse for the vaccine product and for vaccine administration.

Rev. 07/11
Rev. 10/14

602.2 Health Check providers must submit documentation verifying they possess the necessary equipment to perform all components of the periodic screening (See Chapter 900, Section 903 for the equipment list.)

602.3 Health Check providers must determine whether members requesting a preventive health visit have already received that periodic screening. Periodic screenings for foster children in state custody are an exception to this requirement.

602.4 Health Check providers must perform, at the time of the member's preventive health visit, all of the EPSDT required components for that visit as listed below, along with those identified in the Bright Futures Periodicity Schedule (see Section 902.1). The EPSDT required components include:

- A. A comprehensive health and developmental history, developmental appraisal (including mental, emotional and behavioral)
- B. A comprehensive unclothed physical examination (unclothed means to the extent necessary to conduct a full, age-appropriate examination) including measurements
- C. Health education and anticipatory guidance for both the child and caregiver
- D. Dental/oral health assessment
- E. Vision and hearing assessments
- F. Laboratory testing (including blood lead screening appropriate for age and risk factors)
- G. Appropriate immunizations, in accordance with the pediatric and adult schedules for vaccines established by the Advisory Committee on Immunization Practices

Rev. 01/10
Rev. 07/11
Rev. 07/14
Rev. 10/14

602.5 The Health Check provider must:

Rev. 07/11
Rev. 10/11
Rev. 10/14
Rev. 01/15

- A. Use Place of Service (POS) code 99 for all preventive health services and interperiodic visits. Public Health providers – see Section 1003 for additional information;
- B. Document, in the member’s health record, all services provided during the preventive health visit;
- C. Make available for on-site audits by the Division or its agents all records related to EPSDT services. Providers must submit plans for corrective action when requested;
- D. Refer the member to other ancillary service providers for services that are not covered under the Medicaid or PeachCare for Kids® programs;
- E. Provide services in a manner consistent with the policies, procedures and requirements outlined in this manual;
- F. If performing the required laboratory testing, be in compliance with the Clinical Laboratory Improvement Amendment. Providers seeking information concerning laboratory services should contact:

Office of Health Care Facilities Regulations at (404) 657-5700

Rev. 04/11

- G. Maintain an office, clinic, or other similar physical facility, which complies with local business and building license ordinances;
- H. Provide immunizations as needed at the time of the preventive health visit. Providers not enrolled in the VFC program will not be reimbursed for the cost of the vaccine they administer to Medicaid members if those vaccines can be provided under the VFC program. They will only be reimbursed the vaccine administration fee. To enroll in the Vaccines For Children Program, please call 1-800-848-3868. For members nineteen (19) years of age through twenty (20) years of age, VFC stock is not available. Providers must use their own stock of vaccines for these Medicaid eligible members and the Division will reimburse for the vaccine product and for vaccine administration.

Rev. 07/14
Rev. 01/15

- I. Maintain legible, accurate, and complete medical records in order to support and justify the services provided. A *Medical record* is a summary of essential medical information on an individual patient including dated reports supporting claims submitted to the Division for services provided in an office, hospital, outpatient, or other place of service. Records of service shall be entered in chronological order by the practitioner who rendered the service.

Rev. 01/10

602.6 All documentation in the medical record shall be legible and shall include but not be limited to:

1. Date(s) of service
2. Patient's name and date of birth
3. Name and title of person performing the service
4. Pertinent medical history; immunizations
5. Pertinent findings on examination
6. Medications, equipment or supplies prescribed/provided
7. Recommendations for additional treatment, procedures, or consultations
8. Tests and results
9. Plan of treatment/care and outcomes
10. Refusal of care documented with signed form by responsible person for member
11. The signature of the person performing the service. The original handwritten personal signature (electronic or fax signatures are acceptable only if these documents are legible) of the person performing the service must be on each document contained in the patient's medical record. When a fax document or signature is included in the medical record, the document with the original signature must be retrievable from the original source. This includes but is not limited to progress notes and lab reports for each date of service billed.

Rev. 10/14
Rev. 01/15

NOTE: *Electronic signature is defined as "an electronic or digital method executed or adopted by a party with the intent to be bound by or to authenticate a record, which is unique to the person using it, is capable of verification, is under the sole control of the person using it, and is linked to the data in such a manner that if the data are changed the signature is invalidated." O.C.G.A. 10-12-3. (1) (1997)*

12. All documents contained in the medical record must be written in Standard English Language. Records must be available to the Georgia Division of Medicaid or its agents and to the U.S. Department of Health and Human Services upon request. Documentation must be timely, complete, and consistent with the by-laws and medical policies of the office or facility where the services are provided.

Rev. 01/16

PART II - CHAPTER 700

Special Eligibility Conditions

Rev. 01/11
Rev. 07/14
Rev. 10/14

All persons eligible for Medicaid who are less than twenty-one (21) years of age are eligible for the EPSDT benefit with the exception of women aged eighteen to twenty-one (18 to 21) who are enrolled in the Planning for Healthy Babies Program (P4HB[®]). P4HB participants are not eligible for the EPSDT benefit. All persons eligible for the PeachCare for Kids[®] program are eligible for the EPSDT benefit.

PART II - CHAPTER 800

Prior Approval/Authorization

Rev. 01/10
Rev. 07/11
Rev. 10/11
Rev. 10/14

- 801.** The Health Check provider must provide all of the required EPSDT preventive health services, as identified by the periodicity schedule and this manual, during the preventive visit in order to be reimbursed at the Health Check visit rate. If additional service needs are identified, through the screening process, that are outside the scope of practice of the EPSDT primary care provider (PCP), the member must be referred to a provider who can address those needs.

If the provider is not the member's EPSDT PCP, the provider must notify the member's PCP of the preventive health/interperiodic visit and any additional service needs identified during that visit. The member's PCP must make the appropriate referral(s).

Some EPSDT services provided to Medicaid and PeachCare for Kids[®] members may require prior authorization and/or a referral if the member has a PCP/medical home and the member's PCP/medical home does not perform those additional services.

Documentation of care rendered outside of the PCP or the medical home (such as medical records and immunization charting) must be sent to the PCP or the medical home as identified by the member within five (5) business days of the provision of those services.

- 802.** Prior authorization may be required for services rendered by Diagnostic and Treatment providers. These providers should refer to their applicable Medicaid policy and procedure manuals for a listing of the services that require prior approval.

PART II - CHAPTER 900

Scope of Services

901. General

The Health Check Program provides reimbursement for preventive health and interperiodic visits and other services provided during those visits.

Rev. 01/11
Rev. 07/11
Rev. 10/14

The Diagnostic and Treatment components of the EPSDT benefit are covered under other Georgia Medicaid programs as described previously in this manual. Those programmatic policies and procedures should be followed as specified in the appropriate related manuals (i.e., Physician Services program, etc.). Diagnostic and Treatment services are provided for identified suspicious or abnormal conditions by either the Health Check provider, if qualified to perform those services, or upon referral to an appropriate service provider of the member's choice.

902. AAP Periodicity Schedule and Georgia Minimum Standards for Screening Components

Effective April 1, 2023, the Georgia Division of Medical Assistance Plans adopted the AAP 2023 Bright Futures "Recommendations for Pediatric Health Care" Periodicity Schedule as the periodicity schedule for EPSDT visits and services. The schedule is available at [periodicity_schedule.pdf \(aap.org\)](#)

Rev. 07/11
Rev. 10/14
Rev. 10/15
Rev. 01/16
Rev. 04/16
Rev. 04/17
Rev. 07/17
Rev. 04/19
Rev. 07/19
Rev. 04/21
Rev. 07/21
Rev. 10/22
Rev. 01/23
Rev. 04/23

The updated 2023 schedule consists of revisions to Footnote 30 to update the HIV screening recommendation and extended the upper age limit from 18 to 21 years.

A copy of the summary of changes is displayed in Appendix Q.

902.1 Periodicity Schedule and Screening Sequence

The periodic intervals for screening for all Medicaid and PeachCare for Kids[®] Health Check providers, as shown on the following page, are based on the American Academy of Pediatrics' recommendations.

NOTE: *The updated 2023 Periodicity Schedule will be used for all EPSDT preventive health visits completed on or after April 1, 2023. Exception: the prenatal visit and over 21 years of age visit as listed on the schedule are not covered under the Health Check Category of Service (COS)600.*

(continued)

14. Screen for behavioral and social-emotional problems per "Promoting Optimal Development: Screening for Behavioral and Emotional Problems" (<https://doi.org/10.1542/peds.2014-3716>), "Mental Health Competencies for Pediatric Practice" (<https://doi.org/10.1542/peds.2019-2757>), "Clinical Practice Guideline for the Assessment and Treatment of Children and Adolescents With Anxiety Disorders" (<https://pubmed.ncbi.nlm.nih.gov/32439401/>), and "Screening for Anxiety in Adolescent and Adult Women: A Recommendation From the Women's Preventive Services Initiative" (<https://pubmed.ncbi.nlm.nih.gov/32510990/>). The screening should be family centered and may include asking about caregiver emotional and mental health concerns and social determinants of health, racism, poverty, and relational health. See "Poverty and Child Health in the United States" (<https://doi.org/10.1542/peds.2016-0339>), "The Impact of Racism on Child and Adolescent Health" (<https://doi.org/10.1542/peds.2019-1765>), and "Preventing Childhood Toxic Stress: Partnering With Families and Communities to Promote Relational Health" (<https://doi.org/10.1542/peds.2021-052582>).
15. A recommended assessment tool is available at <http://coraff.org>.
16. Screen adolescents for depression and suicide risk, making every effort to preserve confidentiality of the adolescent. See "Guidelines for Adolescent Depression in Primary Care (GLAD-PC): Part I. Practice Preparation, Identification, Assessment, and Initial Management" (<https://doi.org/10.1542/peds.2017-4081>), "Mental Health Competencies for Pediatric Practice" (<https://doi.org/10.1542/peds.2019-2757>), "Suicide and Suicide Attempts in Adolescents" (<https://doi.org/10.1542/peds.2016-1400>), and "The 21st Century Cures Act & Adolescent Confidentiality" (<https://www.adolescenthealth.org/Advocacy/Advocacy-Activities/2019-11/NASPPG-SAHM-Statement.aspx>).
17. At each visit, age-appropriate physical examination is essential, with infant totally undressed and older children undressed and suitably draped. See "Use of Chaperones During the Physical Examination of the Pediatric Patient" (<https://doi.org/10.1542/peds.2011-0322>).
18. These may be modified, depending on entry point to schedule and individual need.
19. Confirm initial screen was accomplished, verify results, and follow up, as appropriate. The Recommended Uniform Screening Panel (<https://www.hrsa.gov/advisory-committees/heritable-disorders/rusp/index.html>), as determined by the Secretary's Advisory Committee on Heritable Disorders in Newborns and Children, and state newborn screening laws/regulations (<https://www.babyfirsttest.org/>) establish the criteria for and coverage of newborn screening procedures and programs.
20. Verify results as soon as possible, and follow up, as appropriate.
21. Confirm initial screening was accomplished, verify results, and follow up, as appropriate. See "Hyperbilirubinemia in the Newborn Infant <35 Weeks' Gestation: An Update With Clarifications" (<https://doi.org/10.1542/peds.2009-0329>).
22. Screening for critical congenital heart disease using pulse oximetry should be performed in newborns, after 24 hours of age, before discharge from the hospital, per "Endorsement of Health and Human Services Recommendation for Pulse Oximetry Screening for Critical Congenital Heart Disease" (<https://doi.org/10.1542/peds.2011-3211>).
23. Schedules, per the AAP Committee on Infectious Diseases, are available at <https://publications.aap.org/redbook/page/immunization-schedules>. Every visit should be an opportunity to update and complete a child's immunizations.
24. Perform risk assessment or screening, as appropriate, per recommendations in the current edition of the AAP Pediatric Nutrition Policy of the American Academy of Pediatrics (Iron chapter).
25. For children at risk of lead exposure, see "Prevention of Childhood Lead Toxicity" (<https://doi.org/10.1542/peds.2016-1493>) and "Low Level Lead Exposure Harms Children: A Renewed Call for Primary Prevention" (https://www.cdc.gov/hcch/lead/docs/final_document_030712.pdf).
26. Perform risk assessments or screenings as appropriate, based on universal screening requirements for patients with Medicaid or in high prevalence areas.
27. Tuberculosis testing per recommendations of the AAP Committee on Infectious Diseases, published in the current edition of the AAP Red Book: Report of the Committee on Infectious Diseases. Testing should be performed on recognition of high-risk factors.
28. See "Integrated Guidelines for Cardiovascular Health and Risk Reduction in Children and Adolescents" (http://www.nhbi.nih.gov/guidelines/cvd_ped/index.html).
29. Adolescents should be screened for sexually transmitted infections (STIs) per recommendations in the current edition of the AAP Red Book: Report of the Committee on Infectious Diseases.
30. Screen adolescents for HIV at least once between the ages of 15 and 21, making every effort to preserve confidentiality of the adolescent, as per "Human Immunodeficiency Virus (HIV) Infection: Screening" (<https://www.uspreventiveservicestaskforce.org/uspstf/recommendation/human-immunodeficiency-virus-hiv-infection-screening>) after initial screening, youth at increased risk of HIV infection should be retested annually or more frequently, as per "Adolescents and Young Adults: The Pediatrician's Role in HIV Testing and Pre- and Postexposure HIV Prophylaxis" (<https://doi.org/10.1542/peds.2021-055207>).
31. Perform a risk assessment for hepatitis B virus (HBV) infection according to recommendations per the USPSTF (<https://www.uspreventiveservicestaskforce.org/uspstf/recommendation/hepatitis-b-virus-infection-screening>) and in the 2021–2024 edition of the AAP Red Book: Report of the Committee on Infectious Diseases, making every effort to preserve confidentiality of the patient.
32. All individuals should be screened for hepatitis C virus (HCV) infection according to the USPSTF (<https://www.uspreventiveservicestaskforce.org/uspstf/recommendation/hepatitis-c-screening>) and Centers for Disease Control and Prevention (CDC) recommendations (<https://www.cdc.gov/mmwr/volumes/69/nnr6902a1.html>) at least once between the ages of 18 and 79. Those at increased risk of HCV infection, including those who are persons with past or current injection drug use, should be tested for HCV infection and reassessed annually.
33. Perform a risk assessment, as appropriate, per "Sudden Death in the Young: Information for the Primary Care Provider" (<https://doi.org/10.1542/peds.2021-052044>).
34. See USPSTF recommendations (<https://www.uspreventiveservicestaskforce.org/uspstf/recommendation/cervical-cancer-screening>). Indications for pelvic examinations prior to age 21 are noted in "Gynecologic Examination for Adolescents in the Pediatric Office Setting" (<https://doi.org/10.1542/peds.2010-1564>).
35. Assess whether the child has a dental home. If no dental home is identified, perform a risk assessment (<https://www.aap.org/in/patient-care/oral-health/oral-health-practice-tool/>) and refer to a dental home. Recommend brushing with fluoride toothpaste in the proper dosage for age. See "Maintaining and Improving the Oral Health of Young Children" (<https://doi.org/10.1542/peds.2014-2084>).
36. Perform a risk assessment (<https://www.aap.org/in/patient-care/oral-health/oral-health-practice-tool/>). See "Maintaining and Improving the Oral Health of Young Children" (<https://doi.org/10.1542/peds.2014-2084>).
37. The USPSTF recommends that primary care clinicians apply fluoride varnish to the primary teeth of all infants and children starting at the age of primary tooth eruption (<https://www.uspreventiveservicestaskforce.org/uspstf/recommendation/prevention-of-dental-caries-in-children-younger-than-age-5-years-screening-and-interventions>). Once teeth are present, apply fluoride varnish to all children every 3 to 6 months in the primary care or dental office based on caries risk. Indications for fluoride use are noted in "Fluoride Use in Caries Prevention in the Primary Care Setting" (<https://doi.org/10.1542/peds.2020-034637>).
38. If primary water source is deficient in fluoride, consider oral fluoride supplementation. See "Fluoride Use in Caries Prevention in the Primary Care Setting" (<https://doi.org/10.1542/peds.2020-034637>).

Summary of Changes Made to the Bright Futures/AAP Recommendations for Preventive Pediatric Health Care (Periodicity Schedule)

This schedule reflects changes approved in December 2022 and published in April 2023. For updates and a list of previous changes made, visit www.aap.org/periodicityschedule.

CHANGES MADE IN DECEMBER 2022

HIV

The HIV screening recommendation has been updated to extend the upper age limit from 18 to 21 years (to account for the range in which the screening can take place) to align with recommendations of the US Preventive Services Task Force and AAP policy ("Adolescents and Young Adults: The Pediatrician's Role in HIV Testing and Pre- and Postexposure HIV Prophylaxis").

- Footnote 30 has been updated to read as follows: "Screen adolescents for HIV at least once between the ages of 15 and 21, making every effort to preserve confidentiality of the adolescent, as per 'Human Immunodeficiency Virus (HIV) Infection: Screening' (<https://www.uspreventiveservicestaskforce.org/uspstf/recommendation/human-immunodeficiency-virus-hiv-infection-screening>); after initial screening, youth at increased risk of HIV infection should be retested annually or more frequently, as per 'Adolescents and Young Adults: The Pediatrician's Role in HIV Testing and Pre- and Postexposure HIV Prophylaxis' (<https://doi.org/10.1542/peds.2021-055207>)."

CHANGES MADE IN NOVEMBER 2021

HEPATITIS B VIRUS INFECTION

Assessing risk for HBV infection has been added to occur from newborn to 21 years (to account for the range in which the risk assessment can take place) to be consistent with recommendations of the USPSTF and the 2021–2024 edition of the AAP Red Book: Report of the Committee on Infectious Diseases.

- Footnote 31 has been added to read as follows: "Perform a risk assessment for hepatitis B virus (HBV) infection according to recommendations per the USPSTF (<https://www.uspreventiveservicestaskforce.org/uspstf/recommendation/hepatitis-b-virus-infection-screening>) and in the 2021–2024 edition of the AAP Red Book: Report of the Committee on Infectious Diseases, making every effort to preserve confidentiality of the patient."

SUDDEN CARDIAC ARREST AND SUDDEN CARDIAC DEATH

Assessing risk for sudden cardiac arrest and sudden cardiac death has been added to occur from 11 to 21 years (to account for the range in which the risk assessment can take place) to be consistent with AAP policy ("Sudden Death in the Young: Information for the Primary Care Provider").

- Footnote 33 has been added to read as follows: "Perform a risk assessment, as appropriate, per 'Sudden Death in the Young: Information for the Primary Care Provider' (<https://doi.org/10.1542/peds.2021-052044>)."

DEPRESSION AND SUICIDE RISK

Screening for suicide risk has been added to the existing depression screening recommendation to be consistent with the GLAD-PC and AAP policy.

- Footnote 16 has been updated to read as follows: "Screen adolescents for depression and suicide risk, making every effort to preserve confidentiality of the adolescent. See 'Guidelines for Adolescent Depression in Primary Care (GLAD-PC): Part I. Practice Preparation, Identification, Assessment, and Initial Management' (<https://doi.org/10.1542/peds.2017-4081>), 'Mental Health Competencies for Pediatric Practice' (<https://doi.org/10.1542/peds.2019-2757>), 'Suicide and Suicide Attempts in Adolescents' (<https://doi.org/10.1542/peds.2016-1400>), and 'The 21st Century Cures Act & Adolescent Confidentiality' (<https://www.adolescenthealth.org/Advocacy/Advocacy-Activities/2019-11/NASPPG-SAHM-Statement.aspx>)."

BEHAVIORAL/SOCIAL/EMOTIONAL

The Psychosocial/Behavioral Assessment recommendation has been updated to Behavioral/Social/Emotional Screening (annually from newborn to 21 years) to align with AAP policy, the American College of Obstetricians and Gynecologists (Women's Preventive Services Initiative) recommendations, and the American Academy of Child & Adolescent Psychiatry guidelines.

- Footnote 14 has been updated to read as follows: "Screen for behavioral and social-emotional problems per 'Promoting Optimal Development: Screening for Behavioral and Emotional Problems' (<https://doi.org/10.1542/peds.2014-3716>), 'Mental Health Competencies for Pediatric Practice' (<https://doi.org/10.1542/peds.2019-2757>), 'Clinical Practice Guideline for the Assessment and Treatment of Children and Adolescents With Anxiety Disorders' (<https://pubmed.ncbi.nlm.nih.gov/32439401/>), and 'Screening for Anxiety in Adolescent and Adult Women: A Recommendation From the Women's Preventive Services Initiative' (<https://pubmed.ncbi.nlm.nih.gov/32510990/>). The screening should be family centered and may include asking about caregiver emotional and mental health concerns and social determinants of health, racism, poverty, and relational health. See 'Poverty and Child Health in the United States' (<https://doi.org/10.1542/peds.2016-0339>), 'The Impact of Racism on Child and Adolescent Health' (<https://doi.org/10.1542/peds.2019-1765>), and 'Preventing Childhood Toxic Stress: Partnering With Families and Communities to Promote Relational Health' (<https://doi.org/10.1542/peds.2021-052582>)."

FLUORIDE VARNISH

- Footnote 37 has been updated to read as follows: "The USPSTF recommends that primary care clinicians apply fluoride varnish to the primary teeth of all infants and children starting at the age of primary tooth eruption (<https://www.uspreventiveservicestaskforce.org/uspstf/recommendation/prevention-of-dental-caries-in-children-younger-than-age-5-years-screening-and-interventions>). Once teeth are present, apply fluoride varnish to all children every 3 to 6 months in the primary care or dental office based on caries risk. Indications for fluoride use are noted in 'Fluoride Use in Caries Prevention in the Primary Care Setting' (<https://doi.org/10.1542/peds.2020-034637>)."

FLUORIDE SUPPLEMENTATION

- Footnote 38 has been updated to read as follows: "If primary water source is deficient in fluoride, consider oral fluoride supplementation. See 'Fluoride Use in Caries Prevention in the Primary Care Setting' (<https://doi.org/10.1542/peds.2020-034637>)."

KEY

- = to be performed
- * = risk assessment to be performed, with appropriate action follow, if positive
- ← • → = range during which a service may be provided, with the symbol indicating the preferred age.

FOOTNOTES

Rev. 10/10
Rev. 10/14
Rev. 04/16
Rev. 07/17
Rev. 07/19
Rev. 01/23

1. If a child comes under care for the first time at any point on the schedule, or if any items are not accomplished at the suggested age, the schedule should be brought up to date at the earliest possible time.
2. A prenatal visit is recommended for parents who are at high risk, for first-time parents, and for those who request a conference. The prenatal visit should include anticipatory guidance, pertinent medical history, and a discussion of benefits of breastfeeding and planned method of feeding, per “The Prenatal Visit” (<https://doi.org/10.1542/peds.2018-1218>).
3. Newborns should have an evaluation after birth, and breastfeeding should be encouraged (and instruction and support should be offered).
4. Newborns should have an evaluation within 3 to 5 days of birth and within 48 to 72 hours after discharge from the hospital to include evaluation for feeding and jaundice. Breastfeeding newborns should receive formal breastfeeding evaluation, and their mothers should receive encouragement and instruction, as recommended in “Breastfeeding and the Use of Human Milk” (<https://doi.org/10.1542/peds.2011-3552>). Newborns discharged less than 48 hours after delivery must be examined within 48 hours of discharge, per “Hospital Stay for Healthy Term Newborn Infants” (<https://doi.org/10.1542/peds.2015-0699>).
5. Screen, per “Expert Committee Recommendations Regarding the Prevention, Assessment, and Treatment of Child and Adolescent Overweight and Obesity: Summary Report” (<https://doi.org/10.1542/peds.2007-2329C>).
6. Screening should occur per “Clinical Practice Guideline for Screening and Management of High Blood Pressure in Children and Adolescents” (<https://doi.org/10.1542/peds.2017-1904>). Blood pressure measurement in infants and children with specific risk conditions should be performed at visits before age 3 years.
7. A visual acuity screen is recommended at ages 4 and 5 years, as well as in cooperative 3-year-olds. Instrument-based screening may be used to assess risk at ages 12 and 24 months, in addition to the well visits at 3 through 5 years of age. See “Visual System Assessment in Infants, Children, and Young Adults by Pediatricians” (<https://doi.org/10.1542/peds.2015-3596>) and “Procedures for the Evaluation of the Visual System by Pediatricians” (<https://doi.org/10.1542/peds.2015-3597>).
8. Confirm initial screen was completed, verify results, and follow up, as appropriate. Newborns should be screened, per “Year 2007 Position Statement: Principles and Guidelines for Early Hearing Detection and Intervention Programs” (<https://doi.org/10.1542/peds.2007-2333>).
9. Verify results as soon as possible, and follow up, as appropriate.

10. Screen with audiometry including 6,000 and 8,000 Hz high frequencies once between 11 and 14 years, once between 15 and 17 years, and once between 18 and 21 years. See “The Sensitivity of Adolescent Hearing Screens Significantly Improves by Adding High Frequencies” (<https://www.sciencedirect.com/science/article/abs/pii/S1054139X16000483>).
11. Screening should occur per “Incorporating Recognition and Management of Perinatal Depression Into Pediatric Practice” (<https://doi.org/10.1542/peds.2018-3259>).
12. Screening should occur per “Promoting Optimal Development: Identifying Infants and Young Children With Developmental Disorders Through Developmental Surveillance and Screening” (<https://doi.org/10.1542/peds.2019-3449>).
13. Screening should occur per “Identification, Evaluation, and Management of Children With Autism Spectrum Disorder” (<https://doi.org/10.1542/peds.2019-3447>).
14. Screen for behavioral and social-emotional problems per “Promoting Optimal Development: Screening for Behavioral and Emotional Problems” (<https://doi.org/10.1542/peds.2014-3716>), “Mental Health Competencies for Pediatric Practice” (<https://doi.org/10.1542/peds.2019-2757>), “Clinical Practice Guideline for the Assessment and Treatment of Children and Adolescents With Anxiety Disorders” (<https://pubmed.ncbi.nlm.nih.gov/32439401>), and “Screening for Anxiety in Adolescent and Adult Women: A Recommendation From the Women’s Preventive Services Initiative” (<https://pubmed.ncbi.nlm.nih.gov/32510990>). The screening should be family centered and may include asking about caregiver emotional and mental health concerns and social determinants of health, racism, poverty, and relational health. See “Poverty and Child Health in the United States” (<https://doi.org/10.1542/peds.2016-0339>), “The Impact of Racism on Child and Adolescent Health” (<https://doi.org/10.1542/peds.2019-1765>), and “Preventing Childhood Toxic Stress: Partnering With Families and Communities to Promote Relational Health” (<https://doi.org/10.1542/peds.2021-052582>).
15. A recommended assessment tool is available at <http://craftt.org>.
16. Screen adolescents for depression and suicide risk, making every effort to preserve confidentiality of the adolescent. See “Guidelines for Adolescent Depression in Primary Care (GLAD-PC): Part I. Practice Preparation, Identification, Assessment, and Initial Management” (<https://doi.org/10.1542/peds.2017-4081>), “Mental Health Competencies for Pediatric Practice” (<https://doi.org/10.1542/peds.2019-2757>), “Suicide and Suicide Attempts in Adolescents” (<https://doi.org/10.1542/peds.2016-1420>), and “The 21st Century Cures Act & Adolescent Confidentiality” ([https://www.adolescenthealth.org/Advocacy/Advocacy-Activities/2019-\(1\)/NASPAG-SAHM-Statement.aspx](https://www.adolescenthealth.org/Advocacy/Advocacy-Activities/2019-(1)/NASPAG-SAHM-Statement.aspx)).
17. At each visit, age-appropriate physical examination is essential, with infant totally unclothed and older children undressed and suitably draped. See “Use of Chaperones During the Physical Examination of the Pediatric Patient” (<https://doi.org/10.1542/peds.2011-0322>).
18. These may be modified, depending on entry point into schedule and individual need.
19. Confirm initial screen was accomplished, verify results, and follow up, as appropriate. The Recommended Uniform Screening Panel (<https://www.hrsa.gov/advisory-committees/heritable-disorders/rusp/index.html>), as determined by Children The Secretary’s Advisory Committee on Heritable Disorders in Newborns and Children,

- and state newborn screening laws/regulations (<https://www.babysfirsttest.org/>) establish the criteria for and coverage of newborn screening procedures and programs
20. Verify results as soon possible, and follow up, as appropriate
 21. Confirm initial screening was accomplished, verify results, and follow up, as appropriate. See “Hyperbilirubinemia in the Newborn Infant \geq 35 Weeks’ Gestation: An Update With Clarifications” (<https://doi.org/10.1542/peds.2009-0329>).
 22. Screening for critical congenital heart disease using pulse oximetry should be performed in newborns, after 24 hours of age, before discharge from the hospital, per “Endorsement of Health and Human Services Recommendation for Pulse Oximetry Screening for Critical Congenital Heart Disease” (<https://doi.org/10.1542/peds.2011-3211>).
 23. Schedules, per the AAP Committee on Infectious Diseases, are available at <https://publications.aap.org/redbook/pages/immunization-schedules>. Every visit should be an opportunity to update and complete a child’s immunizations.
 24. Perform risk assessment or screening, as appropriate, per recommendations in the current edition of the AAP *Pediatric Nutrition: Policy of the American Academy of Pediatrics* (Iron chapter).
 25. For children at risk of lead exposure, see “Prevention of Childhood Lead Toxicity” (<https://doi.org/10.1542/peds.2016-1493>) and “Low Level Lead Exposure Harms Children: A Renewed Call for Primary Prevention” (https://www.cdc.gov/nceh/lead/docs/final_document_030712.pdf).
 26. Perform risk assessments or screenings as appropriate, based on universal screening requirements for patients with Medicaid or in high prevalence areas.
 27. Tuberculosis testing per recommendations of the AAP Committee on Infectious Diseases, published in the current edition of the AAP *Red Book: Report of the Committee on Infectious Diseases*. Testing should be performed on recognition of high-risk factors.
 28. See “Integrated Guidelines for Cardiovascular Health and Risk Reduction in Children and Adolescents” (http://www.nhlbi.nih.gov/guidelines/cvd_ped/index.htm).
 29. Adolescents should be screened for sexually transmitted infections (STIs) per recommendations in the current edition of the AAP *Red Book: Report of the Committee on Infectious Diseases*.
 30. Screen adolescents for HIV at least once between the ages of 15 and 21, making every effort to preserve confidentiality of the adolescent, as per “Human Immunodeficiency Virus (HIV) Infection: Screening” (<https://www.uspreventiveservicestaskforce.org/uspstf/recommendation/human-immunodeficiency-virus-hiv-infection-screening>); after initial screening, youth at increased risk of HIV infection should be retested annually or more frequently, as per “Adolescent and Young Adults: The Pediatrician’s Role in HIV Testing and Pre- and Postexposure HIV Prophylaxis” (<https://doi.org/10.1542/peds.2021-055207>)
 31. Perform a risk assessment for hepatitis B virus (HBV) infection according to recommendations per the USPSTF (<https://www.uspreventiveservicestaskforce.org/uspstf/recommendation/hepatitis-b-virus-infection-screening>) and in the 2021–2024 edition of the AAP *Red Book: Report of the Committee on Infectious Diseases*, making every effort to preserve confidentiality of the patient.

32. All individuals should be screened for hepatitis C virus (HCV) infection according to the USPSTF (<https://www.uspreventiveservicestaskforce.org/uspstf/recommendation/hepatitis-c-screening>) and Centers for Disease Control and Prevention (CDC) recommendations (<https://www.cdc.gov/mmwr/volumes/69/rr/rr6902a1.htm>) at least once between the ages of 18 and 79. Those at increased risk of HCV infection, including those who are persons with past or current injection drug use, should be tested for HCV infection and reassessed annually.
33. Perform a risk assessment, as appropriate, per “Sudden Death in the Young: Information for the Primary Care Provider” (<https://doi.org/10.1542/peds.2021-052044>).
34. See USPSTF recommendations (<https://www.uspreventiveservicestaskforce.org/uspstf/recommendation/cervical-cancer-screening>). Indications for pelvic examinations prior to age 21 are noted in “Gynecologic Examination for Adolescents in the Pediatric Office Setting” (<https://doi.org/10.1542/peds.2010-1564>).
35. Assess whether the child has a dental home. If no dental home is identified, perform a risk assessment (<https://www.aap.org/en/patient-care/oral-health/oral-healthpractice-tools/>) and refer to a dental home. Recommend brushing with fluoride toothpaste in the proper dosage for age. See “Maintaining and Improving the Oral Health of Young Children” (<https://doi.org/10.1542/peds.2014-2984>).
36. Perform a risk assessment (<https://www.aap.org/en/patient-care/oral-health/oralhealthpractice-tools/>). See “Maintaining and Improving the Oral Health of Young Children” (<https://doi.org/10.1542/peds.2014-2984>).
37. The USPSTF recommends that primary care clinicians apply fluoride varnish to the primary teeth of all infants and children starting at the age of primary tooth eruption (<https://www.uspreventiveservicestaskforce.org/uspstf/recommendation/preventionof-dental-carries-in-children-younger-than-age-5-years-screening-and-interventions1>). Once teeth are present, apply fluoride varnish to all children every 3 to 6 months in the primary care or dental office based on caries risk. Indications for fluoride use are noted in “Fluoride Use in Caries Prevention in the Primary Care Setting” (<https://doi.org/10.1542/peds.2020-034637>).
38. If primary water source is deficient in fluoride, consider oral fluoride supplementation. See “Fluoride Use in Caries Prevention in the Primary Care Setting” (<https://doi.org/10.1542/peds.2020-034637>).

902.2 Minimum Standards for Screening Components during the Preventive Health Visits

Required Components are specified here and in the chart and footnotes of the American Academy of Pediatrics Recommendations for Preventive Pediatric Health Care (AAP Periodicity Schedule).

Visit Components:

Every periodic health supervision (well-child) visit must include:

1. A comprehensive health, psycho-social and developmental history;
2. Documentation of vital signs;

3. An unclothed comprehensive physical examination (unclothed means to the extent necessary to conduct a full, age-appropriate examination);
4. Assessment of growth and nutritional status;
5. Assessment of immunization status and provision of appropriate immunizations. (Use the Advisory Committee on Immunization Practices (ACIP) schedules);
6. Screening for vision, hearing, and development, as per AAP guidance;
7. Laboratory testing where appropriate to age and exam findings, and in line with AAP guidance. (Some testing, if not bundled, may be covered under other programs i.e. Physician Services, DSPTS, etc. Please follow those programs' guidelines for reimbursement.);
8. Oral health screening, preventive counseling, and referral to a dentist for ongoing dental care;
9. Screening for and if suspected, reporting of child abuse and neglect;
10. Anticipatory guidance (Health Education); and
11. Referrals /follow-ups where appropriate based on history and exam findings.

Helpful materials:

<https://brightfutures.aap.org>

Rev. 10/14
Rev. 10/15
Rev. 07/16

CDC Positive Parenting Tips (handouts English and Spanish to download for families) at: <http://www.cdc.gov/ncbddd/child/infants.htm>

Immunization schedule (Recommended Immunization Schedule, 0-18 years)
[Red Book \(2021\): Report of the Committee on Infectious Diseases | AAP Books | American Academy of Pediatrics](#)

A. Age

Calculate the child's age. If a child comes under care for the first time at any point on the schedule, or if any items were not accomplished at the suggested age, the member's visit should be brought up to date.

B. History

Initial history: All ages: The history may be obtained at the time of the visit from the parent/guardian or it may be obtained through a form or checklist sent to the parent/guardian prior to the visit for completion.

History must contain, but is not limited to:

1. Present health status and past health history of member;
2. Developmental information;
3. Allergies and immunization history - allergies must be clearly and easily found in records;
4. Family history;
5. Dietary (nutrition) history;
6. Risk assessment of lead exposure; and
7. Refusal of Care documentation form (as necessary).

Documentation: Initial health history is recorded in the medical record.

Interval history: All ages: For known patients, the age-specific history may be confined to the interval since the previous evaluation. The provider must review and supplement these histories at the time of the patient's examination. Include nutrition history.

Documentation: Evidence of review.

C. Measurements

1. **Assessment of Growth:**

All ages: Growth must be measured, plotted on a graph, and recorded as outlined below.

Children younger than 2 years: Age, weight, length, and head circumference are required. Measurements should be plotted on the appropriate World Health Organization (WHO) growth chart(s).

Children 2 years of age and older: Age, weight, height, and BMI are required. Measurements must be plotted on the appropriate Centers for Disease Control and Prevention (CDC) growth chart(s). The BMI number must be plotted on the BMI-for-age growth chart to obtain a BMI percentile ranking.

The CDC and WHO growth charts are available at the following website:
<http://www.cdc.gov/growthcharts/>

Documentation: All measurements in numerical values must be recorded and plotted as indicated. All measurements outside of normal range must have an intervention. Interventions following assessments, as suggested by the CDC, are also acceptable. Please refer to Appendix F for the correct BMI diagnosis codes to be recorded on the claim. The diagnosis code must align with the BMI percentile plotted on the growth chart.

Fee For Service (FFS) EPSDT providers should not link the preventive health visit code to the BMI percentile diagnosis code on the claim.

2. **Blood Pressure Assessment:**

Children younger than 3 years: Infants and children with specific risk conditions need a blood pressure assessment. See the Bright Futures Guidance (BFG).

Children 3 years and older: Blood pressure assessment is performed at every visit.

Documentation: All measurements in numerical values must be recorded. All measurements outside of the normal range must have an intervention.

NOTE: Definitions for High Blood Pressure must follow the range published by the National High Blood Pressure Education Program in "[The Fourth Report on the Diagnosis, Evaluation, and Treatment of High Blood Pressure in Children and Adolescents.](http://www.nhlbi.nih.gov/files/docs/resources/heart/hbp_ped.pdf)"

http://www.nhlbi.nih.gov/files/docs/resources/heart/hbp_ped.pdf

Rev. 10/13
Rev. 07/14
Rev. 10/14

Rev. 10/13
Rev. 04/15

The article, “Simple Table to Identify Children and Adolescents Needing Further Evaluation of Blood Pressure,” can be found in Pediatrics 2009; 123; e972-e974; David C. Kaelber and Frieda Pickett
[Simple Table to Identify Children and Adolescents Needing Further Evaluation of Blood Pressure | Pediatrics | American Academy of Pediatrics \(aap.org\)](#)

D. Sensory Screening

1. Vision Procedure:

Rev. 10/14

Children from birth to 3 years of age: A Vision Risk Assessment is needed at every visit. This risk assessment includes: ocular history, vision assessment, external inspection of the eyes and lids, ocular motility assessment, pupil and red reflex examination.

If the risk assessment is positive, refer to an ophthalmologist.

Rev. 04/16

Children 3 years and older: A Vision Screening is required at the 3, 4, 5, 6, 8, 10, 12, and 15 year old visits. (The routine screening at age 18 has been changed to a risk assessment). A Vision Risk Assessment should be performed at all other visits. Patients uncooperative with screening and with no history, nor signs/symptoms of problems, should be re-screened within 6 months. To test visual acuity, use age appropriate tests. BFG suggests the Snellen letter or Symbol E charts. The use of alternative tests (HOTV or Matching Symbol, Faye Symbol, Allen Pictures) should be considered for preschoolers.

If the risk assessment is positive, conduct a vision screening. If the vision screening is positive, refer to an ophthalmologist.

Reminder: If a child wears eyeglasses, assessment regarding the need for referral for optometric re-evaluation must be made based on screening with eyeglasses and the length of time since the last evaluation.

Documentation: Sensory Screening documentation consists of an age appropriate assessment, assessment results (normal or abnormal) and examinations performed and results (pass/fail) data. Appropriate follow up or referral is needed for results outside of the normal range.

2. Hearing Procedure:

Rev. 10/14
Rev. 07/16

Newborns: All newborns should receive a newborn hearing screening per the AAP “Year 2007 Position Statement: Principles and Guidelines for Early Hearing Detection and Intervention Programs”
(<http://pediatrics.aappublications.org/content/120/4/898.full>)

If the newborn does not pass the hearing screening, refer for a follow-up outpatient rescreening within one month. For newborns who do not pass the rescreening, refer to an audiologist.

Georgia's [Early Hearing Detection and Intervention \(EHDI\) Program](#) is housed in the Georgia Department of Public Health along with the Newborn Metabolic Screening Program and the [Children 1st Program](#). These three programs maintain and support a comprehensive, coordinated, statewide public health screening and referral system. EHDI includes:

- Screening for hearing loss in the birthing hospital;
- Referral of newborns who do not pass the hospital screening for rescreening;
- Referral of newborns who do not pass the rescreening for diagnostic audiological evaluation; and
- Linkage to appropriate intervention for those babies diagnosed with hearing loss.

Refer to the [Georgia EHDI Program \(http://dph.georgia.gov/EHDI\)](http://dph.georgia.gov/EHDI) for further guidance.

Infants and toddlers under age 2 years: These children should be monitored for auditory skills, middle ear status, and developmental milestones (surveillance).

Infancy and Early Childhood visits: Conduct a risk assessment at each preventive visit during the Infancy and Early Childhood years (from the three to five days visit through the 3-year-old visit). If the risk assessment is positive, refer to an audiologist.

Middle Childhood and Adolescent visits: Conduct a risk assessment during the preventive visit at ages 7 years and 9 years. If the risk assessment is positive, refer to an audiologist.

Rev. 07/17

At the 4, 5-, 6-, 8- and 10-year visits: Appropriate universal hearing screening (objective) is required.

At the 11 years through 20 years visits: Screen with audiometry including 6,000 and 8,000 Hz high frequencies once between 11 and 14 years, once between 15 and 17 years, and once between 18 and 20 years. See "*The Sensitivity of Adolescent Hearing Screens Significantly Improves by Adding High Frequencies*"

[The Sensitivity of Adolescent Hearing Screens Significantly Improves by Adding High Frequencies - Journal of Adolescent Health \(jahonline.org\)](#)

Additional Guidance for Hearing Assessment Visits

Concerns identified during surveillance for children less than 2 years of age should be followed by performance of screening using a validated global screening tool (ASQ, PEDS, etc.). Those who do not pass the speech-language portion of the global screen or who have a caregiver concern should be referred immediately for further evaluation. Children with persistent middle ear effusions should be referred for otologic evaluation.

Older children who fail the risk assessment tool or screening should have appropriate intervention. Older children with persistent middle ear effusions should be referred for otologic evaluation.

Documentation: Sensory Screening documentation consists of an age appropriate assessment, assessment results (normal or abnormal) and examinations performed and results (pass/fail) data. Appropriate follow up or referral is needed for results outside of the normal range.

Patients uncooperative with screening and with no history, nor signs/symptoms of problems, should be re-screened within 6 months. This time frame is not appropriate for newborns.

E. Psychosocial/Behavioral Assessment and Developmental Surveillance

1. Surveillance:

Required for all ages: This assessment should occur with each clinical encounter with the child or adolescent. Comprehensive childhood surveillance of development includes activities that will document *social, emotional, communication, cognitive, and physical development* concerns (this content is listed at each health supervision visit in BFG under *Surveillance of Development*). Psychosocial/ behavioral surveillance will encourage activities and interventions to promote mental health and emotional well-being. See BFG Chapter 3.

Documentation: Evidence of surveillance.

2. Developmental Screening:

Required at ages 9 months, 18 months, and 30 months:

Tools must meet the following criteria:

- a) **Developmental domains:** The following domains must be included in the standardized developmental screening tool: motor (fine and gross), language, cognitive, and social-emotional.
- b) **Established Reliability:** Reliability scores of approximately 0.70 or above.
- c) **Established Findings Regarding the Validity:** Validity scores for the tool must be approximately 0.70 or above. Measures of validity must be conducted on a significant number of children and using an appropriate standardized developmental or social-emotional assessment instrument(s).
- d) **Established Sensitivity/Specificity:** Sensitivity and specificity scores of approximately 0.70 or above.

Rev. 04/12
Rev. 07/14
Rev. 01/15
Rev. 07/21

The following tools meet the above criteria and are included in the Bright Futures Recommendations for Preventive Care, which reference the updated January 2020 American Academy of Pediatrics (AAP) Statement.

- Ages and Stages Questionnaire - 3rd Edition (ASQ-3) - 1-60 months
- Parents' Evaluation of Developmental Status (PEDS) – Birth to 8 years
- Parents' Evaluation of Developmental Status - Developmental Milestones (PEDS-DM)
- Survey of Well-Being in Young Children (SWYC) – 1-65 months

Tools included in the 2006 Statement that meet the above criteria but were not listed in the 2020 Statement (as they often are not used by primary care providers in the context of routine well-childcare) include the following:

- Battelle Developmental Inventory Screening Tool (BDI-ST) – Birth to 95 months
- Bayley Infant Neuro-developmental Screen (BINS) - 3 months to 2 yrs
- Brigance Screens-II – Birth to 90 months
- Child Development Inventory (CDI) - 18 months to 6 years
- Infant Development Inventory – Birth to 18 months

The tools listed above are not specific recommendations but are examples of tools cited in Bright Futures that meet the above criteria.

Tools that do NOT meet the criteria: It is important to note that standardized tools specifically focused on one domain of development (e.g. child's socio-emotional development [ASQ-SE] or autism [M-CHAT]) are not included in the list above as this measure is anchored to recommendations related to global developmental screening using tools that identify risk for developmental, behavioral and social delays.

Rev. 04/12
Rev. 07/14
Rev. 01/15
Rev. 07/21
Rev. 04/23

REFERENCE:

Center for Medicaid and CHIP Services, Centers for Medicare & Medicaid Services - Core Set of Children's Health Care Quality Measures for Medicaid and CHIP (Child Core Set) Technical Specifications and Resource Manual for Federal Fiscal Year 2023 (February 2023)

NOTE: *More information about the developmental screening tools that meet the measure criteria is available at*

[Bright Futures Toolkit: Links to Commonly Used Screening Instruments and Tools | AAP Toolkits | American Academy of Pediatrics](#)

Rev. 01/15

Documentation: Evidence of the screening. Documentation in the medical record must include all of the following: a note indicating the date on which the screening was performed; a copy of the completed standardized tool used; and documented evidence of a screening result or screening score. If indicated, document the follow-up assessment, therapeutic interventions used, referrals made, and treatments received.

3. Autism Screening:

Required at ages 18 months and 24 months or any time parents raise a concern: The screening should be performed with an autism-specific screening tool. The Modified Checklist For Autism in Toddlers (MCHAT) is the recommended tool and downloadable at [Screening Tool Finder \(aap.org\)](http://www.aap.org) The M-CHAT is a validated developmental screening tool for toddlers between 16 and 30 months of age and should not be used for children younger than 16 months of age. The M-CHAT is designed to identify children who may benefit from a more thorough developmental and autism evaluation. The M-CHAT can be administered and scored as part of the preventive health visit and can be used by specialists or other professionals to screen for developmental delay and autism. The M-CHAT on-line version features the latest scoring system, Modified Checklist for Autism in Toddlers, Revised with Follow-Up (M-CHAT-R/F), making the results more sensitive in detecting developmental concerns.

Documentation: Evidence of the screening. Documentation in the medical record must include a note indicating the date on which the screening was performed; a copy of the screening tool used; and documented evidence of a screening result or screening score. If indicated, document the follow-up assessment, therapeutic interventions used, referrals made, and treatments received.

For guidelines related to Autism Spectrum Disorder, including a list of acceptable Autism Spectrum Disorder Assessment Tools, refer to the Autism Spectrum Disorder policy manual Section 801.

4. Tobacco, Alcohol, or Drug Use Assessment:

Required at 11 years through 20 years of age: At all adolescent (11-20 years) visits, pre-teens and teens should be asked about substance use. The screening should be performed and documented or the child referred for care at any encounter when a parent raises a concern. AAP recommends using the [CRAFFT screening tool](http://www.crafft.org) (available at <http://www.crafft.org>) for this assessment. A copy of the CRAFFT Screening Interview is available in Appendix P.

Documentation: Evidence of assessment. Screening tool must be standardized and scorable. Document the screening tool used and the screening results (i.e., CRAFFT score). If indicated, document the follow-up assessment, therapeutic intervention used, referrals made, and treatments received.

Rev. 10/14
Rev. 01/15
Rev. 07/16
Rev. 07/17

Rev. 01/22

Rev. 07/14
Rev. 10/14
Rev. 10/15
Rev. 07/16
Rev. 07/17
Rev. 10/18
Rev. 10/19

5. Depression Screening:

Required at 12 years through 20 years of age: At all adolescent (12-20 years) visits, pre-teens and teens should be asked about depression.

AAP recommends using the Patient Health Questionnaire (PHQ)-9 Modified for Teens (PHQ-A) and Patient Health Questionnaire (PHQ)-2 (see Appendix O) or other tools available in the GLAD-PC toolkit. ([Guidelines for Adolescent Depression in Primary Care - The Reach Institute](#))

The Bright Futures Instructions for use of the PHQ 2 Questionnaire are available at <https://brightfutures.aap.org>.

Documentation: Evidence of assessment. Document the screening tool used and the screening results in the medical record. If indicated, document the follow-up assessment, therapeutic intervention used, referrals made, and treatments received.

Fee For Service EPSDT providers should link the depression screening procedure code (96127) to the applicable preventive health visit ICD-10 diagnosis code. The providers should not link a depression screening diagnosis code to the preventive health visit procedure code.

6. Maternal Depression Screening:

Required at the 1-, 2-, 4- and 6-month visits: At all visits (1, 2, 4, and 6 months), the mothers of newborn children should be asked about depression.

The relevant AAP guidance, in concert with Bright Futures recommendations (<http://brightfutures.aap.org>), references screening mothers with one of the following tools the [Edinburgh Postnatal Depression Scale \(EPDS\)](#), the [Patient Health Questionnaire-9 \(PHQ-9\)](#) or the [Patient Health Questionnaire-2 \(PHQ-2\)](#), administered at 1, 2, 4 and 6 months postpartum, with follow up referral for resources and treatment.

Refer to Instruments for Recommended Universal Screening at Specific Bright Futures Visits at

[Screening Tool Finder \(aap.org\)](#) <https://psychology-tools.com/epds/>)

NOTE: Per the BF guidance, "Screening should occur per "Incorporating Recognition and Management of Perinatal and Postpartum Depression Into Pediatric Practice"

<http://pediatrics.aappublications.org/content/early/2010/10/25/peds.2010-2348>

Documentation: Evidence of assessment. Screening tool must be standardized and scorable. Document the screening tool used and the screening results in the medical record. If indicated, document the follow-up assessment, therapeutic intervention used, referrals made, and treatments received.

Rev. 10/14
Rev. 10/15
Rev. 07/16
Rev. 07/17
Rev. 04/23

Rev. 10/14
Rev. 10/15
Rev. 04/16

Rev. 07/17
Rev. 01/18

Rev. 10/18

If evidence of depression is documented, EPSDT providers may refer the mother to the Georgia Crisis and Access Line (GCAL). GCAL, a statewide toll free crisis hotline, provides access to resources and services to individuals in need of crisis management for mental health, addictive disease, and crisis services. GCAL can be reached 24 hours a day, 7 days per week at 1-800-715-4225 (GCAL) or accessed on the web at www.mygcal.com.

F. Physical Exam

1. Physical Exam:

Rev. 10/14

All children: A comprehensive physical exam is required for periodic and catch-up visits. The physical examination is the cornerstone of pediatric evaluation. Per the Federal EPSDT policy guidelines, the physical examination must be an unclothed physical inspection (unclothed means to the extent necessary to conduct a full, age-appropriate examination) that checks the general appearance of the child to determine overall health status. This process can pick up obvious physical defects, including orthopedic disorders, hernias, skin diseases, genital abnormalities and oral health needs. Physical inspection includes an examination of all organ systems such as pulmonary, cardiac, and gastrointestinal.

Documentation: Findings on all organ systems must be documented in the medical record. A checklist type form allowing documentation of normal/ abnormal findings may be utilized for recording the different organ systems. Abnormal findings require further evaluation, follow-up or parental counseling.

2. Nutrition:

Rev. 10/14
Rev. 01/15

The Federal EPSDT policy guidelines mandate assessment of nutritional status but state it can be accomplished during many different parts of the exam. “Accurate measurements of height and weight...are among the most important indices of nutritional status.” “If information suggests dietary inadequacy, obesity or other nutritional problems, further assessment is indicated.”

Documentation: Evidence of the assessment.

G. Procedures

1. Newborn Screenings

a. Newborn Blood Screening

Rev. 01/11
Rev. 10/14
Rev. 07/16

All infants under 4 months: Georgia law requires that every live born infant receive a metabolic screening for selective inherited disorders. The [Georgia Newborn Screening \(NBS\) Program](#) ensures that every newborn in Georgia is screened for 31 heritable disorders for prompt identification and treatment.

If the infant is discharged before twenty-four (24) hours after birth, a blood specimen shall be collected prior to discharge. In this case a second specimen shall be collected prior to 7 days of age.

Refer to the [Georgia NBS Program](#) for further guidance.

Documentation: All infants whose test results are unavailable at the time of the 3–5-day preventive visit must have a specimen collected immediately during this visit unless the results are pending due to processing.

The Newborn Blood Screening process may not be complete with results available before the first scheduled preventive health visit; however, these results should be actively tracked to completion and documented as soon as possible. If the results are outside the normal limits for a newborn screening disorder, the provider should ensure that the child receives prompt appropriate retesting and/or make a referral to an appropriate sub specialist.

Providers may access newborn screening results online through the State Electronic Notification Surveillance System (SendSS). Results are also available through the Georgia Public Health Laboratory’s Newborn Screening eReports web portal.

Rev. 10/12
Rev. 10/14

Information regarding the SendSS registration process is available at:

<https://sendss.state.ga.us/sendss/login.screen>

Providers may register as a user of the Georgia Department of Public Health Laboratory’s Newborn Screening eReports electronic portal by completing the [Newborn Screening eReports Web Portal Registration Form](#). The form is available at

[Microsoft Word - Newborn Screening eReports Registration Form 10-3-12 \(georgia.gov\)](#)

<http://dph.georgia.gov/sites/dph.georgia.gov/files/NBS%20eReports%20Web%20Portal%20Registration%20form.pdf>

NOTE

Rev. 07/17

Newborn Bilirubin: The screening for bilirubin concentration at the newborn visit has been added. See “Hyperbilirubinemia in the Newborn Infant > 35 Weeks’ Gestation: An Update With Clarifications” <http://pediatrics.aappublications.org/content/124/4/1193>

b. **Critical Congenital Heart Defect (CCHD) Screening**

Rev. 10/14
Rev. 01/15
Rev. 04/16
Rev. 07/16
Rev. 07/17

Screening for critical congenital heart disease using pulse oximetry should be performed in newborns, after 24 hours of age, before discharge from the hospital, per the 2011 AAP statement, “Endorsement of Health and Human Services Recommendation for Pulse Oximetry Screening for Critical Congenital Heart Disease” <http://pediatrics.aappublications.org/content/129/1/190.full>

CCHD was added to the Georgia newborn screening panel in 2014. In accordance with the Georgia Newborn Screening Program's Policy and Procedure Manual, hospitals and birthing centers shall be equipped to conduct a critical congenital heart defect (CCHD) screening test on newborns using pulse oximetry. This screening can detect 14 common congenital heart defects in newborns which could result in disability and death. The CCHD screening should be performed in newborns before discharge from the hospital.

Documentation: Refer to the [Georgia Newborn Screening Program](#) for further guidance and documentation requirements.

2. Immunizations

All children: An immunization assessment is required for all children. This is a key element of preventive health services. Immunizations, if needed and appropriate, shall be given at the time of the preventive health visit. The Federal EPSDT policy guidelines mandate the use of the current ACIP schedule at <http://www.cdc.gov/vaccines>

Documentation: All immunizations (historic and current) must be documented in the medical record and recorded in the [Georgia Registry of Immunization Transactions and Services \(GRITS\)](#). Refusals must be documented with a signed document.

3. Anemia Screening (Hematocrit and Hemoglobin)

Anemia Screening Procedure:

At 12 months: Screening must be performed on all members with documentation of a hemoglobin or hematocrit measurement.

At 4 months: Selective screening may be performed on all preterm, low birth weight infants and those not on iron fortified formula.

Anemia Risk Assessment: An anemia risk assessment is required at the 4, 15, 18, 24, and 30 months visits, and annually starting at 3 years.

Documentation: Evidence of screening, if required, and/or test results as well as any further evaluation, treatment or counseling for results outside of the normal limits. Evidence of a risk assessment performed at the 4, 15, 18, 24, and 30 months visits, and annually thereafter starting at 3 years. This can be part of the nutrition assessment.

Rev. 07/17

Rev. 10/14

4. Lead Screening

Blood Lead Risk Assessment:

The Blood Lead Risk Assessment *is* required at 6, 9 and 18 months and 3 to 6 years per the BFG periodicity schedule. A questionnaire, based on currently accepted public health guidelines, should be administered to determine if the child is at risk for lead poisoning. A recommended tool is the Georgia Healthy Homes and Lead Poisoning Prevention Program (GHHLPPP) Blood Lead Risk Assessment Questionnaire which can be found at

[Healthy Homes and Lead Poisoning Prevention | Georgia Department of Public Health](#)

[EnvHealthLeadRiskQuestionnaire \(1\).pdf](#)

When using the questionnaire, a blood lead test should be done immediately if the child is at high risk (one or more “yes” or “I don’t know” answers on the lead risk assessment questionnaire) for lead exposure. Completing this questionnaire does not count as a blood lead screening.

Note: Assessment questions are not needed if a Blood Lead Level (BLL) screening (test) will be done at the visit.

Documentation: Risk assessment findings per the Bright Futures periodicity schedule with selective BLL screening (test) if there is a positive response or a change in risk.

Blood Lead Level (BLL) Screen:

A BLL screening (test) is required at 12 and 24 months.

Children between the ages of 36 months and 72 months: All children in this age range must receive one BLL screening IF they have not previously been tested for lead exposure.

ALL **venous** sample lead screening tests conducted using any Magellan Diagnostic lead testing system should be laboratory analyzed by a properly accredited laboratory.

Documentation: Test results as well as any further evaluation, treatment or counseling for results outside of the normal limits must be documented in the medical record.

Note: Completing a lead risk assessment questionnaire **DOES NOT** count as a blood lead level screening and does not meet Medicaid requirements.

Rev. 10/14

Rev. 07/17

Rev. 04/14
Rev. 10/14
Rev. 07/15

NOTE: The Georgia Healthy Homes and Lead Poisoning Prevention Program (GHHLPPP) at 404-657-6534 has a Lead Risk Assessment Questionnaire that the provider may choose to use at [Healthy Homes and Lead Poisoning Prevention | Georgia Department of Public Health EnvHealthLeadRiskQuestionnaire \(1\).pdf](https://dph.georgia.gov/lead-screening-guidelines-children)
<https://dph.georgia.gov/lead-screening-guidelines-children>

Resource: See Appendix A: Guidelines for Elevated BLL. These must be used if a child has results outside normal limits.

5. Tuberculin Risk Assessment and Test:

Tuberculin Risk Assessment: Required at the 1, 6, 12, and 24 month visits then annually beginning at age 3 years. An assessment is given using a risk assessment questionnaire. The questionnaire should assess at least four (4) major risk factors:

Rev. 10/14
Rev. 01/15
Rev. 10/15

- Contact with TB disease
- Foreign birth
- Foreign travel to TB endemic countries; and
- Household contact with TB

The AAP Bright Futures Guidelines for Health Supervision of Infants, Children, and Adolescents recommends asking the following questions:

- 1) Was your child born in a country at high risk for tuberculosis?
- 2) Has your child traveled (had contact with resident populations) for longer than 1 week to a country a high risk for tuberculosis?
- 3) Has a family member or contact had tuberculosis or a positive tuberculin skin test?

Documentation: Validated risk assessment and responses. If positive on initial risk assessment questions, there should be a TB test recorded.

Resources: <https://brightfutures.aap.org/materials-and-tools/Pages/default.aspx>

Tuberculin Test

TB testing is not required at any age. The TB test is only administered to a child when questions are positive on the TB risk assessment or as the practitioner designates.

Documentation: If administered, a recorded Tuberculin skin test. If the practitioner needs to defer testing for reasons that cannot be validated with professionally written guidelines, consult with state TB experts. If a child cannot be given the screening test on this day, a follow-up visit is necessary. Document risk appropriate attempts to contact and re-schedule the appointment if the parent fails to keep the follow up appointment.

Rev. 01/15

If the TB skin test result for a high-risk child less than six (6) months is negative, please retest the child at six (6) months of age.

Rev. 10/14
Rev. 10/19

For more information visit the [Georgia Department of Public Health Tuberculosis \(TB\) Prevention and Control Program](#). [TB Public Health Clinic Forms](#) (reporting & notification forms, legal forms, case management/clinic forms, TB assessments), including the [Pediatric Tuberculosis \(TB\) Risk Assessment](#), are available.

6. Dyslipidemia

Risk Assessment and selective screening when indicated: At the 2, 4, 6, 8 year and adolescence (12 through 16 year) visits.

Screening: Once between 9 and 11 years and once between 17 and 20 years: Universal screening is needed if not done previously in late adolescence (see periodicity schedule).

Rev. 10/14
Rev. 07/17

Documentation: Results of risk assessment and screening. Abnormal findings during assessment or screening require further evaluation, follow-up or parental counseling.
Resource: See AAP-endorsed 2011 guidelines from the National Heart Blood and Lung Institute, “Integrated Guidelines for Cardiovascular Health and Risk Reduction in Children and Adolescents” (http://www.nhlbi.nih.gov/guidelines/cvd_ped/index.htm).

7. Sexually Transmitted Infections (STIs):

Risk Assessment: At the 11 through 20 year visits.

Screening: Adolescents should be screened for sexually transmitted infections (STIs) per recommendations in the current edition of the AAP *Red Book: Report of the Committee on Infectious Diseases.*”

Documentation: Results of risk assessment and screening. Abnormal findings during assessment or screening require further evaluation, treatment, follow-up, referral or parental counseling.

Rev. 10/14
Rev. 10/15
Rev. 07/17

8. Human Immunodeficiency Virus (HIV):

Rev. 07/17

Risk Assessment: At the 11 through 14 year and 19 through 20 year visits.

Screening: Adolescents should be screened for HIV once between the ages of 15 and 18 years, according to the USPSTF recommendations,

[Recommendation: Human Immunodeficiency Virus \(HIV\) Infection: Screening | United States Preventive Services Taskforce \(uspreventiveservicestaskforce.org\)](https://www.uspreventiveservicestaskforce.org)

making every effort to preserve confidentiality of the adolescent. Those at increased risk of HIV infection, including those who are sexually active, participate in injection drug use, or are being tested for other STIs, should be tested for HIV and reassessed annually.

Documentation: Results of risk assessment and screening. Abnormal findings during assessment or screening require further evaluation, treatment, follow-up, referral or parental counseling.

The following in-office procedure codes are available for reimbursement for the routine HIV screening:

Rev. 10/15

86701 – Analysis for antibody to HIV-1 virus

86702 – Analysis for antibody to HIV-2 virus

86703 – Analysis for antibody to HIV-1 and HIV-2 virus (Oraquick test)

87389 – Detection test for HIV-1 and HIV-2

87390 – Detection test for HIV-1

87391 – Detection test for HIV-2

Refer to the Part II Policies and Procedures For Independent Lab Services Program Manual.

9. Cervical Dysplasia (Pap Test)

Rev. 10/12

Rev. 10/14

Adolescents should not be routinely screened for cervical dysplasia prior to age 21 years.

Indications for pelvic exams prior to age 21 are noted in the “Gynecologic Examination for Adolescents in the Pediatric Office Setting” (<http://pediatrics.aappublications.org/content/126/3/583.full>)

H. Oral Health

Rev. 04/11
Rev. 10/14
Rev. 10/16
Rev. 01/17
Rev. 07/17

Every child should begin to receive oral health risk assessments by 6 months of age. The AAP recommends both the establishment of a dental home and the first dental exam no later than 12 months of age. Assessing for a dental home should occur at the 12-month and 18-month through 6-year visits.

Risk Assessment: At the 6 and 9 month visits, conduct an oral health risk assessment [oralhealth_RiskAssessmentTool.pdf \(aap.org\)](#).

Encourage the parent to select a dental home.

For the 12-, 18-, 24-, and 30-month visits, assess whether the child has a dental home. If no dental home is identified, perform a risk assessment and refer to a dental home. Recommend brushing with fluoride toothpaste in the proper dosage for age. See *Maintaining and Improving the Oral Health of Young Children*” (<http://pediatrics.aappublications.org/content/134/6/1224>). For those at high risk, consider application of fluoride varnish for caries prevention.

At 3 and 6 years: Determine if the patient has a dental home. If not, a referral must be made. If a dental home has not been established, perform a risk assessment. For those at high risk, consider application of fluoride varnish for caries prevention.

Documentation: Document a referral or inability to refer to a dental home if one has not been established. Document the risk assessment if less than 6 years and dental home not established. Document dental appointment for older children and care per AAPD periodicity schedule. Any abnormal findings must have an appropriate intervention for all children.

An oral health risk assessment tool has been developed by the AAP/Bright Futures. This tool can be accessed at [oralhealth_RiskAssessmentTool.pdf \(aap.org\)](#) (www.aap.org/oralhealth)

Rev. 10/14
Rev. 10/16
Rev. 01/17
Rev. 07/17

1. Fluoride Varnish

Rev. 04/16
Rev. 10/16

Once teeth are present, the application of fluoride varnish is required and may be applied every 3-6 months in the primary care or dental office for children between the ages of 6 months and 5 years. Indications for fluoride use are noted in the 2014 AAP clinical report “Fluoride Use in Caries Prevention in the Primary Care Setting” (<http://pediatrics.aappublications.org/cgi/doi/10.1542/peds.2014-1699>)

Documentation: Evidence that fluoride varnish was applied once between the ages of 6 months and 5 years OR evidence that the provider addressed the fluoride varnish requirement and/or its importance with the parent.

2. Fluoride Supplementation

Rev. 07/17

Starting at tooth eruption, fluoridated toothpaste is recommended.
[AAP clarifies recommendations on fluoride use in primary care | AAP News | American Academy of Pediatrics](https://www.aapd.org/media/policies_guidelines/g_fluoridetherapy.pdf)
(http://www.aapd.org/media/policies_guidelines/g_fluoridetherapy.pdf)

If primary water source is deficient in fluoride, consider oral fluoride supplementation. See “Fluoride Use in Caries Prevention in the Primary Care Setting” (<http://pediatrics.aappublications.org/content/134/3/626>)

Risk Assessment: Required at the 6 through 12-month visits, 18 month through 30 month visits, and then annually beginning at 3 years through 16 years.

Documentation: Evidence that the provider addressed the fluoride supplementation requirement and/or its importance with the parent.

I. Anticipatory Guidance Procedure:

Rev. 04/11
Rev. 10/14

For all ages: Anticipatory guidance and health education must be offered. It is a federally required component of the EPSDT preventive health visit. Age-appropriate topics/information must be presented during each visit. Providers may use oral and written information. Providers may refer to the specific guidance by age as listed in the Bright Futures Guidelines.

Note: Providers must document discussion or provision of guidance for all children on Injury and Violence Prevention. Bright Futures Guidelines recommend and DCH requires sleep positioning counseling and documentation of such at every visit for members aged birth to six (6) months. DCH encourages sleep positioning counseling through the nine (9) month visit.

Documentation: Topics or name of handout given.

J. Referral/Treatment noted between the PCP and Specialist *or* Follow-Up for Abnormal Values

Rev. 10/14

All suspicious or abnormal findings identified during an EPSDT visit must be treated or be further evaluated. The provider must either treat (if qualified) or refer all members with abnormal findings.

Documentation: Evidence of appropriate plan of care, treatment or referral for all components, results, and overriding concerns.

903. Required Equipment and Required Location Where Services Are To Be Provided

Rev. 01/11
Rev. 04/11
Rev. 10/14
Rev. 07/15
Rev. 07/17
Rev. 07/23

In addition to an examination table and routine supplies, providers **must** have appropriate equipment in order to complete the EPSDT preventive health exam, such as:

1. Scale for weighing infants and other children;
2. Measuring board or appropriate device for measuring length or height in the recumbent position for infants and children up to the age of two (2) years;
3. Measuring board or accurate device for measuring height in the vertical position for children who are over two (2) years old;
4. Blood pressure apparatus with infant, child and adult cuffs;
5. Screening audiometer;
6. Eye charts appropriate for age of the child;
7. Ophthalmoscope and otoscope;

8. Developmental/Behavioral Health screening tools and supplies for the following:
Developmental Screening - *The required developmental screenings at ages 9 months, 18 months, and 30 months **must** be accomplished using one or more of the recommended standardized developmental screening tools specified in Section 902.2.*
 - Autism Screening
 - Depression Screening
 - Maternal Depression Screening
 - Tobacco, Alcohol, or Drug Use Assessment
9. Vaccines and immunization administration supplies; and
10. Lab supplies for appropriate lab tests/screenings.
The provider may also have a Centrifuge or other device for measuring hematocrit or hemoglobin.

904. Periodic, Catch-up and Interperiodic Visits

Rev. 10/10
Rev. 10/12
Rev. 04/14
Rev. 07/14
Rev. 10/14
Rev. 01/16
Rev. 07/17
Rev. 01/23

The Georgia Department of Community Health, Division of Medicaid adopted the updated Bright Futures Periodicity Schedule and the schedule's components as the guidelines for each EPSDT preventive health visit. Please use these guidelines, the following tables and the EPSDT HIPAA referral codes (See Section 911) when billing for the EPSDT visit.

Table A: Use this table when billing for the EPSDT periodic visits of children who are on time for their visits according to the updated Bright Futures Periodicity schedule. One visit from each sequence may be billed.

Table B: Use this table when billing for the EPSDT periodic visits of children who have missed one or more of their EPSDT periodic visits according to the updated Bright Futures Periodicity schedule and need to get caught up with the Periodicity schedule.

Table C: Use this table when billing for the EPSDT interperiodic visits of children who are up to date on their periodic visits but have a medical necessity for another visit: i.e. referred to the EPSDT provider because of a suspected problem by a health, developmental, or educational professional who comes into contact with the child outside of the formal health care system or a need identified by the provider or parent.

Table A

On Time EPSDT Periodic Visit Procedure Codes

Rev. 07/11 Use the preventive visit codes (99381-99385, 99391-99395) for Medicaid-eligible and
Rev. 10/11 PeachCare for Kids® (PCK)-eligible children. All preventive visits must be coded with the EP
Rev. 04/12 modifier (Refer to Table A) and appropriate diagnosis code (Refer to Table A-2). The 25
Rev. 03/13 modifier must be included when a vaccine is administered during the preventive visit. A
Rev. 01/14 preventive visit that is not performed as specified in the periodicity schedule should be coded
Rev. 04/14 as a catch-up visit (Refer to Table B.) Catch-up visits are only available for children younger
Rev. 07/14 than three (3) years of age.

Rev. 10/14 If an abnormality/ies is encountered or a preexisting problem is addressed during the EPSDT
Rev. 01/15 Periodic visit, and if the problem/abnormality is significant enough to require additional work
Rev. 07/15 to perform the key components of a problem-oriented E/M (evaluation and management)
Rev. 10/15 service, then the appropriate Office/Outpatient code (99202-99205, 99211-99215) should also be
Rev. 01/16 reported. If the member is a new patient, *defined as one who has not received any EPSDT
Rev. 04/16 services (face-to-face services reported with a CPT code) from a practitioner or any
Rev. 07/16 practitioner within the same group practice of the exact same specialty or subspecialty within
Rev. 04/24 the past 3 years*, code the EPSDT preventive visit using the 9938x codes. If the member is an
established patient, *defined as one who has received an EPSDT service from a practitioner or
any practitioner within the same group practice of the same specialty or subspecialty within the
past 3 years*, code the EPSDT preventive visit using the 9939x codes. Use the appropriate E/M
code (99202-99205, 99211-99215) for the office visit component. (There is a reimbursement rate
differential for PCK's members as identified in Table C.)

Modifier EP and 25 should be added to the Office/Outpatient code to indicate that a significant,
separately identifiable Evaluation and Management service was provided by the same EPSDT
provider on the same day as the EPSDT Periodic visit. The appropriate EPSDT Periodic visit
code is additionally reported. If an abnormality/ies is encountered use the appropriate HIPAA
diagnosis code which relates to the medical service(s) provided.

Other helpful information

The blood lead level screening is due at the 12- and 24-month preventive visits and the preventive
visit will not be reimbursed without documentation that the blood lead level screening occurred.

Rev. 10/14 FFS providers must submit the CPT code 83655 with modifier EP and 90 or 91 along with the CPT
Rev. 10/15 code 36415 or 36416 modifier EP and appropriate ICD-10 diagnosis code (i.e., Z13.88, Z00.121,
Rev. 10/21 Z00.129, Z77.011) to signify blood lead level screening.

Reimbursement for immunization administration will be provided when vaccines are administered
and properly coded on the claim by the provider.

The appropriate EPSDT Referral Code should be documented on the EPSDT claim when an
EPSDT preventive visit has occurred. (See Section 911 – EPSDT HIPAA Referral Codes and
Appendix K – EPSDT HIPAA Referral Code Examples.)

Table A

Sequence Numbers	Periodic Exam to take place at stated age. Otherwise, code as Catch-Up Visit	Modifier = EP HIPAA Procedure Code
1	<i>Newborn Visit (performed in the hospital)</i>	
2	3-5 days	99381 or 99391
3	by 1 month	
4	2 months	
5	4 months	
6	6 months	
7	9 months	
8	12 months	99382 or 99392
9	15 months	
10	18 months	
11	24 months	
12	30 months	
13	3 years	
14	4 years	99383 or 99393
15	5 years	
16	6 years	
17	7 years	99383 or 99393
18	8 years	
19	9 years	
20	10 years	
21	11 years	99384 or 99394
22	12 years	
23	13 years	
24	14 years	
25	15 years	
26	16 years	
27	17 years	99385 or 99395
28	18 years	
29	19 years	
30	20 years	

Table A-1

Age	Preventive Visit Codes (EP Modifier)	Rates
Birth – 11 months	99381	\$107.75
	99391	\$99.99
12 months – 4 years	99382	\$112.38
	99392	\$106.92
5 years – 11 years	99383	\$117.35
	99393	\$106.58
12 years – 17 years	99384	\$132.94
	99394	\$116.82
18 years – 20 years	99385	\$129.03
	99395	\$115.86

Rev. 07/16
 Rev. 10/16
 Rev. 04/18
 Rev. 04/23
Rev. 04/24

Table A-2

ICD-10 diagnosis codes to be used when billing preventive health visits.

CPT Codes	ICD-10-CM Codes
99381, 99391 Infant (birth to 28 days)	Z00.110 Health supervision for newborn under 8 days old OR Z00.111 Health supervision for newborn 8 to 28 days old OR Z02-Z02.89 (age 0 through 20 years)
99381, 99391 Infant (29 days to 1 year)	Z00.121 Routine child health exam <i>with abnormal findings</i> OR Z00.129 Routine child health exam <i>without abnormal findings</i> OR Z02-Z02.89
99382, 99392 Early childhood (age 1-4 years) 99383, 99393 Late childhood (age 5-11 years) 99384, 99394 Adolescent (age 12-14 years)	Z00.121 or Z00.129 or Z02-Z02.89
99384, 99394 Adolescent (age 15-17 years)	Z00.121 or Z00.129 or Z00.00 General adult medical exam <i>without abnormal findings</i> OR Z00.01 General adult medical exam <i>with abnormal findings</i> Z02-Z02.89
99385, 99395 (age 18-20 years)	Z00.00 OR Z00.01 OR Z02-Z02.89

Rev. 07/16

TABLE B

Catch-Up EPSDT Visit Procedure Codes

Use Table B for Medicaid-eligible and PeachCare for Kids[®]-eligible children who have missed their EPSDT periodic visit(s). Catch-up preventive visits are only available for children younger than three (3) years of age. All catch-up preventive visits must be coded with the EP and HA modifiers and appropriate diagnosis code (see Table B). The 25 modifier must be included when a vaccine is administered during the catch-up preventive visit.

The Health Check provider must complete all missed components during this catch-up visit but may only bill for one catch-up visit (*Example - Child presents to the Health Check provider at eight (8) months of age and has missed the four- and six-month periodic visits. All components of the four- and six-month periodic visits must be included during the present catch-up visit and documentation must be provided for all periodic visit components included during this catch-up visit.*) The appropriate EPSDT Catch-Up visit procedure code along with the EP and HA modifiers and appropriate diagnosis code (see Table B) must be included on the claim.

If abnormalities are encountered or a preexisting problem is addressed during the EPSDT Catch-Up visit, and if the problem/abnormality is significant enough to require additional work to perform the key components of a problem-oriented E/M service, then the appropriate Office/Outpatient code (99202-99205, 99211-99215) should also be reported. The provider should use the appropriate HIPAA diagnosis code that relates to the medical service(s) provided and include modifiers EP and 25.

If the member is a new patient, *one who has not received any EPSDT services (defined as face-to-face services reported with a CPT code) from a practitioner or any practitioner within the same group practice of the exact same specialty or subspecialty within the past 3 years*, code the EPSDT preventive visit using the 9938x codes. If the member is an established patient, *defined as one who has received an EPSDT service from a practitioner or any practitioner within the same group practice of the same specialty or subspecialty within the past 3 years*, code the EPSDT preventive visit using the 9939x codes. Use the appropriate E/M code (99202-99205, 99211-99215) for the office visit component. (There is a reimbursement rate differential for PCK's members as identified in Table C.)

Other helpful information

The blood lead level screening is due at the 12- and 24-month preventive visits and the preventive visit will not be reimbursed without documentation that the blood lead level screening occurred.

FFS providers must submit CPT code 83655 with modifier EP and 90 or 91 along with the CPT code 36415 or 36416 modifier EP and appropriate ICD-10 diagnosis code (i.e., Z13.88, Z00.121, Z001.129, Z77.011) to signify blood lead level screening.

Reimbursement for immunization administration will be provided when vaccines are administered and properly coded on the claim by the provider.

Rev. 04/11
Rev. 07/11
Rev. 10/11
Rev. 04/12
Rev. 10/12
Rev. 03/13
Rev. 10/13
Rev. 01/14
Rev. 10/14
Rev. 01/15
Rev. 07/15
Rev. 10/15
Rev. 07/16
Rev. 04/24

Rev. 10/14
Rev. 10/15
Rev. 10/21

The appropriate EPSDT Referral Code should be documented on the EPSDT claim when an EPSDT periodic visit has occurred. (See Section 911 – EPSDT HIPAA Referral Codes and Appendix K – EPSDT HIPAA Referral Code Examples.)

Table B

Catch-Up Preventive Health Visits				
Age of Child	HIPAA Modifiers	HIPAA Proc Code	Fee For Service Reimbursement	ICD-10-CM Codes
0 days through 11 months	EP HA	99381	\$107.75	<u>Z00.110</u> Health supervision for newborn under 8 days old
		99391	\$99.99	<u>Z00.111</u> Health supervision for newborn 8 to 28 days old
12 months up to 3 years	EP HA	99382	\$112.38	<u>Z00.121</u> Routine child health exam <i>with abnormal findings</i>
		99392	\$106.92	<u>Z00.129</u> Routine child health exam <i>without abnormal findings</i>
Z02 – Z02.89 (age 0-3 years)				

Rev. 04/14
Rev. 07/14
Rev. 10/15
Rev. 01/16
Rev. 07/16
Rev. 10/16
Rev. 04/18
Rev. 04/24

TABLE C

Interperiodic EPSDT Visit Procedure Codes

Per the Federal EPSDT policy guidelines, interperiodic screening, vision, hearing, and dental services which are medically necessary to determine the existence of suspected physical or mental illnesses or conditions are to be provided.

Rev. 07/11
Rev. 04/12
Rev. 01/14
Rev. 10/14
Rev. 01/15
Rev. 07/16

Use Table C-1 for Medicaid-eligible and PeachCare for Kids®-eligible children who are up to date on their periodic exams but have been referred because of a suspected problem to a qualified health provider by a health, developmental, or educational professional who comes into contact with the child outside of the formal health care system or a need identified by the provider or parent. The complete set of EPSDT preventive health visit components does not have to be performed. EPSDT providers must document the correct level of care when using office visit codes. The rate differential for PCK’s members are shown in Table C-1.

Rev. 01/21
Rev. 04/24

Code the EPSDT interperiodic visit (99202-99205, 99211-99215) with the EP modifier. When vaccines are administered during the interperiodic visit, code the EPSDT interperiodic visit (99202-99205, 99212-99215) with the EP and 25 modifiers. The National Correct Coding Initiative (NCCI) does not allow reimbursement of the 99211 code when it is billed together with any of the vaccine administration codes (90460, 90471-90474).

Table C-1

Rev. 07/11
Rev. 10/11
Rev. 04/12
Rev. 07/12
Rev. 10/14
Rev. 07/16
Rev. 04/18
Rev. 01/21
Rev. 04/24

HIPAA Procedure Code	HIPAA Modifier	HIPAA Diagnosis Code	Medicaid Fee For Service (FFS) Reimbursement
99202 (new patient)	EP	*	\$72.04
99203 (new patient)	EP	*	\$109.35
99204 (new patient)	EP	*	\$167.09
99205 (new patient)	EP	*	\$202.13
99211 (established patient)	EP	*	\$19.16 \$19.99 PCK®
99212 (established patient)	EP	*	\$46.19
99213 (established patient)	EP	*	\$76.15
99214 (established patient)	EP	*	\$110.43
99215 (established patient)	EP	*	\$148.33

*Use the appropriate HIPAA diagnosis code that relates to the medical service(s) provided.

Table C

Interperiodic Vision and Hearing Screening

Rev. 07/11
Rev. 10/11
Rev. 04/14

An enrolled provider may use the codes as indicated below in Table C-2 when billing for vision and/or hearing screening only. For example, a recheck of a failed hearing screening or a child who needs Form 3300 (Certificate of Vision, Hearing, Dental and Nutrition Screening) completed. Separate reimbursement is not allowed when these screenings are performed during the periodic preventive visit. Please see Table C-2 below.

Table C-2

Rev. 04/14
Rev. 07/14
Rev. 10/15

HIPAA Procedure Code	HIPAA Modifier	Interperiodic Screening	ICD-10 Diagnosis Code	PCK and FFS Rate
99173	EP	Interperiodic Vision	Z01.00 or Z01.01 Or appropriate abnormal results code	\$5.62
V5008, 92551-92553, 92555-92556	EP	Interperiodic Hearing	Z01.10 or Z01.110 Or Z01.118 Or appropriate abnormal results code	\$5.62

Rev. 01/18

Note: The Georgia Department of Public Health’s Form 3300 has been revised to document a nutrition screening. More information and a copy of the form can be found at: http://www.gachd.org/DPH_Form_3300.pdf

Rev. 04/11

The determination of whether an interperiodic screen is medically necessary may be made by a health, developmental, or educational professional who comes into contact with the child outside of the formal health care system [e.g., State early intervention or special education programs, Head Start and day care programs, the Special Supplemental Food Program for Women, Infants and Children (WIC), and other nutritional assistance programs]. Providers may not bill interperiodic and periodic visits on the same day.

905. Immunizations

905.1 Recommended Immunization Schedule:

The Recommended Childhood Immunization Schedule should be used as the guideline for administering immunizations.

905.2 Delayed Immunizations:

Practitioners who begin the immunization process on children who are late or at times other than the recommended optimal immunization schedule may use recommendations from the Advisory Committee on Immunization Practices (ACIP) - see schedule for Children and Adolescents Who Start Late.

905.3 Vaccines for Children (VFC) Program:

It is recommended that all Health Check providers enroll in the Vaccines For Children program to provide immunizations to Medicaid eligible children whose ages are birth through eighteen (18) years of age. If the Health Check provider giving the EPSDT preventive health exam does not wish to participate in VFC, it is expected that they administer vaccines at the time of service and understand that only the administration fee will be reimbursed. The Vaccines For Children (VFC) program is a federally funded and state operated vaccine supply program that began October 1, 1994. The program supplies, at no cost to all public health and private health care providers, federally purchased vaccines to be administered to children in certain groups. Children eligible to receive VFC-provided vaccines include the following:

- A. children enrolled in Medicaid;
- B. children who do not have health insurance;
- C. children who are American Indian or Alaskan native; and
- D. children who have health insurance but for whom vaccines are not a covered benefit.

Note: PeachCare for Kids® members may receive state purchased vaccines.

Questions regarding enrollment and vaccine orders should be directed to the appropriate VFC program (1-800-848-3868).

Since vaccines are provided at no cost to the Health Check provider for children eighteen (18) years and younger, only administration costs are allowed to be submitted for reimbursement for vaccines administered to this age group.

Rev. 04/09
Rev. 10/14

Rev. 01/10
Rev. 04/12
Rev. 03/13
Rev. 10/14

905.4 Tdap and Meningococcal Requirements

7th Grade Immunization Requirements

Rev. 07/14
Rev. 04/20

Beginning in the 2014-2015 academic school year (effective July 1, 2014), the Georgia Department of Public Health (DPH) Rule (511-2-2) requires all students born on or after January 1, 2002 entering or transferring into seventh (7th) grade and any “new entrant” entering into 8th-12th grades in Georgia must provide proof that the student has received one dose of Tdap (tetanus, diphtheria, pertussis) vaccine and one dose of meningococcal conjugate (MCV4) vaccine. DPH guidelines state that the student must have received the meningococcal vaccine on or after their 10th birthday for entry into the 7th grade. A student that received the meningococcal vaccine before their 10th birthday will need to be revaccinated on or after his/her 10th birthday. There is a minimum time interval of 8 weeks between the previous dose of the MCV4 vaccine and the newly required dose.

This law affects all public and private schools including, but not limited to, charter schools, community schools, juvenile court schools and other alternative school settings (excluding homeschool). “New Entrant” means any child entering any school in Georgia for the first time or entering after having been absent from a Georgia school for more than twelve months or one school year.

11th Grade Immunization Requirements

Rev. 04/20

Georgia's immunization requirements for students entering or transferring into the eleventh grade have been revised to align with the current recommendations of the Advisory Committee on Immunization Practices (ACIP), the American Academy of Pediatrics (AAP), and the American Academy of Family Physicians (AAFP).

Beginning in the 2020-2021 academic school year (effective July 1, 2020), all students who are new entrants or transfers into a Georgia school in the eleventh grade, will require proof of a booster dose of the meningococcal conjugate vaccine, unless their first dose was received on or after their sixteenth birthday. Georgia law requires students be vaccinated against this disease, unless the child has an exemption.

<https://dph.georgia.gov/immunization-section>

Recommended Immunization Schedule for Children and Adolescents Aged 18 Years or Younger, United States, 2024

(For those who fall behind or start late, see the catch-up schedule)

Rev. 04/16
 Rev. 04/17
 Rev. 04/18
 Rev. 04/19
 Rev. 04/20
 Rev. 04/21
 Rev. 04/22
 Rev. 04/23
 Rev. 01/24

Table 1 Recommended Child and Adolescent Immunization Schedule for Ages 18 Years or Younger, United States, 2024

These recommendations must be read with the notes that follow. For those who fall behind or start late, provide catch-up vaccination at the earliest opportunity as indicated by the green bars. To determine minimum intervals between doses, see the catch-up schedule (Table 2).

Vaccine and other immunizing agents	Birth	1 mo	2 mos	4 mos	6 mos	9 mos	12 mos	15 mos	18 mos	19-23 mos	2-3 yrs	4-6 yrs	7-10 yrs	11-12 yrs	13-15 yrs	16 yrs	17-18 yrs	
Respiratory syncytial virus (RSV-mAb [Nirsevimab])	1 dose depending on maternal RSV vaccination status, See Notes					1 dose (8 through 19 months), See Notes												
Hepatitis B (HepB)	1 st dose	← 2 nd dose →		← 3 rd dose →														
Rotavirus (RV): RV1 (2-dose series), RVS (3-dose series)				1 st dose	2 nd dose	See Notes												
Diphtheria, tetanus, acellular pertussis (DTaP <7 yrs)				1 st dose	2 nd dose	3 rd dose	← 4 th dose →						5 th dose					
Haemophilus influenzae type b (Hib)				1 st dose	2 nd dose	See Notes		← 3 rd or 4 th dose → See Notes										
Pneumococcal conjugate (PCV15, PCV20)				1 st dose	2 nd dose	3 rd dose	← 4 th dose →											
Inactivated poliovirus (IPV <18 yrs)				1 st dose	2 nd dose	← 3 rd dose →					4 th dose		See Notes					
COVID-19 (1vCOV-mRNA, 1vCOV-aPS)	1 or more doses of updated (2023-2024 Formula) vaccine (See Notes)																	
Influenza (IIV4)	Annual vaccination 1 or 2 doses																	
Influenza (LAIV4)											Annual vaccination 1 or 2 doses		Annual vaccination 1 dose only					
Measles, mumps, rubella (MMR)						See Notes		← 1 st dose →					2 nd dose					
Varicella (VAR)							← 1 st dose →					2 nd dose						
Hepatitis A (HepA)						See Notes		2-dose series, See Notes										
Tetanus, diphtheria, acellular pertussis (Tdap ≥7 yrs)														1 dose				
Human papillomavirus (HPV)															See Notes			
Meningococcal (MenACWY-CRM ≥2 mos, MenACWY-TT ≥2years)				See Notes											1 st dose	2 nd dose		
Meningococcal B (MenB-4C, MenB-FHbp)																See Notes		
Respiratory syncytial virus vaccine (RSV [Abrysvo])														Seasonal administration during pregnancy, See Notes				
Dengue (DEN4CYD; 9-16 yrs)														Seropositive in endemic dengue areas (See Notes)				
Mpox																		

Range of recommended ages for all children
 Range of recommended ages for catch-up vaccination
 Range of recommended ages for certain high-risk groups
 Recommended vaccination can begin in this age group
 Recommended vaccination based on shared clinical decision-making
 No recommendation/not applicable

NOTE: The above recommendations must be read along with the footnotes of this schedule
 Refer to footnotes at <https://www.cdc.gov/vaccines/schedules/downloads/child/0-18yrs-child-combined-schedule.pdf>

Catch-up Immunization Schedule for Children and Adolescents who start late or who are more than 1-month behind, United States, 2024

Rev. 04/16
 Rev. 04/17
 Rev. 04/18
 Rev. 04/19
 Rev. 04/20
 Rev. 04/21
 Rev. 04/22
 Rev. 04/23
 Rev. 01/24

Table 2 Recommended Catch-up Immunization Schedule for Children and Adolescents Who Start Late or Who Are More than 1 Month Behind, United States, 2024					
The table below provides catch-up schedules and minimum intervals between doses for children whose vaccinations have been delayed. A vaccine series does not need to be restarted, regardless of the time that has elapsed between doses. Use the section appropriate for the child's age. Always use this table in conjunction with Table 1 and the Notes that follow.					
Children age 4 months through 6 years					
Vaccine	Minimum Age for Dose 1	Minimum Interval Between Doses			
		Dose 1 to Dose 2	Dose 2 to Dose 3	Dose 3 to Dose 4	Dose 4 to Dose 5
Hepatitis B	Birth	4 weeks	8 weeks and at least 16 weeks after first dose minimum age for the final dose is 24 weeks		
Rotavirus	6 weeks Maximum age for first dose is 14 weeks, 6 days.	4 weeks	4 weeks maximum age for final dose is 8 months, 0 days		
Diphtheria, tetanus, and acellular pertussis	6 weeks	4 weeks	4 weeks	6 months	6 months A fifth dose is not necessary if the fourth dose was administered at age 4 years or older and at least 6 months after dose 3
Haemophilus influenzae type b	6 weeks	No further doses needed if first dose was administered at age 15 months or older. 4 weeks if first dose was administered before the 1 st birthday. 8 weeks (as final dose) if first dose was administered at age 12 through 14 months.	No further doses needed if previous dose was administered at age 15 months or older 4 weeks if current age is younger than 12 months and first dose was administered at younger than age 7 months and at least 1 previous dose was PRP-T (ActHib®, Pentacef®, Hibrix®), Vaxelis® or unknown 8 weeks and age 12 through 59 months (as final dose) if current age is younger than 12 months and first dose was administered at age 7 through 11 months; OR if current age is 12 through 59 months and first dose was administered before the 1 st birthday and second dose was administered at younger than 15 months; OR if both doses were PedvaxHIB® and were administered before the 1st birthday	8 weeks (as final dose) This dose only necessary for children age 12 through 59 months who received 3 doses before the 1 st birthday.	
Pneumococcal conjugate	6 weeks	No further doses needed for healthy children if first dose was administered at age 24 months or older 4 weeks if first dose was administered before the 1 st birthday 8 weeks (as final dose for healthy children) if first dose was administered at the 1 st birthday or after	No further doses needed for healthy children if previous dose was administered at age 24 months or older 4 weeks if current age is younger than 12 months and previous dose was administered at <7 months old 8 weeks (as final dose for healthy children) if previous dose was administered between 7-11 months (wait until at least 12 months old); OR if current age is 12 months or older and at least 1 dose was administered before age 12 months	8 weeks (as final dose) This dose is only necessary for children age 12 through 59 months regardless of risk, or age 60 through 71 months with any risk, who received 3 doses before age 12 months.	
Inactivated poliovirus	6 weeks	4 weeks	4 weeks if current age is <4 years 6 months (as final dose) if current age is 4 years or older	6 months (minimum age 4 years for final dose)	
Measles, mumps, rubella	12 months	4 weeks			
Varicella	12 months	3 months			
Hepatitis A	12 months	6 months			
Meningococcal ACWY	2 months MenACWY-CRM 2 years MenACWY-TT	8 weeks	See Notes	See Notes	
Children and adolescents age 7 through 18 years					
Meningococcal ACWY	Not applicable (N/A)	8 weeks			
Tetanus, diphtheria, tetanus, diphtheria, and acellular pertussis	7 years	4 weeks	4 weeks if first dose of DTaP/DT was administered before the 1 st birthday 6 months (as final dose) if first dose of DTaP/DT or Tdap/Td was administered at or after the 1 st birthday	6 months if first dose of DTaP/DT was administered before the 1 st birthday	
Human papillomavirus	9 years	Routine dosing intervals are recommended.			
Hepatitis A	N/A	6 months			
Hepatitis B	N/A	4 weeks	8 weeks and at least 16 weeks after first dose		
Inactivated poliovirus	N/A	4 weeks	6 months A fourth dose is not necessary if the third dose was administered at age 4 years or older and at least 6 months after the previous dose.	A fourth dose of IPV is indicated if all previous doses were administered at <4 years OR if the third dose was administered <6 months after the second dose.	
Measles, mumps, rubella	N/A	4 weeks			
Varicella	N/A	3 months if younger than age 13 years. 4 weeks if age 13 years or older			
Dengue	9 years	6 months	6 months		

NOTE: The above recommendations must be read along with the footnotes of this schedule
 Refer to footnotes at <https://www.cdc.gov/vaccines/schedules/downloads/child/0-18yrs-child-combined-schedule.pdf>

Recommended Child and Adolescent Immunization Schedule by Medical Indication, United States, 2024

Rev. 04/16
 Rev. 04/17
 Rev. 04/18
 Rev. 04/19
 Rev. 04/20
 Rev. 04/21
 Rev. 04/22
 Rev. 04/23
 Rev. 01/24

Table 3 Recommended Child and Adolescent Immunization Schedule by Medical Indication, United States, 2024

Always use this table in conjunction with Table 1 and the Notes that follow. Medical conditions are often not mutually exclusive. If multiple conditions are present, refer to guidance in all relevant columns. See Notes for medical conditions not listed.

Vaccine and other immunizing agents	Pregnancy	Immunocompromised (excluding HIV infection)	HIV infection CD4 percentage and count ^a		CSF leak or cochlear implant	Asplenia or persistent complement deficiencies	Heart disease or chronic lung disease	Kidney failure, End-stage renal disease or on Dialysis	Chronic liver disease	Diabetes
			<15% or <200mm	≥15% and ≥200mm						
RSV-mAb (nirsevimab)		2nd RSV season	1 dose depending on maternal RSV vaccination status, See Notes				2nd RSV season for chronic lung disease (See Notes)			1 dose depending on maternal RSV vaccination status, See Notes
Hepatitis B										
Rotavirus		SCID ^b								
DTaP/Tdap	DTaP Tdap: 1 dose each pregnancy									
Hib		HSCT: 3 doses	See Notes			See Notes				
Pneumococcal										
IPV										
COVID-19			See Notes							
IIV4										
LAIV4							Asthma, wheezing: 2–4 years ^c			
MMR	*									
VAR	*									
Hepatitis A										
HPV	*	3 dose series. See Notes								
MenACWY										
MenB										
RSV (Abrysvo)	Seasonal administration, See Notes									
Dengue										
Mpox	See Notes									

 Recommended for all age-eligible children who lack documentation of a complete vaccination series
 Not recommended for all children, but is recommended for some children based on increased risk for or severe outcomes from disease
 Recommended for all age-eligible children, and additional doses may be necessary based on medical condition or other indications. See Notes.
 Precaution: Might be indicated if benefit of protection outweighs risk of adverse reaction
 Contraindicated or not recommended
 *Vaccinate after pregnancy, if indicated
 No Guidance/Not Applicable

a. For additional information regarding HIV laboratory parameters and use of live vaccines, see the General Best Practice Guidelines for Immunization, "Altered Immunocompetence," at www.cdc.gov/vaccines/hcp/acip-recs/general-recs/immunocompetence.html and Table 4-1 (footnote J) at www.cdc.gov/vaccines/hcp/acip-recs/general-recs/contraindications.html.
b. Severe Combined Immunodeficiency
c. LAIV4 contraindicated for children 2–4 years of age with asthma or wheezing during the preceding 12 months

Recommended Adult Immunization Schedule by Age Group, United States, 2024

Rev. 04/16
 Rev. 04/17
 Rev. 04/18
 Rev. 04/19
 Rev. 04/20
 Rev. 04/21
 Rev. 04/22
 Rev. 04/23
 Rev. 01/24

Table 1 Recommended Adult Immunization Schedule by Age Group, United States, 2024

Vaccine	19–26 years	27–49 years	50–64 years	≥65 years
COVID-19	1 or more doses of updated (2023-2024 Formula) vaccine (See Notes)			
Influenza inactivated (IIV4) or Influenza recombinant (RIV4)	1 dose annually			
Influenza live, attenuated (LAIV4)	1 dose annually			
Respiratory Syncytial Virus (RSV)	Seasonal administration during pregnancy. See Notes.			≥60 years
Tetanus, diphtheria, pertussis (Tdap or Td)	1 dose Tdap each pregnancy; 1 dose Td/Tdap for wound management (see notes)			
	1 dose Tdap, then Td or Tdap booster every 10 years			
Measles, mumps, rubella (MMR)	1 or 2 doses depending on indication (if born in 1957 or later)			For healthcare personnel, see notes
Varicella (VAR)	2 doses (if born in 1980 or later)	2 doses		
Zoster recombinant (RZV)	2 doses for immunocompromising conditions (see notes)		2 doses	
Human papillomavirus (HPV)	2 or 3 doses depending on age at initial vaccination or condition	27 through 45 years		
Pneumococcal (PCV15, PCV20, PPSV23)				See Notes
				See Notes
Hepatitis A (HepA)	2, 3, or 4 doses depending on vaccine			
Hepatitis B (HepB)	2, 3, or 4 doses depending on vaccine or condition			
Meningococcal A, C, W, Y (MenACWY)	1 or 2 doses depending on indication, see notes for booster recommendations			
Meningococcal B (MenB)	19 through 23 years	2 or 3 doses depending on vaccine and indication, see notes for booster recommendations		
Haemophilus influenzae type b (Hib)	1 or 3 doses depending on indication			
Mpox				

Recommended vaccination for adults who meet age requirement, lack documentation of vaccination, or lack evidence of immunity
 Recommended vaccination for adults with an additional risk factor or another indication
 Recommended vaccination based on shared clinical decision-making
 No recommendation/Not applicable

The above recommendations are located at <https://www.cdc.gov/vaccines/schedules/downloads/adult/adult-combined-schedule.pdf>

Recommended Adult Immunization Schedule by Medical Condition and Other Indications, 2024

Rev. 04/16
 Rev. 04/17
 Rev. 04/18
 Rev. 04/19
 Rev. 04/20
 Rev. 04/21
 Rev. 04/22
 Rev. 04/23
 Rev. 01/24

Table 2 Recommended Adult Immunization Schedule by Medical Condition or Other Indication, United States, 2024

Always use this table in conjunction with Table 1 and the Notes that follow. Medical conditions or indications are often not mutually exclusive. If multiple medical conditions or indications are present, refer to guidance in all relevant columns. See Notes for medical conditions or indications not listed.

VACCINE	Pregnancy	Immunocompromised (excluding HIV infection)	HIV infection CD4 percentage and count		Men who have sex with men	Asplenia, complement deficiency	Heart or lung disease	Kidney failure, End-stage renal disease or on dialysis	Chronic liver disease; alcoholism*	Diabetes	Healthcare Personnel ^c
			<15% or <200mm	≥15% and ≥200mm							
COVID-19		See Notes									
IIV4 or RIV4	1 dose annually										
LAIV4					1 dose annually if age 19 - 49 years		1 dose annually if age 19 - 49 years				
RSV	Seasonal administration. See Notes	See Notes					See Notes				
Tdap or Td	Tdap: 1 dose each pregnancy	1 dose Tdap, then Td or Tdap booster every 10 years									
MMR	*										
VAR	*	See Notes									
RZV		See Notes									
HPV	*	3 dose series if indicated									
Pneumococcal											
HepA											
Hep B	See Notes								Age ≥ 60 years		
MenACWY											
MenB											
Hib		HSCT: 3 doses ^a					Asplenia: 1 dose				
Mpox	See Notes				See Notes						

 Recommended for all adults who lack documentation of vaccination, OR lack evidence of immunity
 Not recommended for all adults, but recommended for some adults based on either age OR increased risk for or severe outcomes from disease
 Recommended based on shared clinical decision-making
 Recommended for all adults, and additional doses may be necessary based on medical condition or other indications. See Notes.
 Precaution: Might be indicated if benefit of protection outweighs risk of adverse reaction
 Contraindicated or not recommended. *Vaccinate after pregnancy, if indicated
 No Guidance/ Not Applicable

a. Precaution for LAIV4 does not apply to alcoholism.
 b. See notes for influenza; hepatitis B; measles, mumps, and rubella; and varicella vaccinations.
 c. Hematopoietic stem cell transplant.

The above recommendations are located at <https://www.cdc.gov/vaccines/schedules/downloads/adult/adult-combined-schedule.pdf>

906. Diagnostic, Treatment and Referral Services

Rev. 07/10
Rev. 10/14

All suspicious or abnormal findings identified during an EPSDT preventive health visit must be treated or be further evaluated.

When an EPSDT service is needed but not performed during the EPSDT preventive health visit, the child provider should be appropriately referred for diagnosis.

Rev. 04/12
Rev. 10/14

For non-CMO (care management organization) Medicaid and PeachCare for Kids[®] Fee-For Service (FFS) members, the screening provider must either treat (if qualified) or refer all members with abnormal findings. Members needing referrals must be appropriately referred to Medicaid or PeachCare for Kids[®] enrolled providers. For information on billing levels allowed for treatment during the EPSDT periodic and interperiodic visits, see section 1003. Billing Tips.

If the provider is not the member's EPSDT PCP, the provider must notify the member's PCP of the preventive health/interperiodic visit to discuss any clinical findings which require prompt medical attention. Referral and prior authorization may be required for children who are assigned a PCP.

Rev. 01/10
Rev. 10/14

The level of treatment required should determine whether additional services are billed or provided during the EPSDT scheduled visit.

Rev. 07/10

907. Lead Risk Assessment and Screening

907.1 Purpose:

The purpose of screening for lead absorption is to identify children who have either symptomatic or asymptomatic lead poisoning and to intervene as quickly as possible to reduce their blood lead levels.

907.2 Lead Screening:

Rev. 07/08
Rev. 07/10
Rev. 10/14

- A. Since 1989, Federal law has required that children enrolled in Medicaid must have their blood lead level measured at 12 and 24 months of age.
- B. A blood lead test, *capillary* or *venous*, must be used when screening Medicaid-eligible children. A capillary Blood Lead Test that is elevated ($\geq 3.5\mu\text{g}/\text{dL}$ reported by a certified lab) must be confirmed with a repeat Blood Lead Test (confirmatory venous specimen is preferred) at a certified laboratory.
- C. ALL venous sample lead screening tests conducted using any Magellan Diagnostic lead testing system should be laboratory analyzed by a properly accredited laboratory.

Rev. 07/17

907.3 Lead Health Education and Anticipatory Guidance

Health education is a required component of screening services (EPSDT benefit in accordance with section 1905(r) of the Act) and includes anticipatory guidance.

Anticipatory Guidance regarding Lead Exposure

Anticipatory guidance should be provided to families when children are:

- A. 3-6 months of age and again at 12 months.
- B. Between the ages of 24 and 72 months at well-child visits and when a lead risk assessment questionnaire is administered.

The following topics should be covered with anticipatory guidance:

- A. Effects of lead poisoning on children
- B. Sources of lead poisoning
- C. Pathways of exposure (including placental exposure)
- D. How to prevent a child's exposure to lead hazards
- E. Appropriate schedule for testing children for lead poisoning

907.4 Lead Case Management

All children whose initial screening test shows an elevated blood lead level should follow the Georgia Healthy Homes and Lead Poisoning Prevention Program's (GHHLPPP) Case Management Guidelines. Georgia Department of Public Health (DPH) Lead Hazard Risk Assessors, under the guidance of the GHHLPPP, will perform an environmental lead risk assessment for all children with a confirmed blood lead level of ≥ 3.5 ug/dL, as well as provide education about low cost methods to reduce identified lead hazards. As a primary care provider, you will be notified by a DPH Lead Hazard Risk Assessor of the results of the environmental lead risk assessment and remediation recommendations.

Refer to the GHHLPPP Case Management Guidelines located in Appendix A, Guidelines in Screening and Reporting Elevated Blood Lead Levels.

Rev. 10/12
Rev. 04/14
Rev. 04/23

908. Oral Health and Dental Services

Rev. 10/14

Per the Federal EPSDT policy guidelines, dental services must be provided to eligible members under twenty-one years of age. Dental services required under the EPSDT benefit include:

- Dental care needed for relief of pain, infection, restoration of teeth, and maintenance of dental health (provided at as early an age as necessary); and
- Emergency, preventive, and therapeutic services for dental disease that, if left untreated, may become acute dental problems or cause irreversible damage to the teeth or supporting structures.

In addition, medically necessary oral health and dental services, including those identified during an oral screening or a dental exam, are covered under the EPSDT benefit.


Refer to the Dental Services Program Policy and Procedures Manual.

Rev. 07/12
Rev. 10/19
Rev. 07/23

In accordance with the American Academy of Pediatric Dentistry (AAPD) – [Oral Health Policies & Recommendations \(The Reference Manual of Pediatric Dentistry\)](#), dental providers should refer to the [Periodicity of Examination, Preventive Dental Services, Anticipatory Guidance/Counseling, and Oral Treatment for Infants, Children, and Adolescents for best practice recommendations about anticipatory guidance and timing of other clinical modalities which promote oral health during infancy, childhood, and adolescence](#). The AAPD intends these recommendations to help practitioners make clinical decisions concerning preventive oral health interventions, including anticipatory guidance and preventive counseling, for infants, children, and adolescents. The [Recommended Dental Periodicity Schedule for Pediatric Oral Health Assessment, Preventive Services, and Anticipatory Guidance/Counseling](#) may be assessed at https://www.aapd.org/globalassets/media/policies_guidelines/bp_recdentperiodschedule.pdf

Recommended Dental Periodicity Schedule for Pediatric Oral Health Assessment, Preventive Services, and Anticipatory Guidance/Counseling

Since each child is unique, these recommendations are designed for the care of children who have no contributing medical conditions and are developing normally. These recommendations will need to be modified for children with special health care needs or if disease or trauma manifests variations from normal. The American Academy of Pediatric Dentistry emphasizes the importance of very early professional intervention and the continuity of care based on the individualized needs of the child. Refer to the text of this best practice for supporting information and references.

 AMERICA'S PEDIATRIC DENTISTS THE BIG AUTHORITY on little teeth[®]	AGE				
	6 TO 12 MONTHS	12 TO 24 MONTHS	2 TO 6 YEARS	6 TO 12 YEARS	12 YEARS AND OLDER
Clinical oral examination ¹	•	•	•	•	•
Assess oral growth and development ²	•	•	•	•	•
Caries-risk assessment ³	•	•	•	•	•
Radiographic assessment ⁴	•	•	•	•	•
Prophylaxis and topical fluoride ^{3,4}	•	•	•	•	•
Fluoride supplementation ⁵	•	•	•	•	•
Anticipatory guidance/counseling ⁶	•	•	•	•	•
Oral hygiene counseling ^{3,7}	Parent	Parent	Patient/parent	Patient/parent	Patient
Dietary counseling ^{3,8}	•	•	•	•	•
Counseling for nonnutritive habits ⁹	•	•	•	•	•
Injury prevention and safety counseling ¹⁰	•	•	•	•	•
Assess speech/language development ¹¹	•	•	•		
Assessment developing occlusion ¹²			•	•	•
Assessment for pit and fissure sealants ¹³			•	•	•
Periodontal-risk assessment ^{3,14}			•	•	•
Counseling for tobacco, vaping, and substance misuse				•	•
Counseling for human papilloma virus/vaccine				•	•
Counseling for intraoral/perioral piercing				•	•
Assess third molars					•
Transition to adult dental care					•

1 First examination at the eruption of the first tooth and no later than 12 months. Repeat every six months or as indicated by child's risk status/susceptibility to disease. Includes assessment of pathology and injuries.
 2 By clinical examination.
 3 Must be repeated regularly and frequently to maximize effectiveness.
 4 Timing, types, and frequency determined by child's history, clinical findings, and susceptibility to oral disease.
 5 Consider when systemic fluoride exposure is suboptimal. Up to at least 16 years.
 6 Appropriate discussion and counseling should be an integral part of each visit for care.
 7 Initially, responsibility of parent; as child matures, jointly with parent; then, when indicated, only child.
 8 At every appointment, initially discuss appropriate feeding practices, then the role of refined carbohydrates and frequency of snacking in caries development and childhood obesity. Monitor body mass index beginning at age two.
 9 At first, discuss the need for nonnutritive sucking; digits vs. pacifiers; then the need to wean from the habit before malocclusion or deleterious effect on the dentofacial complex occurs. For school-aged children and adolescent patients, counsel regarding any existing habits such as fingernail biting, clenching, or bruxism.

10 Initially pacifiers, car seats, play objects, electric cords; secondhand smoke; when learning to walk; with sports and routine playing, including the importance of mouthguards; then motor vehicles and high-speed activities.
 11 Observation for age-appropriate speech articulation and fluency as well as achieving receptive and expressive language milestones.
 12 Identify: transverse, vertical, and sagittal growth patterns; asymmetry; occlusal disharmonies; functional status including temporomandibular joint dysfunction; esthetic influences on self-image and emotional development.
 13 For caries-susceptible primary molars, permanent molars, premolars, and anterior teeth with deep pits and fissures; placed as soon as possible after eruption.
 14 Periodontal probing should be added to the risk-assessment process after the eruption of the first permanent molars.

American Academy of Pediatric Dentistry (AAPD) – “The Reference Manual of Pediatric Dentistry”, 2022-2023/p. 165

RECOMMENDED DENTAL PERIODICITY SCHEDULE FOR PEDIATRIC ORAL HEALTH ASSESSMENT, PREVENTIVE SERVICES, AND ANTICIPATORY GUIDANCE/COUNSELING

Footnotes

1. First examination at the eruption of the first tooth and no later than 12 months. Repeat every 6 months or as indicated by child's risk status/susceptibility to disease. Includes assessment of pathology and injuries.
2. By clinical examination.
3. Must be repeated regularly and frequently to maximize effectiveness.
4. Timing, types, and frequency determined by child's history, clinical findings, and susceptibility to oral disease.
5. Consider when systemic fluoride exposure is suboptimal. Up to at least 16 years.
6. Appropriate discussion and counseling should be an integral part of each visit for care.
7. Initially, responsibility of parent; as child matures, jointly with parent; then, when indicated, only child.
8. At every appointment; initially discuss appropriate feeding practices, then the role of refined carbohydrates and frequency of snacking in caries development and childhood obesity. Monitor body mass index beginning at age two.
9. At first, discuss the need for nonnutritive sucking: digits vs pacifiers; then the need to wean from the habit before malocclusion or skeletal dysplasia deleterious effect on the dentofacial complex occurs. For school-aged children and adolescent patients, counsel regarding any existing habits such as fingernail biting, clenching, or bruxism.
10. Initially pacifiers, car seats, play objects, electric cords; secondhand smoke; when learning to walk; with sports and routine playing, including the importance of mouthguards; then motor vehicles and high-speed activities.
11. Observation for age-appropriate speech articulation and fluency as well as achieving receptive and expressive language milestones.
12. Identify: transverse, vertical, and sagittal growth patterns; asymmetry; occlusal disharmonies; functional status including temporomandibular joint dysfunction; esthetic influences on self-image and emotional development.
13. For caries-susceptible primary molars, permanent molars, premolars, and anterior teeth with deep pits and fissures; placed as soon as possible after eruption.
14. Periodontal probing should be added to the risk-assessment process after the eruption of the first permanent molars.

909. Other Related Medicaid Programs (This is not an inclusive list)

See the policies and procedures manual of the related programs for complete information. Provider Manuals relevant to EPSDT providers include, but may not be limited to:

- Advanced Nurse Practitioner Services
- Autism Spectrum Disorder (ASD) Services
- Children's Intervention Services (CIS)
- Children's Intervention School Services (CISS)
- Dental Services
- Diagnostic Screening and Preventive Services (DSPS)
- Durable Medical Equipment (DME) Services
- Federally Qualified Health Center (FQHC) Services
- Georgia Pediatric Program (GAPP)
- Hospice Services

- Hospital Services
- Medicaid Medicaid/PeachCare for Kids® Provider Billing Manuals
- Nurse Midwifery Services
- Orthotic and Prosthetic Services
- Pharmacy Services
- Physician Assistant Services
- Physician Services
- Rural Health Clinic (RHC) Services
- Vision Care Services

Provider Manuals are available for downloading. Contact Gainwell Technologies at 1-800-766-4456 or visit the website at www.mmis.georgia.gov for more information.

Rev. 01/21

910. Summary of Non-covered Services

Non-Covered Services include:

Services provided in a manner inconsistent with the provisions of this manual.

Rev. 10/11
Rev. 07/23

911. EPSDT Profile (Appointment Tracking System)

The purpose of the EPSDT Appointment Tracking System is to track enrolled Fee-For-Service children eligible for services and to assist providers in conducting and documenting outreach and follow-up activities to EPSDT families and children.

The EPSDT Appointment Tracking System fully supports the State's goals of providing appropriate and continuing screening and treatment services to Georgia's children and of preventing more costly health problems by encouraging regular health care.

This system provides immediate access to medical and dental information on EPSDT members through online inquiry and provides a reminder call system at no cost to the EPSDT provider. These capabilities enhance the control and operation of the EPSDT program and allow information gathering to support research and program development.

In collaboration with the monthly EPSDT roster (Periodic Screenings Due Report), the EPSDT Profile (Appointment Tracking System) provides:

1. Member's demographic information in addition to the last dates for Hearing (Interperiodic Hearing), Vision (Interperiodic Vision), EPSDT Medical and Dental screenings.
2. Detailed information on the member's entire EPSDT history. This allows the provider to view the member's entire EPSDT history and document outreach attempts as a result of letters/rosters distributed. Based on the notice type distributed by Gainwell Technologies, all the provider has to do is document the member's response and a response date. For example, if the provider arranges a future appointment with the member, he/she will select scheduled appointment under the drop down box for response type and enter the date of the appointment under response date.

Rev. 01/21

3. The Response Type options on the drop-down box are:
 - a. Set Appointment (EPSDT preventive health screening visit)
 - b. Set Appointment (Dental)
 - c. Set Appointment (Blood Lead).
 - d. Screen Completed
4. The last section of the EPSDT Profile is the critical health information. EPSDT medical and dental providers are encouraged to enter information determined to be useful to another Health Care professional in the delivery of care to the member (For example, allergic to Penicillin).
5. If you need further instructions, feel free to click on the help link.

912. EPSDT HIPAA Referral Codes

The Centers for Medicare and Medicaid Services (CMS) defines an EPSDT referral as a member scheduled for another appointment with the EPSDT Provider or a referral to another provider for further needed diagnostic and treatment services as a result of at least one health problem identified during the EPSDT preventive health visit. Effective with HIPAA implementation, CMS and DCH require documentation of EPSDT Referral Codes when submitting EPSDT Screening Code Claims (See Appendix K for examples). When completing the Health Insurance Claim Form [CMS-1500], the EPSDT Referral Codes must be entered in the shaded area of box 24H. (See Appendix L)

Rev. 10/14
Rev. 04/15
Rev. 07/15

Example 1: If the EPSDT screening is normal, the referral code is NU (No follow up visit needed)

Example 2: If the EPSDT screening indicates the need for further diagnostic and treatment services and a follow-up visit is necessary, use the applicable referral code(s):

- AV Available, Not Used: Patient refused referral
- S2 Under Treatment: Patient is currently under treatment for health problem and has a return appointment.
- ST New Services Requested: Referral to another provider for diagnostic or corrective treatment/scheduled.

913. Access to Mental Health Services

Health Link operates the Georgia Crisis and Access Line (GCAL) through a contract with the Department of Behavioral Health and Developmental Disabilities (DBHDD). To access mental health, addictive disease, and crisis services 24 hours a day, 7 days per week call 1-800-715-4225 (GCAL) or go to www.mygcal.com

Rev. 10/14

914. Services for Foster Care Children

As of March 3, 2014, children, youth, and young adults in state custody, children receiving adoption assistance and a select group of children under the Juvenile Justice system, were transitioned to Medicaid managed care under the Georgia Families 360° Program. Amerigroup Community Care is the single Care Management Organization (CMO) managing this population (see Appendix I). Children in state custody under the Kenny A. Consent Decree are required to have an EPSDT preventive health visit and a dental visit within 10 days of official transition to state custody. Unless otherwise noted, EPSDT services for all other children enrolled in the Georgia Families 360° Program should follow the Bright Futures Periodicity Schedule.

Rev. 07/11
Rev. 07/12
Rev. 04/14
Rev. 07/14
Rev. 10/14
Rev. 10/15

PART II - CHAPTER 1000

Basis for Reimbursement

1001. Fee for Service Reimbursement Methodology

Rev. 10/10
Rev. 07/14

The Division will pay the lower of the lowest price regularly and routinely offered to any segment of the general public for the same service or items on the same date(s) of service, the lowest price charged to other third party payers, or effective with dates of service July 1, 2003, the statewide maximum allowable reimbursement which is 84.645% of Medicare's Resource Based Relative Value Scale (RBRVS) for 2000 for Region IV (Atlanta). All procedure codes recognized and adopted after the 2000 RBRVS are subject to the statewide maximum allowable reimbursement – 84.645% of the Region IV Medicare RBRVS in effect at the time the procedure code was adopted.

1002. Vaccines for Children

Rev. 10/10
Rev. 07/11
Rev. 10/11
Rev. 03/13
Rev. 04/13
Rev. 07/13
Rev. 10/14

Since the Vaccines For Children (VFC) program supplies vaccines to providers at no cost to the provider for children birth through eighteen (18) years who have Medicaid, the Division will reimburse an administration fee only for immunizations given to Medicaid enrolled children of this age group. These fees cover the cost of administering the immunizations as well as any paperwork involved (including an immunization or health certificate). Refer to Appendix C-2 for the reimbursement rates for the administration of vaccines provided by the VFC program for the Medicaid-eligible children birth through eighteen (18) years. Appendix C-2 also includes the reimbursement rate for the administration of state-purchased vaccines for the PeachCare for Kids[®] Fee for Service population, for children birth through age eighteen (18) years.

For members nineteen (19) years of age through twenty (20) years of age, VFC stock is not available. Providers must use their own stock of vaccines for these Medicaid eligible members. Health Check providers may include a vaccine administration code on their EPSDT claims when vaccines are administered to members nineteen (19) years of age through twenty (20) years of age. The Division will reimburse for the vaccine product and for vaccine administration. Billing tips for vaccine administration and the effective date for this change are in Appendix C-1 of this manual and providers are strongly encouraged to follow those tips.

1003. Billing Tips

The following are tips to assist with billing for EPSDT screening services and interperiodic visits.

1. The EPSDT preventive health visit is reimbursed as a package of services. All of the age appropriate EPSDT preventive health visit components (as identified in the Bright Futures periodicity schedule) must be completed for each screening visit and billed under one procedure code except where indicated. All preventive or well-child services, except normal newborn care in the hospital, must be billed under the EPSDT Program following the policies and procedures as outlined in this manual.
2. EPSDT preventive health visits must be referred by or performed by the child's primary care practitioner for those services to be reimbursed. Only one (1) periodic preventive or well-child visit will be reimbursed per member at each age-appropriate interval, as specified in the periodicity schedule (excluding foster care members).
3. Providers must perform the age-appropriate hearing and vision screening in order to be reimbursed for the complete EPSDT preventive health exam. Providers may not refer the child to another provider for hearing and vision screening which is required at the time of the EPSDT preventive health visit.
4. When a visit is found to be medically necessary between periodic visit sequences, the EPSDT provider may be reimbursed by billing the appropriate interperiodic visit procedure code. An interperiodic visit cannot be billed on the same date of service as a complete EPSDT preventive health visit.
5. The Georgia Medicaid program reimburses for many of the Diagnostic and Treatment services under other Medicaid programs.

Rev. 01/10
Rev. 10/10
Rev. 07/13
Rev. 10/14
Rev. 10/15
Rev. 01/16
Rev. 07/17
Rev. 07/18

Rev. 04/11
Rev. 10/14
Rev. 07/15

6. Developmental Screenings

Rev. 04/12
Rev. 07/12
Rev. 07/13
Rev. 04/14
Rev. 07/14
Rev. 04/15
Rev. 10/15
Rev. 01/16
Rev. 10/17
Rev. 04/19

A developmental screening should be performed at the following periodic visits: 9, 18, and 30 months. Providers must bill code 96110 with the EP modifier and the appropriate preventive ICD-10 diagnosis code in order to receive reimbursement for this screening.

Only one (1) developmental screening will be reimbursed at each of these intervals. If the child is not seen at the 9-, 18-, or 30-month visit, a developmental screening should be performed during the catch-up visit for the missed periodic visit. This catch-up developmental screening should be billed, using the EP and HA modifiers with code 96110 and the appropriate preventive ICD-10 code. The provider can only bill one (1) catch-up developmental screening during any one (1) catch-up interval.

ICD-10 Diagnosis Codes (preventive)
Z00.121 or Z00.129
Z02 – Z02.89

Rev. 01/24

Developmental screenings when performed outside of the Bright Futures requirements can also be reported. A developmental screening may be performed, as needed, for members ages 3 years through 6 years of age. Providers must bill code 96110 with the EP,U1 modifiers and the appropriate ICD-10 diagnosis code in order to receive reimbursement for this screening.

7. Autism Screenings

Rev. 07/17
Rev. 10/17
Rev. 10/18
Rev. 04/19

Autism screenings should be performed at the 18- and 24-month periodic visits (or catch-up visits). Providers must bill code 96110 with the EP,UA or EP,UA,HA modifiers and the appropriate ICD-10 diagnosis code in order to receive reimbursement for this screening.

Autism screenings when performed outside of the Bright Futures requirements can also be reported. Providers must bill code 96110 with the EP,UA modifiers and Z13.41 or the applicable ICD-10 diagnosis code to receive reimbursement for the screening.

Rev. 01/22

For guidelines related to Autism Spectrum Disorder, including a list of acceptable Autism Spectrum Disorder Assessment Tools, refer to the Autism Spectrum Disorder policy manual Section 801.

8. **Brief Emotional/Behavioral Assessments**

Periodic Screening Visits

Rev. 01/16
Rev. 04/16
Rev. 07/17
Rev. 07/18
Rev. 04/19

An annual depression screening should be performed for members ages 12 years through 20 years during the EPSDT periodic screening visit. When completed during the periodic visit, the depression screening can be reported as a brief emotional/behavioral assessment. Providers should bill code 96127 with the EP modifier, POS 99 and the appropriate ICD-10 diagnosis code in order to receive reimbursement.

Non-Periodic Screening Visits (Sick Visits)

Brief emotional /behavioral assessments, when performed during non-periodic screening visits, can also be reported for the following:

- depression screenings (outside of the BF requirements); AND
- emotional/behavioral assessments conducted for other conditions, such as ADHD, suicidal risk, anxiety, eating disorders, etc.

Brief emotional /behavioral assessments performed during non-periodic visits should be billed with the E/M office visit code (992xx) and reported with procedure code 96127, the appropriate ICD-10 diagnosis code, along with the EP modifier and POS 99.

Procedure code 96127:

- is reimbursed at the current default rate
- should be listed only once per claim for multiple units
 - units submitted should not exceed acceptable medically unlikely edit (MUE) maximum established by CMS

9. **Patient-Focused Health Risk Assessments**

Rev. 07/18
Rev. 10/18
Rev. 04/19

Patient Focused health risk assessments performed during the EPSDT periodic screening visit in accordance with the Bright Futures requirements may be reported (i.e., tobacco, alcohol, or drug use assessment; anemia risk assessment; lead risk assessment).

Providers should bill procedure code 96160 with the EP modifier, POS 99 and the appropriate ICD-10 diagnosis code in order to receive reimbursement. Code the health risk assessment (96160) with the EP, 59 modifiers when reporting the health risk assessment(s) and vaccine administration codes for the same visit.

Health risk assessments when performed outside of the Bright Futures requirements can also be reported. Health risk assessments performed during non-periodic visits should be reported with procedure code 96160, the appropriate ICD-10 diagnosis code, along with the EP modifier and POS 99. Code the health risk assessment (96160) with the EP, 59 modifiers when reporting the health risk assessment(s) and vaccine administration codes for the same visit.

Procedure Code 96160:

- is reimbursed at the current default rate.
- should be listed only once per claim for multiple units.
- units submitted should not exceed acceptable medically unlikely edit (MUE) maximum established by CMS.

10. Caregiver-Focused Health Risk Assessments

Caregiver focused health risk assessments performed during the EPSDT periodic screening visit in accordance with the Bright Futures requirements may be reported, (i.e., maternal depression screening).

A maternal depression screening should be performed during the following EPSDT periodic screening visits:

- By 1 month
- 2 month
- 4 month
- 6 month

Providers should bill procedure code 96161 with the EP modifier, POS 99 and the appropriate ICD-10 diagnosis code in order to receive reimbursement. Code the caregiver focused health risk assessment (96161) with the EP, 59 modifiers when reporting the health risk assessment(s) and vaccine administration codes for the same visit.

Caregiver focused health risk assessments when performed outside of the Bright Futures requirements can also be reported. Health risk assessments performed during non-periodic visits should be reported with procedure code 96161, the appropriate ICD-10 diagnosis code, along with the EP modifier and POS 99. Code the health risk assessment (96161) with the EP, 59 modifiers when reporting the health risk assessment(s) and vaccine administration codes (90460, 90471-74) for the same visit.

Procedure Code 96161:

- is reimbursed at the current default rate.
- should be listed only once per claim for multiple units.
- units submitted should not exceed acceptable medically unlikely edit (MUE) maximum established by CMS.

11. Hematocrit and Hemoglobin Level

The laboratory tests due at the twelve (12) month visit for hematocrit and hemoglobin levels may be performed as in office tests at the time of the EPSDT preventive health visit by the PCP; or the blood sample may be obtained by the PCP and submitted to a Medicaid contracted lab; or the member may be sent to a Medicaid contracted lab for the blood draw and laboratory analysis. The PCP must document in the medical record which option was selected. These tests cannot be sent to a non-participating laboratory for analysis.

Rev. 07/18
Rev. 10/18
Rev. 04/19

12. Federally required Blood Lead Level (BLL) screening

Rev. 07/09
Rev. 10/09
Rev. 04/11
Rev. 04/14
Rev. 07/14
Rev. 10/14
Rev. 04/15
Rev. 10/15
Rev. 10/21

If FFS EPSDT providers use private laboratories for BLL screening or perform BLL screening using an in-office Lead Analyzer, the EPSDT provider cannot file a claim for reimbursement of the BLL test.

The Georgia Public Health Laboratory provides analysis of blood lead specimen and charges a laboratory fee. Fee for Service providers may submit claims to DCH for this fee if the blood sample is obtained by them during the visit and sent to the GPL for analysis. To ensure accurate reimbursement, FFS providers must submit the CPT code 83655 with modifier EP and 90 or 91 along with the CPT code 36415 or 36416 modifier EP and appropriate ICD-10 diagnosis code (i.e., Z13.88, Z00.121, Z00.129, Z77.011).

Additional details regarding this process are contained in Appendix C-4.

13. Vaccine Administration

Rev. 07/11
Rev. 04/12
Rev. 04/13
Rev. 07/13
Rev. 01/14
Rev. 10/14

In order to receive the administration fee from the Division for administering federally or state purchased vaccines for children 0 – 18 years of age, the child must be a Medicaid or PeachCare for Kids® member. Since the federally or state purchased vaccines are provided at no cost to the provider, the Division will only reimburse an administration fee based upon the Division's maximum allowable rate. (See Chapter 1000, Section 1002). The vaccine's National Drug Code (NDC) is not required to be included on the EPSDT claim for reimbursement for the administration of federally or state purchased vaccines for children 0-18 years of age.

Beginning April 1, 2013, providers should bill any of the following appropriate vaccine administration codes, when administering VFC vaccines, as they apply: 90460, 90471, 90472, 90473, 90474. Additional details regarding the use of the vaccine administration codes are contained in Appendix C-1.

Rev. 04/14
Rev. 07/14
Rev. 10/14

EPSDT providers may bill the EPSDT Program for vaccines administered to members nineteen (19) years of age through twenty (20) years of age. Include the vaccine CPT code, diagnosis code, and NDC, along with the appropriate vaccine administration code(s) [90471, 90472, 90473, 90474] on the EPSDT claim. The Division will reimburse for the vaccine product (providers must use their own stock of vaccines) and for vaccine administration. Additional details regarding the use of the vaccine administration codes and the effective date for this change are contained in Appendix C-1.

Rev. 10/07
Rev. 10/10
Rev. 07/11
Rev. 04/14
Rev. 10/14
Rev. 07/15
Rev. 07/17

14. Office Visit Codes

Providers must use place of service (POS) code 99 when billing office visits for EPSDT preventive health screening services. All diagnostic x-ray, laboratory testing (except hematocrit, hemoglobin) and/or treatment services provided to the EPSDT member at the time of the preventive health visit, can be billed on the same CMS 1500 claim form as the EPSDT preventive health visit if the EPSDT provider uses a CMS 1500 form to bill Diagnostic and Treatment Services (i.e., Physician Services, Nurse Practitioner Services, etc.).

Rev. 01/21

Effective May 1, 2015, paper claims are no longer accepted by Gainwell Technologies. As part of the Georgia Paperless Initiative, providers are required to submit CMS 1500 claims electronically over the GAMMIS web portal. For more information regarding the Paperless Initiative, please access the web portal and review all related Banner Messages.

If an EPSDT provider uses a UB 04 to bill Diagnostic and Treatment services (i.e., Hospitals, Rural Health Clinics, etc.), they may also bill the EPSDT preventive health visit services on the UB 04.

Rev. 10/11
Rev. 04/12
Rev. 10/21
Rev. 07/23

15. School-Based Vaccine Clinics

Public Health providers must use place of service (POS) code 03 when billing the vaccine administration fee for vaccines administered during school-based clinics held within their county of jurisdiction.

Rev. 04/14
Rev. 10/14
Rev. 04/16

16. School-Based Telemedicine Services

LEAs enrolled as Health Check providers to serve as telemedicine originating sites only will be allowed to bill the telemedicine originating site facility fee (procedure code Q3014). The LEA provider should report procedure code Q3014 along with the EP and GT modifiers, POS 03, and the appropriate ICD-10 diagnosis code(s). The diagnosis code(s) should be the same diagnosis code(s) listed on the distant site (rendering) provider's claim.

The rendering provider serving as the telemedicine distant site should report the E/M office visit code (992xx) along with the GT modifier (including any other applicable modifiers), the appropriate POS, and the ICD-10 diagnosis code(s). In order for the originating site (LEA) provider to receive reimbursement for procedure code Q3014, a corresponding paid history claim from the distant site provider must be found in GAMMIS. The distant site provider's claim billed for the same member, same date of service, with an E/M office visit code (992xx), the same ICD-10 diagnosis code(s) and the GT modifier, will confirm that a telemedicine service was rendered. If no record of the E/M claim is found that aligns with the LEA provider's originating site claim, the originating site claim will suspend up to 30 days after submission in search of the E/M claim. If no record of an E/M claim is found within 30 days after submission of the LEA provider's originating site claim, reimbursement to the LEA provider will be denied. It is the responsibility of the LEA provider to contact the provider who rendered the distant site service to determine if the E/M visit was billed.

The telemedicine originating facility fee is reimbursed at the current DEFAULT rate.

17. Consultation Services

Rev. 07/14
Rev. 10/14
Rev. 07/23

Effective July 1, 2014, providers enrolled in the EPSDT Program, may bill for reimbursement of the following office/outpatient consultation codes: 99242, 99243, 99244, 99245. The EP modifier must be added to the applicable code along any other applicable modifiers. A consultation service can be rendered once every three years.

18. Tobacco Cessation Counseling Services

Rev. 10/14
Rev. 04/15

Effective January 1, 2014, the Division began coverage of tobacco cessation counseling services to all Medicaid members. Providers enrolled in the EPSDT Program may bill 99406 and 99407 for the reimbursement of tobacco cessation counseling. The EP modifier must be added to the applicable code.

The tobacco cessation counseling must be rendered in a face-to-face setting with the member. Only two 12-week tobacco cessation treatment periods will be allowed per member per year. The provider must document the services in the member's medical record every 30 days during the 12-week treatment period.

19. Incontinence Supplies

Rev. 01/16
Rev. 01/20

Incontinence supplies are covered for children ages 2 through 20 years who have an underlying medical condition that prevents control of the bowels or bladder. Incontinence supplies are not covered for convenience. Children under the age of 2 years will be considered for coverage on a case-by-case basis. Since incontinence supplies are not covered for members over 20 years of age, or on a general basis for members under 21 years, providers must ensure the following for services to be considered for coverage:

- the item is considered durable medical equipment (DME);
- there is a current order prescribed by a physician; and
- a prior authorization (PA) must be submitted which includes documentation of medical necessity.

The following procedure codes are covered under DME services with unique HCPCS codes but are without Medicare-based or nationally accepted rates (T4521-T4535, T4541, T4544). These HCPCS codes require a PA. Refer to the DME Manual for further guidance.

20. Fluoride Varnish

Rev. 04/16
Rev. 10/16

Once teeth are present, the application of fluoride varnish is required and may be applied every 3-6 months in the primary care or dental office for children between the ages of 6 months and 5 years.

21. Other Procedure Codes

Rev. 04/15

When billing for EPSDT screening services and interperiodic visits, only the procedure codes for those services found in this manual may be reimbursed under the EPSDT Program. Reimbursement for other services billable to Medicaid is covered under the program areas overseeing the delivery of those services.

22. National Correct Coding Initiative (NCCI) Medically Unlikely Edits (MUEs) Limits

Rev. 04/16

Providers are reminded to bill in compliance with the NCCI MUE limit for procedure codes and to check the MUE file, at minimum, on a quarterly basis for updates. Procedure codes submitted with frequencies greater than the allowed MUE will be denied according to the NCCI MUE regulations set by CMS.

23. NCCI Procedure-To-Procedure (PTP) Edits

Rev. 10/18

Providers are reminded to bill in compliance with the NCCI PTP edits. The NCCI PTP edits define pairs of Healthcare Common Procedure Coding System (HCPCS)/ Current Procedural Terminology (CPT) codes that should not be reported together. The purpose of the PTP edits is to prevent improper payments when incorrect code combinations are reported. Procedure codes that should not be reported together will be denied according to the NCCI PTP edits defined by CMS.

24. **Other Reimbursement Rates**

Rev. 04/16
 Rev. 10/16
 Rev. 07/17
 Rev. 07/18
 Rev. 04/22
 Rev. 04/23
 Rev. 07/23
 Rev. 01/24
 Rev. 04/24

<u>Procedure Code & Description</u>	<u>Modifier(s)</u>	<u>Rate</u>
96110 – Developmental Screening	EP or EP HA or EP U1	\$11.77
96110 – Autism Screening	EP UA or EP UA HA	\$11.77
96127 – Brief Emotional/ Behavioral Assessment (depression screening)	EP	\$4.55
96160 – Patient-Focused Health Risk Assessment (i.e., tobacco, alcohol, drug use risk assessment)	EP	\$3.95
96161 – Caregiver-Focused Health Risk Assessment (i.e., maternal depression screening)	EP	\$3.95
Q3014 – Telehealth Originating Site Facility Fee	EP	\$20.52
99401 – Preventive Medicine Counseling	EP	\$28.29
99242 – Patient Office Consultation, typically 30 minutes	EP	\$89.66
99243 – Patient Office Consultation, typically 40 minutes	EP	\$122.60
99244 – Patient Office Consultation, typically 60 minutes	EP	\$182.06
99245 – Patient Office Consultation, typically 80 minutes	EP	\$223.01
99406 – Smoking and Tobacco Use Intermediate Counseling, Greater than 3 minutes up to 10 minutes	EP	\$13.73
99407 – Smoking and Tobacco Use Intensive Counseling, Greater than 10 minutes	EP	\$27.18

Reviewed 03/26/2024.

Questions regarding Medicaid billing should be directed to Georgia Health Partnership (GHP) at 1-800-766-4456 or ‘Contact Us’ at www.mmis.georgia.gov

APPENDIX A

Guidelines in Screening and Reporting Elevated Blood Lead Levels

Rev. 04/12
Rev. 04/14

The mission of the Georgia Healthy Homes and Lead Poisoning Prevention Program (GHHLPPP) is to eliminate childhood lead poisoning in Georgia.

Screening Guidelines: Children

Rev. 04/12
Rev. 07/15
Rev. 07/16
Rev. 01/20

Screening for lead poisoning helps identify children who need interventions to reduce their blood lead levels. Many children who may have been exposed to lead or who are at risk for lead poisoning go without being screened. This makes their chances of being harmed by lead greater. Parents and providers should know when a child should be tested for lead poisoning.

Rev. 04/20
Rev. 07/21
Rev. 04/23



Guidelines in Screening and Reporting Elevated Blood Lead Levels: Lead Screening Requirements and Medical Management Recommendations For children ages 6 to 72 months

GHHLPPP RISK FACTORS ASSESSMENT QUESTIONNAIRE

-- ask these questions at the 6, 9, 18 months and 3, 4, 5, 6 years of age preventive visits --

1. Is your child living in or regularly visiting, or has your child lived in or regularly visited, a house or childcare center built before 1978?
2. Does your child have a sibling or playmate who has or has had lead poisoning?
3. Does your child come in contact with an adult who works in an industry or has a hobby that uses lead (battery factory, steel smelter, stained glass, fishing or hunting)?
4. Has your child spent more than 1 week in South or Central American, Africa or Asia since their last blood test?
5. Does anyone in your family use ethnic or folk remedies, cosmetics or eat candies or use pottery imported from South or Central America, Africa or Asia?

If the answer is YES or UNKNOWN to any of the questions, a blood lead test is necessary!

<https://dph.georgia.gov/environmental-health/healthy-homes-and-lead-poisoning-prevention>

Test the blood of ALL Medicaid children for lead poisoning at 12 and 24 months of age AND children 3 to 6 years of age if never tested *regardless* of their risk factors.

Lead Screening Requirements and Medical Management Recommendations for Children

Test all Medicaid children at 12 and 24 months of age, and children 3 to 6 years of age if never tested *regardless of their risk factors!*
IT IS A FEDERAL REQUIREMENT

Recommended Medical and Case Management Actions

Rev. 10/12
 Rev. 04/14
 Rev. 07/15
 Rev. 10/19
 Rev. 07/21
 Rev. 04/23

Blood Lead Levels (BLL)	Recommended Medical and Case Management Actions							
	Confirmatory Blood Lead Test	Hospitalization	Chelation Therapy (A)	Blood Lead Level Retest	Referrals (B)	History and Physical (C)	Lead Poisoning Education (D)	Reducing Exposure and Absorption(E)
< 3.5 µg/dL	No	No	No	No	No	No	No	No
3.5 - 9 µg/dL	Within ¹ day to 3 months, venous or capillary	No	No	See <i>Retest Chart</i> below	Yes	Yes	YES	YES
10-19 µg/dL	within 1 day to 3 months, venous or capillary	No	No	see <i>Retest Chart</i> below	YES	YES	YES	YES
20-44 µg/dL	within 1 day to 1 Month, venous or capillary	No	No	see <i>Retest Chart</i> below	YES	YES	YES	YES
45-69 µg/dL	within 24- 48 hours, venous preferred	No, if home is lead-safe	YES	see <i>Retest Chart</i> below	YES	YES	YES	YES
70 µg/dL or higher	within 24 hours, venous only	YES MEDICAL EMERGENCY	YES	see <i>Retest Chart</i> below	YES	YES	YES	YES
THERE IS NO SAFE LEVEL OF LEAD IN THE BODY – DAMAGE CAUSED BY LEAD POISONING IS PERMANENT AND IRREVERSIBLE!								

¹ If the child's blood lead level persists between 10-19 ug/dL (2 blood lead tests 3 months apart) proceed according to the level of care for 20-44 µg/dL.



Guidelines in Screening and Reporting Elevated Blood Lead Levels: Lead Screening Requirements and Medical Management Recommendations

For children ages 6 to 72 months

Explanation of Recommended Medical and Case Management Actions

- (A) **Chelation Therapy:** if chelation therapy is indicated, the child should be immediately removed from the hazardous environment until the child's environment is made lead-safe; however, if the home is already lead-safe, the child may remain in the home unless hospitalization is indicated.
- (B) **Referrals:** contact local health department and/or **GHHLPPP** to assist in case management and environmental investigations.
- (C) **History and Physical:** take medical, environmental, and nutritional history, test for anemia and iron deficiency, assess neurological, psychosocial, and language development, screen all siblings under age 6, and evaluate risk of other family members, especially pregnant women.
- (D) **Lead Poisoning Education:** discuss sources of lead, effects of lead, lead-based paint hazards associated with living in a pre-1978 and/or renovating a pre-1978 home. Discuss how lead affects prenatal care and well childcare at ages 3, 6, and 12 months and explain what blood lead levels mean and their significance. Lastly, contact **GHHLPPP** for information.
- (E) **Reducing Exposure and Absorption:** discuss damp cleaning to remove lead dust on surfaces, eliminating access to deteriorating lead paint surfaces, and ensuring regular meals which are low in fat and rich in calcium and iron; contact **GHHLPPP** for materials.

Rev. 10/12
Rev. 04/14
Rev. 07/15
Rev. 10/19

ELEVATED BLOOD LEAD RE-TEST CHART

Use this chart to determine when to retest children who are *confirmed as lead-poisoned*.
Venous testing is **strongly preferred**, but capillary testing is acceptable.

Rev. 10/12
Rev. 04/14
Rev. 07/15
Rev. 10/19
Rev. 04/23
Rev. 07/23

If the child's last confirmed BLL was...	and...	
	if the child's blood lead level HAS NOT DROPPED at least 3 µg/dl over a span of at least 3 months...	if the child's blood lead level HAS DROPPED at least 3 µg/dl over a span of at least 3 months...
then test the child again in...		
3.5-9 µg/dL	3 months	6 months
10-19 µg/dL	3 months	3 months
20-24 µg/dL	1 month	2 months
25-44 µg/dL	1 month	1 month
45-69 µg/dL	1 month after chelation	1 month after chelation
≥70 µg/dL	1 month after chelation	1 month after chelation
Retesting should occur until the blood lead level is less than 3.5 µg/dL for six months, all lead hazards have been removed, housing is made lead-safe, and no new exposure exists.		

**Department of Public Health, 200 Piedmont Avenue, SE, Atlanta, GA, 30334,
Phone (404) 463-3754 | Fax (404) 463-4039**

Lead Screening Guidelines for Children

Screening for lead poisoning helps identify children who need interventions to reduce their blood lead levels. Many children who may have been exposed to lead or who are at risk for lead poisoning go without being screened. This makes their chances of being harmed by lead greater. Parents and providers should know when a child should be tested for lead poisoning.

Medicaid and PeachCare for Kids

All children enrolled in Medicaid and PeachCare for Kids® should be tested for lead poisoning and offered certain services based on the following schedule:

Age	Lead Blood Test	Lead Risk Assessment Questionnaire	Anticipatory Guidance
6 months		X	X
9 months		X	X
12 months	X (risk assessment if not enrolled in Medicaid)		X
24 months	X (risk assessment if not enrolled in Medicaid)		X
36-72 months	X – If there's no record of previous test at 12 and 24 months	X - Complete annually unless blood lead test performed	X

Source: [Healthy Homes and Lead Poisoning Prevention | Georgia Department of Public Health](#)

Lead Risk Assessment Questionnaire

Rev. 04/12
Rev. 04/14
Rev. 10/14
Rev. 04/23

The GHHLPPP Childhood Lead Risk Questionnaire can be found at:

[Healthy Homes and Lead Poisoning Prevention | Georgia Department of Public Health](#) in [English](#), [Spanish](#) and [Vietnamese](#).

When using the questionnaire, blood lead tests should be done right away if the child is at high risk (one or more "yes" or "I don't know" answers on the lead risk assessment questionnaire) for lead exposure. Completing this questionnaire does not count as a lead screening.

Blood Lead Test

Rev. 04/12
Rev. 07/15
Rev. 10/19

A blood test is the preferred method for lead screening. There are two tests used to obtain blood lead specimens: capillary blood test or venous blood test. Finger stick capillary blood tests (the Lead Care II Analyzer uses capillary blood) can be done as the initial screening. However, a lab analyzed sample is necessary for confirmation. Safety measures should be taken to reduce the risk of contamination of the capillary blood sample. These measures include:

- Rinsing powder from the examination gloves
- Thoroughly washing patient's hands with soap and water, then drying them before taking a sample.

A venous blood test can be done as the initial screening as well. This method should always be used to confirm elevated blood lead test results when a capillary test was used as the initial screening. Alternatively, a second lab analyzed capillary test can be used to confirm an initial capillary test when the test is conducted according to the schedule in Table 1 - Lead Screening Requirements and Medical Management Recommendations for Children ages 6 to 72 months.

Rev. 04/12
Rev. 07/15
Rev. 07/17

All venous sample lead screening tests conducted using any Magellan Diagnostic lead testing system should be laboratory analyzed by a properly accredited laboratory.

Lab Submission

Rev. 10/19
Rev. 04/23

The Decatur Regional Laboratory provides an analysis of blood lead specimens to Georgia children less than 72 months of age. The provider's office should contact the laboratory to use this service. GHHLPPP does not recommend or endorse the use of another lab.

Decatur Public Health Laboratory
1749 Clairmont Road
Decatur, Georgia 30033

Reporting Guidelines

Rev. 10/19

Laboratories attempt to test each lead specimen on the day it arrives. The reports are mailed back to providers on the same day. All laboratory data is sent monthly in electronic format to GHHLPPP.

Providers should report the results of all screening and follow-up blood lead level (BLL) tests to GHHLPPP. Because data from laboratories often do not include demographic information, complete reports from providers' offices are very important. If reports are not complete, GHHLPPP may contact providers' offices for missing information.

Results must be reported by:

Rev. 04/12
Rev. 04/14
Rev. 10/19
Rev. 04/23

- [State Electronic Notification Disease Surveillance System \(SendSS\)](#)
- SendSS is a web-based reporting system designed to collect information about [notifiable diseases](#) in Georgia.
 - o [Click here](#) for reporting instructions.
 - o [Click here](#) for Blood Lead Test Reporting Log spreadsheet template.

Providers who utilize the Lead Care II Analyzer to perform blood lead tests should report the Lead Care II Analyzer generated results to the GHHLPPP. On a weekly basis, providers should upload their test results to the [State Electronic Notification Disease Surveillance System \(SendSS\)](#). SendSS recommends providers keep track of their Lead Care II Analyzer test results by documenting those results in a Blood Lead Test Reporting Log provided by GHHLPPP (link above). If reports are not complete, GHHLPPP may contact providers' offices for missing information. Complete a new Blood Lead Reporting each week. Upload the Blood Lead Test Reporting Log and submit weekly to the GHHLPPP via SendSS. For uploading instructions, refer to the *SendSS Registration and Login Manual for Uploading Lead Report Files*.

Blood Lead Test Reporting Log: [Click here](#)



Registration and Login Manual for Uploading Lead Report Files may be accessed at:

https://dph.georgia.gov/sites/dph.georgia.gov/files/related_files/site_page/EnvHealthLeadReportingInstructions.pdf

- Before you can enter the system, you must accept the Privacy Statement [1]. Selecting “**I disagree with this statement**” will terminate your login and return you to the login page.
- The home page can be accessed with the following URL and is best viewed using Microsoft Internet Explorer Version 6.x.x. <https://sendss.state.ga.us>
- In order to gain access to SENDSS, you will first need to fill in a registration form and create a login.



Rev. 02/12
 Rev. 10/19
 Rev. 04/23

Georgia Comprehensive Childhood Lead Poisoning Prevention Program Case Management Guidelines

Blood Lead Level	Recommendations
< 3.5 ug/dl	<ul style="list-style-type: none"> • No safe threshold above “0” has been identified. Medical provider should provide anticipatory guidance for any blood lead level (BLL) above “0”.
3.5 – 9µg/dL	<p>Per GHHLPPP recommendations, the medical provider will:</p> <ul style="list-style-type: none"> • Conduct diagnostic (confirmatory) test (venous preferred) within 3 months. If child is < 12 months old or it is believed the BLL may be increasing rapidly, the test should be done earlier. • Test other children in the home <72 months of age who have not been tested in the last 6 months. • Conduct nutritional assessment • Continue testing at 3 month intervals until all the following conditions are met: <ul style="list-style-type: none"> ○ BLL has remained <3.5µg/dL for at least 6 months (two tests at least 3 months apart) ○ Lead hazards have been controlled. ○ There are no new sources of lead exposure <p>GHHLPPP or the Case Management Provider will send, by mail, email or deliver the following information to the caregiver:</p> <ul style="list-style-type: none"> • Child should receive a diagnostic (confirmatory) test (venous preferred) within 3 months. • Recommendation to have other children in the home <72 months of age who have not been tested in the last 6 months, tested. <p>GHHLPPP will refer the case to a DPH Regional Healthy Homes Coordinator (RHHC)</p> <p>Regional Healthy Homes Coordinator or Case Management Provider will give in-person (or in some cases by phone or mail, or email) to caregiver:</p> <ul style="list-style-type: none"> • Information on lead poisoning cause and effect, environmental lead hazard reduction, and nutrition. • Information on WIC services available. • Information on Children 1st Program information (newborns to 5 years old) • Information on Children’s Medical Services (CMS) if child =>5 years old <p>Regional Healthy Homes Coordinator will also:</p> <ul style="list-style-type: none"> • Conduct or arrange an environmental risk assessment by a certified risk assessor. Risk assessment should occur within 2 weeks of receiving referral from GHHLPPP.

10 – 44µg/dL	<p>Per GHHLPPP recommendations, the medical provider will:</p> <ul style="list-style-type: none">• Conduct diagnostic (confirmatory) test (venous preferred) within 1 week-1 month.• Test other children in the home <72 months of age who have not been tested in the last 6 months.• Conduct comprehensive medical evaluation including nutritional assessment.• Continue testing at 1-3 month intervals until all the following conditions are met:<ul style="list-style-type: none">- BLL has remained <3.5µg/dL for at least 6 months (two tests at least 3 months apart)- Lead hazards have been controlled- There are no new sources of lead exposure. <p>GHHLPPP or the Case Management Provider will send, by mail, the following information to the caregiver:</p> <ul style="list-style-type: none">• Child should receive a diagnostic (confirmatory) test (venous preferred) within 1 week to 1 month,• Recommendation to have other children in the home <72 months of age who have not been tested in the last 6 months, tested. <p>GHHLPPP will refer the case to a DPH Regional Healthy Homes Coordinator (RHHC)</p> <p>Regional Healthy Homes Coordinator or the Case Management Provider will give in-person (or in some cases by phone, email or mail) to caregiver:</p> <ul style="list-style-type: none">• Information on lead poisoning cause and effect, environmental lead hazard reduction, and nutrition.• Information on WIC services available.• Information on Children 1st Program information (newborns to 5 years old)• Information on Children’s Medical Services (CMS) if child =>5 years old <p>Regional Healthy Homes Coordinator will also:</p> <ul style="list-style-type: none">• Conduct or arrange an environmental risk assessment by a certified risk assessor. Risk assessment should occur within 1-2 weeks of receiving referral from GHHLPPP.• Send by mail or email (or in some cases call) a summary of risk assessment and recommendations to the caregiver, property owner and medical provider.• Provide a copy of the risk assessment in approved format to GHHLPPP.
---------------------	---

<p>45 – 69µg/dL</p>	<p>URGENT Per GHHLPPP recommendations, the medical provider will:</p> <ul style="list-style-type: none">○ Conduct diagnostic (confirmatory) test (venous preferred) within 24-48 hours.○ Test other children in the home <72 months of age who have not been tested in the last 6 months.○ Conduct comprehensive medical evaluation, including nutrition assessment and consider pharmacologic treatment. Contact the Georgia Poison Center for consultation.○ Continue testing at 1–2-month intervals until all the following conditions are met:<ul style="list-style-type: none">- BLL has remained <45µg/dL for at least 4 months (two tests at least 2 months apart) then start follow up blood lead testing at 3-month intervals until BLL has remained <3.5µg/dL for at least 6 months (two tests at least 3 months apart)- All identified lead hazards have been controlled Note: A child receiving chelation therapy <u>MAY NOT</u> return to the home until all lead hazards have been controlled.- There are no new sources of lead exposure. <p>GHHLPPP or the Case Management Provider will give, by phone, the following recommendation to the caregiver:</p> <ul style="list-style-type: none">○ Child should receive a diagnostic (confirmatory) test (venous preferred) within 24-48 hours,○ Recommendation to have other children in the home <72 months of age who have not been tested in the last 6 months, tested. <p>GHHLPPP will refer the case to a Regional Healthy Homes Coordinator (RHHC)</p> <p>Regional Healthy Homes Coordinator or the Case Management Provider will give in-person (or in some cases by phone) to caregiver:</p> <ul style="list-style-type: none">○ Information on lead poisoning cause and effect, environmental lead hazard reduction, and nutrition.○ Information on WIC services available.○ Information on Children 1st Program information (newborns to 5 years old)○ Information on Children’s Medical Services (CMS) if child =>5 years old <p>Regional Healthy Homes Coordinator will also:</p> <ul style="list-style-type: none">○ Conduct or arrange an environmental risk assessment by a certified risk assessor. Risk assessment should occur within 48 hours of receiving referral from GHHLPPP.○ Call-in a summary of risk assessment and recommendations to the caregiver, property owner and medical provider.○ Provide a copy of the risk assessment in approved format to GHHLPPP.
----------------------------	--

≥ 70µg/dL

MEDICAL EMERGENCY.

DO NOT DELAY MEDICAL TREATMENT.

Per GHHLPPP recommendations, the **medical provider** will:

- Conduct diagnostic (confirmatory) test (venous preferred) as emergency lab test.
- Conduct immediate medical evaluation and pharmacologic treatment. Contact the Georgia Poison Center for consultation.
- Test other children in the home <72 months of age who have not been tested in the last 6 months.
- Continue testing at 1–2-month intervals until all the following conditions are met:
 - BLL remains <45µg/dL for at least 4 months (two tests at least 2 months apart) then start follow up blood lead testing at 3-month intervals until BLL has remained <3.5µg/dL for at least 6 months (two tests at least 3 months apart)
 - All identified lead hazards have been controlled Note: A child receiving chelation therapy **MAY NOT** return to the home until all lead hazards have been controlled.
 - There are no new sources of lead exposure.

GHHLPPP or the Case Management Provider will give, by phone, the following recommendation to the caregiver:

- Child should receive a diagnostic (confirmatory) test (venous preferred) immediately.
- Recommendation to have other children in the home <72 months of age who have not been tested in the last 6 months, tested.

GHHLPPP will refer the case to a Regional Healthy Homes Coordinator (RHHC)

Regional Healthy Homes Coordinator or the Case Management Provider will give in-person (or in some cases by phone) to caregiver:

- Information on lead poisoning cause and effect, environmental lead hazard reduction, and nutrition.
- Information on WIC services available.
- Information on Children 1st Program information (newborns to 5 years old)
- Information on Children’s Medical Services (CMS) if child =>5 years old

Regional Healthy Homes Coordinator will also:

- Conduct or arrange an environmental risk assessment by a certified risk assessor. Risk assessment should occur within 24 hours of receiving referral from GHHLPPP.
- Call-in a summary of risk assessment and recommendations to the caregiver, property owner and medical provider
- Provide a copy of the risk assessment in approved format to GHHLPPP.
- If child must go to different housing unit post chelation, RHHC will inspect the new unit for lead hazards and inform medical provider that home is lead safe prior to child’s release from hospital.

Access guidelines at:

<https://dph.georgia.gov/sites/dph.georgia.gov/files/EnvHealth/Lead/EnvHealthLeadCaseManagementGuidelines2018.pdf>

*GA-AAP recommends a follow-up blood test within 3 months if the initial test is 5-9 µg/dL

Rev. 07/23

For questions on the GHHLPPP guidelines, please contact:

Georgia Healthy Homes and Lead Poisoning Prevention Program

200 Piedmont Avenue, SE | East Tower | Atlanta, GA 30334

Phone: 404-657-6534 Fax: 404-463-4039

<https://dph.georgia.gov/lead>

ENVIRONMENTAL LEAD RISK ASSESSMENTS

Rev. 01/13

Certified Lead Risk Assessors who conduct the initial environmental lead risk assessment should bill Medicaid using code T1028 and the appropriate diagnosis code. For follow up clearance inspections following removal of the lead hazards, the certified lead risk assessor should bill Medicaid using code T1028 with the U-1 modifier. For additional information, please consult the Diagnostic Screening and Preventive Services (DPS) Manual.

	<u>Procedure Code</u>
Initial lead investigation	T1028
Post hazard abatement	T1028 - (Modifier U-1)

Georgia and other Lead Resources

Rev. 04/14

Lead Information for Professionals and Parents

For information on lead poisoning and prevention, professionals and parents can call GHHLPPP (Georgia Healthy Homes and Lead Poisoning Prevention Program) at 404-657-6534 or the National Lead Information Center at 1-800-424-lead (5323).

Georgia Public Health Laboratory (GPHL)

The Georgia Public Health Laboratory, which has locations in Decatur and Waycross performs blood lead testing on children for the Georgia Healthy Homes and Lead Poisoning Prevention Program (GHHLPPP). Contact information:

GPHL
1749 Clairmont Road
Decatur, GA 30033
Phone 1-800-GEORGIA.

Emergency Information on Lead:

Call the Georgia Poison Center at 1-800-222-1222.

Sources of Lead

Common Sources of Lead

- Lead-based paint
- Lead dust, which is produced by aging lead-based paint
- Soil, which is contaminated by lead emissions from gasoline (prior to 1978), lead-based paint chips, storage of old batteries, etc.
- Water which flows through lead pipes or copper pipes soldered with lead (prior to 1986)
- Improperly glazed ceramic pottery and cooking utensils
- Construction workers, particularly those doing:
 - Department of Transportation (DOT) Sign Makers
 - Painting
 - Remodeling
 - Renovation
 - Road work (specifically painters)

Industries

- Battery manufacturers or reclamation
- Window replacement
- Bronze manufacture
- Firing range instructors
- Gas station attendants
- Glass manufacturers
- Lead pigment manufacture
- Lead smelters and refiners
- Plumbers, pipe fitters
- Policemen who work in automobile tunnels
- Printers
- Radiator manufacture or repair
- Shipbuilders
- Welders or Cutters – Steel burning or cutting (dismantling bridges, ships, etc.)
- Bridge or ship workers (including airports and boats)

Hobbies, Sports, Other

- Moonshine whiskey
- Car or boat repair
- Fishing
- Glazed pottery making
- Home remodeling
- Lead soldering
- Making lead shot or bullet
- Shooting at firing range
- Stained glass manufacture
- Additives to some “health foods” and imported candies
- Substance Use
- Toy soldiers (leaded)
- Folk Remedies – Most commonly found in Mexican, Asian Indian, and Middle Eastern groups. Names include: *Alarcon, Alkohl, Azarcon, Bali Goli, Coral, Ghasard, Greta, Liga, Pay-loo-ah, Rueda*. Cosmetics, used commonly by those from the Middle East and India.

APPENDIX B

Guidelines in Screening and Reporting for TB Disease and Infection

Tuberculin Skin Testing

Mantoux tuberculin skin testing is the standard method of identifying persons infected with *M. tuberculosis*. Multiple puncture tests should NOT be used to determine whether a person is infected.

The Mantoux test is performed by giving an intradermal injection of 0.1 ml of purified protein derivative (PPD) tuberculin containing 5 tuberculin units (TU) into either the volar or dorsal surface of the forearm. The injection should be made with a disposable tuberculin syringe, just beneath the surface of the skin, with the needle bevel facing upward. This should produce a discrete, pale elevation of the skin (a wheal) 6 mm to 10 mm in diameter.

The reaction to the Mantoux test should be read by a trained health care worker 48 to 72 hours after the injection. If a patient fails to show up for the scheduled reading, a positive reaction may still be measurable up to 1 week after testing.

However, if a patient who fails to return within 72 hours has a negative reaction, tuberculin testing should be repeated.

The area of induration (palpable swelling) around the site of injection is the reaction to tuberculin. The diameter of the indurated area should be measured across the forearm (perpendicular to the long axis). Erythema (redness) should not be measured. All reactions should be recorded in millimeters of induration, even those classified as negative. If no induration is found, “0 mm” should be recorded.

Reporting requirements

In Georgia, all tuberculosis must be reported immediately to the local county health department. Physicians, hospitals, laboratories and other health care providers are required to report any of the following:

- Any child less than 5 years discovered with Latent TB Infection
- Any confirmed case of TB
- Any suspected case of TB
- Any person being treated with two (2) or more anti-tuberculosis drugs
- Any positive culture for *Mycobacterium tuberculosis*
- Any positive smear for AFB (Acid Fast Bacilli)

How to report

- Report cases electronically through the [State Electronic Notifiable Disease Surveillance System \(SENDSS\)](#)
- Complete a Georgia [Notifiable Disease/Condition Report Form](#) and mail in an envelope marked CONFIDENTIAL, or
- Call your [District Health Office](#)
- If your District Health Office cannot be reached, call the Georgia Department of Public Health, TB Section at 404-657-2634.

Rev. 07/11
Rev. 10/14
Rev. 10/15

Childhood TB Risk Assessment Questionnaires

The AAP Bright Futures Guidelines for Health Supervision of Infants, Children, and Adolescents (3rd edition) recommends asking the following questions:

- 1) Was your child born in a country at high risk for tuberculosis?
- 2) Has your child traveled (had contact with resident populations) for longer than 1 week to a country a high risk for tuberculosis?
- 3) Has a family member or contact had tuberculosis or a positive tuberculin skin test?

Resource: <https://brightfutures.aap.org/materials-and-tools/Pages/default.aspx>

For more information about tuberculosis in Georgia: contact

Georgia Tuberculosis (TB) Section
2 Peachtree St. NW
12th Floor
Atlanta, GA 30303
(Phone) 404-657-2634
(Fax) 404-463-3460

The [Georgia TB Program](#) has the legal responsibility for all TB clients in Georgia regardless of who provides the direct services. TB services are available to all who fall within the service criteria without regard to the client's ability to pay.

Information also available at

[Tuberculosis \(TB\) Prevention and Control | Georgia Department of Public Health](#)

APPENDIX C-1

Vaccine Administration Codes

Rev. 04/24

Vaccine Administration Codes			
HIPAA Procedure Code	HIPAA Modifier	Procedure Code Description	Reimbursement Rate
90460	EP	Pediatric Immunization Administration Code. Immunization administration through 18 years of age via any route of administration, with counseling by physician or other qualified health care professional.	\$22.15
90471	EP	Non-Age Specific Immunization Administration Code. Immunization administration (includes percutaneous, intradermal, subcutaneous, or intramuscular injections); one vaccine (single or combination vaccine/toxoid)	\$23.78
90472	EP	Non-Age Specific Immunization Administration Code. Immunization administration (includes percutaneous, intradermal, subcutaneous, or intramuscular injections); each additional vaccine (single or combination vaccine/ toxoid)	\$12.99 \$18.69 (PCK)
90473	EP	Non-Age Specific Immunization Administration Code. Immunization administration (includes intranasal or oral administration); one vaccine (single or combination vaccine/toxoid)	\$23.78
90474	EP	Non-Age Specific Immunization Administration Code. Immunization administration (includes intranasal or oral administration); each additional vaccine (single or combination vaccine/toxoid)	\$12.10 \$18.69 (PCK)

Vaccine Administration Code for Face-to-Face Counseling

90460 - Vaccine administration through 18 years of age via any route of administration, with counseling by physician or other qualified health care professional.

Rev. 04/13
Rev. 07/13
Rev. 07/14
Rev. 10/14

90460 is reported when both of the following requirements are met: (1) The patient must be 18 years of age or younger. (2) The physician or other qualified health care professional must perform face-to-face vaccine counseling associated with the administration. *(Any clinical staff can do the actual administration of the vaccine per the physician's or the qualified health care professional's orders.)*

If both of these requirements are not met, report a non-age specific vaccine administration code(s) (90471-90474) instead.

90460 may be reported for more than one (1) unit of vaccine administered during a single office visit.

Note: Local Public Health Departments May use vaccine administration code 90460 only if a physician or other qualified health care professional performs face-to-face vaccine counseling associated with administration of the vaccine

A 'qualified health care professional' is an individual who by education, training, licensure/regulation, facility credentialing (when applicable), and payer policy is able to perform a professional service within their scope of practice and independently report a professional service. These professionals are distinct from 'clinical staff.' A clinical staff member is a person who works under the supervision of a physician or other qualified health care professional and who is allowed by law, regulation, facility, and payer policy to perform or assist in the performance of specified professional services, but who does not individually report any professional services. (CPT 2012)

Non-Age Specific Vaccine Administration Codes

Rev. 01/14

These codes must be used when there is no face-to-face physician counseling associated with vaccine administration. The add-on codes (90472, 90474) may also be used in conjunction with the 90460 code.

Codes 90471 and 90473 are used to code for the first vaccine given during a single office visit. Codes 90472 and 90474 are considered add-on codes (*hence the + symbol next to them*) to 90471 and 90473, respectively. This means that the provider will use 90472 and 90474 in addition to 90471 or 90473 if more than one vaccine is administered during a visit. Providers may use 90460 for the first (i.e., counseled) vaccine and 90472 or 90474 for the second (i.e., non-counseled) vaccine. Note that there can only be one first administration during a given visit.

Rev. 04/13

90471 - Vaccine administration (includes percutaneous, intradermal, subcutaneous, or intramuscular injections); one vaccine (single or combination vaccine/toxoid)
Do not report 90471 in conjunction with 90473 or 90460

+90472 - Vaccine administration (includes percutaneous, intradermal, subcutaneous, or intramuscular injections); each additional vaccine (single or combination vaccine/toxoid) (List separately in addition to code for primary procedure.)
May use 90472 in conjunction with 90471 or 90473 or 90460

90473 - Vaccine administration (includes intranasal or oral administration); one vaccine (single or combination vaccine/toxoid)
Do not report 90473 in conjunction with 90471 or 90460

+90474 - Vaccine administration (includes intranasal or oral administration); each additional vaccine (single or combination vaccine/toxoid) (List separately in addition to code for primary procedure.)
May use 90474 in conjunction with 90471 or 90473 or 90460

Vaccine Administration Codes for 19 – 20-year-olds

Rev. 07/13

Rev. 10/14

EPSDT providers may bill the EPSDT benefit for vaccines administered to members nineteen (19) years of age through twenty (20) years of age. Providers must include the vaccine's CPT, NDC, and diagnosis code, along with the appropriate vaccine administration code(s) [90471, 90472, 90473, 90474] on the EPSDT claim. The Division will reimburse for the vaccine product (providers must use their own stock of vaccines) and for vaccine administration.

Billing Tips

Rev. 04/13
Rev. 07/13

- Code the vaccine administration with the appropriate vaccine administration code and the EP modifier.
- For the vaccine, code the vaccine product code, the associated diagnosis code and the EP modifier.
- The primary vaccine administration code (90460, 90471, or 90473) must precede the add-on vaccine administration code(s) (90472 or 90474), if applicable.
- The vaccine product code must immediately follow the corresponding vaccine administration code.

Rev. 07/15

- Code the vaccine administration code(s), the vaccine product code(s), and the preventive or interperiodic visit on the same claim when vaccines are administered during a preventive or interperiodic visit. Each vaccine administration code should be listed only once per claim. If multiple vaccine product codes correspond to the same vaccine administration code, the vaccine administration code is listed once with the appropriate number of units indicated.
- 90460 may be reported for more than one (1) unit of vaccine administered during a single office visit.

Rev. 07/14
Rev. 10/15
Rev. 04/18
Rev. 07/19
Rev. 01/20

- May report diagnosis code Z00.121 or Z00.129 or Z23 with each of the vaccine administration codes ONLY when vaccines are administered during EPSDT preventive health visits for members through age 17 years.
- May report diagnosis code Z00.00 or Z00.01 or Z23 with the applicable vaccine administration code ONLY when vaccines are administered during EPSDT preventive health visits for members aged 15 years through 20 years.

Rev. 04/14
Rev. 07/14
Rev. 10/14
Rev. 04/15
Rev. 01/21
Rev. 04/24

- Use the appropriate vaccine diagnosis code with the vaccine administration code when the vaccine is administered outside of the EPSDT preventive health visit.
- Code the EPSDT preventive visit (9938x or 9939x) with the EP and the 25 modifiers when vaccines are administered during the preventive health visit.
- Code the EPSDT interperiodic visits (99202-99205 or 99212-99215) with the EP and the 25 modifiers when vaccines are administered during the interperiodic health visit.
- The National Correct Coding Initiative (NCCI) does not allow reimbursement of the 99211 code when it is billed together with any of the vaccine administration codes regardless of whether the 25 modifier is appended to the 99211 code.

Rev. 01/24
Rev. 04/24

- Code the administration of Beyfortus with vaccine administration code 96380 or 96381 and the EP modifier. The Beyfortus product code should immediately follow the vaccine administration code. Codes 96380 and 96381 replaces the previously instructed generic cpt code 96372. For claims beginning with effective date 3/1/2024, 96372 should no longer be reported for the administration of Beyfortus.

Rev. 04/24

RSV administration codes	Reimbursement Rates
96380 96381	\$10.00 \$18.50 (PCK)

APPENDIX C-2

Vaccine Procedure and Diagnosis Codes (Ages Birth through 18 years)

The following vaccine procedure and diagnosis codes must be included on the claim, following the vaccine administration code, when billing for vaccine administration.

HIPAA Procedure Code	HIPAA Modifier	Procedure Code Description
90380		Respiratory Syncytial Virus (RSV) monoclonal antibody, recombinant, 50 mg (0.5 ml) dosage, for intramuscular use (Beyfortus) (birth up to 24 months)
90381	EP	Respiratory Syncytial Virus (RSV) monoclonal antibody, recombinant, 100 mg (1.0 ml) dosage, for intramuscular use (Beyfortus) (birth up to 24 months)
90678		Respiratory Syncytial Virus (RSV) vaccine, pref, subunit, bivalent, for intramuscular use (Abrysvo)
90619	EP	Meningococcal conjugate vaccine, serogroups A, C, W, Y, quadrivalent, tetanus toxoid carrier (MENACWY-TT), for intramuscular use (2-18 years) [MenQuadfi]
90620	EP	Meningococcal recombinant protein and outer membrane vesicle vaccine, serogroup B (MenB-4C), 2 dose schedule, for intramuscular use (10-18 years) [Bexsero]
90621	EP	Meningococcal recombinant lipoprotein vaccine, serogroup B (MenB-FHbp), 2 or 3 dose schedule, for intramuscular use (10-18 years) [Trumenba]
90633	EP	Hepatitis A vaccine (HepA), pediatric/adolescent dosage-2 dose schedule, for intramuscular use
90636	EP	Combination Hepatitis A/Hepatitis B vaccine, adult dosage, 3 dose, for intramuscular use (18 years)
90647	EP	Haemophilus influenzae type B vaccine (Hib), PRP-OMP conjugate, 3 dose schedule, for intramuscular use
90648		Haemophilus influenzae type b vaccine (Hib), PRP-T conjugate, 4 dose schedule, for intramuscular use
90651	EP	Human Papillomavirus (HPV) vaccine types 6, 11, 16, 18, 31, 33, 45, 52, 58 nonvalent (9vHPV), 2 or 3-dose schedule, for intramuscular use (females and males 9 years and older) [Gardasil 9]
90674		Influenza virus vaccine, quadrivalent (ccIIV4), derived from cell cultures, subunit preservative and antibiotic free, 0.5mL dosage, for intramuscular use
90685		Influenza virus vaccine, quadrivalent (IIV4), split virus, preservative free, 0.25 mL dosage, for intramuscular use
90686	EP	Influenza virus vaccine, quadrivalent (IIV4), split virus, preservative free, 0.5 mL dosage, for intramuscular use
90687		Influenza virus vaccine, quadrivalent (IIV4), split virus, 0.25 mL dosage, for intramuscular use
90688 90672	EP	Influenza virus vaccine, quadrivalent (IIV4), split virus, 0.5 mL dosage, for intramuscular use Influenza virus vaccine (FluMist), quadrivalent, live (LAIV4), for intranasal use (2 years and older)

Rev. 01/08
Rev. 07/08
Rev. 07/10
Rev. 07/11
Rev. 07/12
Rev. 01/13
Rev. 03/13
Rev. 01/14
Rev. 04/14
Rev. 07/14
Rev. 10/14
Rev. 01/15
Rev. 04/15
Rev. 10/15
Rev. 01/16
Rev. 04/16
Rev. 10/16
Rev. 01/17
Rev. 01/18
Rev. 04/18
Rev. 10/18
Rev. 01/19
Rev. 10/21
Rev. 01/24
Rev. 04/24

Rev. 01/08
 Rev. 07/08
 Rev. 07/10
 Rev. 07/11
 Rev. 07/12
 Rev. 01/13
 Rev. 03/13
 Rev. 01/14
 Rev. 04/14
 Rev. 07/14
 Rev. 10/14
 Rev. 01/15
 Rev. 04/15
 Rev. 10/15
 Rev. 01/16
 Rev. 04/16
 Rev. 10/16
 Rev. 01/17
 Rev. 01/18
 Rev. 04/18
 Rev. 07/19
 Rev. 10/21
 Rev. 04/23

HIPAA Procedure Code	HIPAA Modifier	Procedure Code Description
90756	EP	Influenza virus vaccine, quadrivalent (ccIV4), derived from cell cultures, subunit, antibiotic free, 0.5mL dosage, for intramuscular use
90670 90671 90677	EP	Pneumococcal conjugate vaccine, 13 valent (PCV13), for intramuscular use Pneumococcal conjugate vaccine, 15 valent (PCV15), for intramuscular use Pneumococcal conjugate vaccine, 20 valent (PCV20), for intramuscular use
90680	EP	Rotavirus vaccine, pentavalent (RV5), 3 dose schedule, live, for oral use (RotaTeq)
90681	EP	Rotavirus vaccine, human, attenuated (RV1), 2 dose schedule, live, for oral use (Rotarix)
90696	EP	Diphtheria, tetanus toxoids, and acellular pertussis vaccine and inactivated poliovirus vaccine, (DTaP-IPV), for intramuscular use (ages 4-6 years)
90697		Diphtheria, tetanus toxoids, acellular pertussis (whooping cough), haemophilus influenzae type B, hepatitis B and polio vaccine, (DTaP-IPV-Hib-HepB), for intramuscular use (ages 6 weeks to 5 years) [Vaxelis]
90698		Diphtheria, tetanus toxoids, acellular pertussis vaccine, Haemophilus influenza type B, and inactivated poliovirus vaccine (DTaP-IPV/Hib), for intramuscular use
90700		Diphtheria, tetanus toxoids, and acellular pertussis vaccine (DTaP), for intramuscular use (ages younger than 7 years)
90707	EP	Measles, mumps, and rubella virus vaccine (MMR), live, for subcutaneous use
90710	EP	Measles, mumps, rubella, and varicella vaccine (MMRV), live, for subcutaneous use
90713	EP	Poliovirus vaccine (IPV), inactivated, for subcutaneous or intramuscular use
90714	EP	Tetanus and diphtheria toxoids (Td) adsorbed, preservative free, for intramuscular use (7 years to 18 years, 11 months)
90715	EP	Tetanus, diphtheria toxoids and acellular pertussis vaccine (Tdap), for intramuscular use (7 years to 18 years, 11 months)
90716	EP	Varicella virus vaccine (VAR), live, for subcutaneous use
90723	EP	Diphtheria, tetanus toxoids, acellular pertussis vaccine, Hepatitis B and inactivated poliovirus vaccine (DTaP-Hep B-IPV), for intramuscular use
90732	EP	Pneumococcal polysaccharide vaccine, 23-valent (PPSV23), adult or immunosuppressed patient dosage, when administered to 2 years or older, for subcutaneous or intramuscular use
90734	EP	Meningococcal conjugate vaccine, serogroups A, C, W, Y, quadrivalent, diphtheria toxoid carrier, (MenACWY-D) or CRM197 carrier (MenACWY-CRM), for intramuscular use
90739		Hepatitis B vaccine (HepB), CPG-Adjuvanted, adult dosage, 2 or 4 dose schedule, for intramuscular use
90744	EP	Hepatitis B vaccine (HepB), pediatric/adolescent dosage, 3 dose schedule, for intramuscular use
90747		Hepatitis B vaccine (HepB), dialysis or immunosuppressed patient dosage, 4 dose schedule, for intramuscular use

Vaccine Procedure and Diagnosis Codes

(Ages 19 through 20 years)

The Division will reimburse for the vaccine product (providers must use their own stock of vaccines) and for vaccine administration for vaccines administered to Medicaid-eligible members nineteen (19) years of age through twenty (20) years of age. Providers who administer any one of the vaccines listed below will receive reimbursement for administering each vaccine PLUS reimbursement for the vaccine product. The reimbursement rates for the vaccine products may be found in the Physicians' Injectable Drug List Manual. Providers must include the vaccine's procedure and diagnosis codes on the claim.

Rev. 10/13
 Rev. 01/14
 Rev. 04/14
 Rev. 07/14
 Rev. 10/14
 Rev. 04/15
 Rev. 07/15
 Rev. 10/15
 Rev. 01/16
 Rev. 04/16
 Rev. 10/16
 Rev. 01/17
 Rev. 10/17
 Rev. 01/18
 Rev. 04/18
 Rev. 10/18
 Rev. 01/19
 Rev. 10/20
 Rev. 10/21
 Rev. 04/23
 Rev. 10/23

Vaccine Procedure and Diagnosis Codes		
(Ages 19 through 20 years)		
HIPAA Procedure Code	HIPAA Modifier	Procedure Code Description
90619		Meningococcal conjugate vaccine, serogroups A, C, W, Y, quadrivalent, tetanus toxoid carrier (MENACWY-TT), for intramuscular use (MenQuadfi)
90620	EP	Meningococcal recombinant protein and outer member vesicle vaccine, serogroup B (MenB-4C), 2 dose schedule, for intramuscular use (Bexsero)
90621		Meningococcal recombinant lipoprotein vaccine, serogroup B (MenB-FHbp), 2 or 3 dose schedule, for intramuscular use (Trumenba)
90632		Hepatitis A vaccine (HepA), adult dosage, for intramuscular use
90636	EP	Combination Hepatitis A/Hepatitis B vaccine, adult dosage, 3 dose, for intramuscular use
90739	EP	Hepatitis B vaccine (HepB), CPG-Adjuvanted, adult dosage, 2 dose or 4 dose schedule, for intramuscular use
90746		Hepatitis B vaccine (HepB), adult dosage 3 dose schedule, for intramuscular use
90747		Hepatitis B vaccine (HepB), dialysis or immunosuppressed patient dosage 4 dose, for intramuscular use
90759		Hepatitis B vaccine (HepB), 3AG 10 mcg, 3 dose schedule, for intramuscular use (Hevbrio)
90651	EP	Human Papillomavirus HP vaccine types 6, 11, 16, 18, 31, 33, 45, 52, 58, nonvalent (9vHPV), 2 or 3-dose schedule, for intramuscular use (females & males) [Gardasil 9]
90674	EP	Influenza virus vaccine, quadrivalent (ccIIV4), derived from cell cultures, subunit, preservative and antibiotic free, 0.5mL dosage, for intramuscular use
90682		Influenza virus vaccine, quadrivalent (RIV4), derived from recombinant DNA, hemagglutinin (HA) protein only, preservative and antibiotic free, for intramuscular use
90686		Influenza virus vaccine, quadrivalent (IIV4), split virus, preservative free, 0.5 mL dosage, for intramuscular use
90688		Influenza virus vaccine, quadrivalent (IIV4), split virus, 0.5 mL dosage, for intramuscular use
90672		Influenza virus vaccine (FluMist), quadrivalent, live (LAIV4), for intranasal use (2 years and older)

Vaccine Procedure and Diagnosis Codes

(Ages 19 through 20 years)

HIPAA Procedure Code	HIPAA Modifier	Procedure Code Description
90756		Influenza virus vaccine, quadrivalent (ccIIV4), derived from cell cultures, subunit, antibiotic free, 0.5mL dosage, for intramuscular use
90707	EP	Measles, mumps, and rubella virus vaccine (MMR), live, for subcutaneous use
90713	EP	Poliovirus vaccine (IPV), inactivated, for adults at high risk, for subcutaneous or intramuscular use
90714	EP	Tetanus and diphtheria toxoids (Td) absorbed, preservative free, for intramuscular use
90715	EP	Tetanus, diphtheria toxoids and acellular pertussis vaccine (Tdap), for intramuscular use
90716	EP	Varicella virus vaccine (VAR), live, for subcutaneous use
90670 90671 90677	EP	Pneumococcal conjugate vaccine, 13 valent (PCV13), for intramuscular use Pneumococcal conjugate vaccine, 15 valent (PCV15), for intramuscular use Pneumococcal conjugate vaccine, 20 valent (PCV20), for intramuscular use
90732		Pneumococcal polysaccharide vaccine, 23-valent (PPSV23), adult or immunosuppressed patient dosage, for subcutaneous or intramuscular use
90734	EP	Meningococcal conjugate vaccine, serogroups A, C, W, Y, quadrivalent, diphtheria toxoid carrier, (MenACWY-D) or CRM197 carrier (MenACWY-CRM), for intramuscular use
90678	EP	Respiratory Syncytial Virus (RSV) vaccine, pref, subunit, bivalent, for intramuscular use (Abrysvo)

Rev. 10/13
Rev. 01/14
Rev. 04/14
Rev. 07/14
Rev. 10/14
Rev. 04/15
Rev. 07/15
Rev. 10/15
Rev. 01/16
Rev. 04/16
Rev. 10/16
Rev. 01/17
Rev. 10/17
Rev. 01/18
Rev. 04/18
Rev. 10/18
Rev. 01/19
Rev. 07/19
Rev. 04/23
Rev. 04/24

APPENDIX C-3

Tuberculin (TB) Skin Test Procedure Codes

The maximum reimbursement rate for the TB skin test provided by private providers is \$8.13 and \$3.00 for public health providers.

Use the following procedure and diagnosis codes to document the Tuberculin Skin Test.

HIPAA Procedure Code	ICD-10 Diagnosis Code	HIPAA Modifier	Procedure Code Description	Fee For Service Reimbursement
86580	Z11.1	EP	TB Skin Test	\$3.00 (public) \$8.13 (private)

APPENDIX C-4

Blood Lead Level Testing Procedure Codes

Blood Lead Level (BLL):

Blood Lead Screenings should be performed at the 12 and 24 month periodic visits (or catch-up visits). There are two tests used to obtain blood lead level (BLL) specimens: capillary blood test or venous blood test. Finger stick capillary blood tests can be done as the initial screening; however, a lab analyzed sample is necessary for confirmation. The specimen may be sent to a private laboratory for analysis, the analysis may be performed using an in-office blood lead level analyzer (the Lead Care II Analyzer uses capillary blood) or sent to the Georgia Public Health Laboratory (GPHL).

Rev. 07/11
Rev. 07/12
Rev. 03/13
Rev. 04/14
Rev. 07/14
Rev. 04/15
Rev. 10/15
Rev. 01/16
Rev. 10/19
Rev. 10/21

Providers should bill code 83655 with the EP modifier and appropriate ICD-10 diagnosis code which indicates the child is receiving a screening blood lead test (*i.e.*, Z13.88, Z00.121, Z00.129, Z77.011) on the same claim with the appropriate blood test code 36415 (blood lead venous) or 36416 (blood lead capillary).

Providers who send BLL specimen to the Georgia Public Health Laboratory (GPHL) will be billed \$10.00 lab handling fee assessed by the GPHL. Therefore, Fee For Service (FFS) providers should bill code 83655 with modifiers EP, 90 or EP, 91 and appropriate ICD-10 diagnosis code (*i.e.*, Z13.88, Z00.121, Z00.129, Z77.011) on the same claim with the appropriate codes (36415 or 36416). This billing will result in a FFS reimbursement of \$10.00 for the lab handling fee assessed by the GPHL. This reimbursement is only available when documentation supports that the BLL specimen was sent to the GPHL.

APPENDIX C-5

Covid-19 Vaccines & Administration Codes

Rev. 10/22

For listing of Covid-19 Vaccines & Administration Codes, see the '*Provider's Administered Drug List – Covid-19 Vaccine Administration Codes*' Fee Schedule located on GAMMIS.

APPENDIX D

Children's Intervention Services

The Children's Intervention Services (CIS) program offers coverage for restorative and/or rehabilitative services to eligible members in non-institutional settings. Services must be determined medically necessary and be recommended and documented as appropriate interventions by a physician or other licensed practitioner of the healing arts, within their scope of practice under state law, for the maximum reduction of physical disability or developmental delay and restoration of the member to the best possible functional level.

The CIS program is comprised of seven intervention services that must be provided by licensed practitioners of the healing arts. The seven services are:

- Audiology
- Nursing
- Nutrition provided by licensed dietitians
- Occupational Therapy
- Physical Therapy
- Counseling provided by licensed clinical Social Workers
- Speech-language Pathology


Qualified providers must be currently licensed as audiologists, registered nurses, dietitians, occupational therapists, physical therapists, clinical social workers, or speech-language pathologists.

APPENDIX E

Rev. 4/12
Rev. 4/21
Rev. 07/23

Non-Emergency Medical Transportation (NEMT)

FACT SHEET



**GEORGIA DEPARTMENT
OF COMMUNITY HEALTH**

Frequently Asked Questions about Non-Emergency Medical Transportation (NEMT)

The Georgia Department of Community Health's (DCH) Non-Emergency Medical Transportation (**NEMT**) services are defined as medically necessary, cost-effective transportation for any eligible Medicaid member (and escort, if required) with no other means of transportation available to any Medicaid-reimbursable service to receive treatment, medical evaluation, obtain prescription drugs or medical equipment. NEMT is a ride-share program and multiple members may be riding in the same vehicle.

To download a copy of the NEMT policies and procedures manual go to: <https://www.mmis.georgia.gov/portal>, select *Provider Information*, then select *Provider Manuals*.

For Members

1. How does the NEMT program operate?

The NEMT program provides transportation through a NEMT Broker System. Five NEMT regions have been established in the state: North, Atlanta, Central, East and Southwest. DCH has contracted with a broker in each of the five NEMT regions to administer and provide non-emergency medical transportation for eligible members.

2. Who are the NEMT Brokers and how do I arrange transportation?

There are two brokers providing NEMT services: Modivcare Solutions, LLC and Verida, Inc. A member or person acting on behalf of a member must contact the broker serving the county in which he/she lives (see chart below) between the hours of 7 a.m. to 6 p.m., Monday through Friday to request NEMT services. Requests must be made at least three workdays prior to the non-urgent, scheduled appointment. The three-day advance scheduling includes the day of the call but not the day of the appointment. Requests for urgent care situations and other exceptions may be arranged more quickly.

For purposes of the NEMT contract, urgent care is defined as an unscheduled episodic situation, in which there is no immediate threat to life or limb. However, the member must be seen on the date of the request and treatment cannot be delayed until the next day. Valid requests for urgent care transport will be honored within three hours of the time the request is made. *The NEMT Broker call centers are operational 24/7 for "Where's My Ride" and hospital discharges.*

Region	Broker/Phone Number	Counties Served
North	Verida • Toll free: 866-388-9844 • Local: 678-510-4555	Banks, Barrow, Bartow, Catoosa, Chattooga, Cherokee, Cobb, Dade, Dawson, Douglas, Fannin, Floyd, Forsyth, Franklin, Gilmer, Gordon, Habersham, Hall, Haralson, Jackson, Lumpkin, Morgan, Murray, Paulding, Pickens, Polk, Rabun, Stephens, Towns, Union, Walker, Walton, White and Whitfield
Atlanta	Verida • 404-209-4000	Fulton, DeKalb and Gwinnett
Central	Modivcare • Toll free: 888-224-7961	Baldwin, Bibb, Bleckley, Butts, Carroll, Clayton, Coweta, Dodge, Fayette, Heard, Henry, Jasper, Jones, Lamar, Laurens, Meriwether, Monroe, Newton, Pike, Putnam, Rockdale, Spalding, Telfair, Troup, Twiggs and Wilkinson
East	Modivcare • Toll free: 888-224-7988	Appling, Bacon, Brantley, Bryan, Bulloch, Burke, Camden, Candler, Charlton, Chatham, Clarke, Columbia, Effingham, Elbert, Emanuel, Evans, Glascock, Glynn, Greene, Hancock, Hart, Jeff Davis, Jefferson, Jenkins, Johnson, Liberty, Lincoln, Long, Madison, McDuffie, McIntosh, Montgomery, Oconee, Oglethorpe, Pierce, Richmond, Screven, Taliaferro, Tattnall, Toombs, Treutlen, Ware, Warren, Washington, Wayne, Wheeler and Wilkes
Southwest	Modivcare • Toll free: 888-224-7985	Atkinson, Baker, Ben Hill, Berrien, Brooks, Calhoun, Chattahoochee, Clay, Clinch, Coffee, Colquitt, Cook, Crawford, Crisp, Decatur, Dooly, Dougherty, Early, Echols, Grady, Harris, Houston, Irwin, Lanier, Lee, Lowndes, Macon, Marion, Miller, Mitchell, Muscogee, Peach, Pulaski, Quitman, Randolph, Schley, Seminole, Stewart, Sumter, Talbot, Taylor, Terrell, Thomas, Tift, Turner, Upson, Webster, Wilcox and Worth

1 of 3

March 2023

FACT SHEET



Frequently Asked Questions about Non-Emergency Medical Transportation (NEMT)

3. What modes of transportation are available and how are they selected?

The broker uses the most appropriate mode of transportation – minibus, wheelchair vans, automobiles, stretcher vans, public or para-transit that meets the member's health needs. In addition, the broker may use gas reimbursement and taxicabs for urgent care and/or back-up services when necessary.

The broker determines the most appropriate and cost-effective mode of transportation based on the information provided by the member at the time the reservation is made. All transportation must be scheduled and authorized by the NEMT broker.

4. How do NEMT brokers use public transit?

Brokers have procedures in place to determine whether public transportation is accessible to and appropriate for the member requesting service. Both NEMT brokers can identify electronic route information from public transit providers to determine if a member's pickup and drop-off locations are on a public transit route. There are also procedures in place for timely distribution of transit tokens/passes to the member and or escort (if applicable) using public transit.

5. May an escort or attendant accompany me to an appointment?

The broker must allow, without charge to the escort or member, one escort to accompany a member when medically necessary. An escort is defined as an individual whose presence is required to assist a member during transport and while at the place of treatment. An escort must be 18 years of age or older.

An attendant is defined as a staff person of the broker or provider present during the trip when in the broker's judgement it is necessary to have an adult helper to assure the safety of all passengers. The attendant remains with the vehicle after the member has left the vehicle. Attendants must complete a training program to include first aid training, a passenger assistance orientation program, and a safety and sensitivity program.

6. May a parent or guardian take someone else with them to an appointment with a family member?

Only one escort may accompany a member to an appointment. NEMT is a ride-share program and there may be other members scheduled to ride with you in the same vehicle. As such, seating may not be available for the additional individual or family members who are not scheduled to receive services.

NOTE: NEMT Policy does not prohibit the broker from allowing a minor to accompany an adult member to an appointment because there is no one else to stay with the minor. If there is room or an available seat that is not otherwise occupied, the broker may allow a minor to be transported with the adult member requiring treatment. This arrangement must be authorized by the broker prior to transport.

7. Is a co-pay required to ride NEMT?

There is no co-payment required.

FACT SHEET



Frequently Asked Questions about Non-Emergency Medical Transportation (NEMT)

8. What if I have a question about or problem/complaint with NEMT services?

If you have a question, comment, or complaint about NEMT, you may call the NEMT Broker directly or call the Member Contact Center at 1-866-211-0950. You may also contact DCH at 404-656-4646, e-mail tmalcolm@dch.ga.gov or 404-651-6911, e-mail sramsue@dch.ga.gov. To check the status of your ride, please call your broker and press option #2 for the "Where's My Ride" line.

9. How is the quality of services monitored for NEMT?

NEMT staff monitors the quality of service provided, including but not limited to: reviewing various required monthly, quarterly and annual reports; conducting on-site reviews to ensure compliance and possibly riding on a NEMT vehicle to monitor service; and attending broker/provider meetings. In addition, the broker must contract with an independent agent to conduct annual customer service satisfaction surveys and provide the methodology for analyzing the data and report results to DCH.

For Providers

1. How do I become a NEMT provider with Georgia Medicaid?

Transportation providers must contact Modivcare and/or Verida to provide NEMT services to our members.

For more information on the NEMT program, please feel free to visit our website, call or email us at the information provided in this fact sheet.

Rider Tips

- ✓ Be ready!
- ✓ Be respectful and courteous of others!
- ✓ Become familiar with the Broker issued *Member Brochure/Rider's Guide!*
- ✓ Always call the Broker directly to inquire about your ride!
- ✓ Always immediately notify Broker of any cancellations or changes in your schedule!
- ✓ Always carry your Broker's telephone number with you!

March 2023

NEMT REGIONS & COUNTIES SERVED

Rev. 4/12
 Rev. 7/12
 Rev. 4/21
 Rev. 7/23

Region	NEMT Broker & Phone Number	Counties Served
North	<p>Verida <i>(formerly Southeastrans)</i></p> <p><i>Toll free</i> 1-866-388-9844</p> <p><i>Local</i> 678-510-4555</p>	Banks, Barrow, Bartow, Catoosa, Chattooga, Cherokee, Cobb, Dade, Dawson, Douglas, Fannin, Floyd, Forsyth, Franklin, Gilmer, Gordon, Habersham, Hall, Haralson, Jackson, Lumpkin, Morgan, Murray, Paulding, Pickens, Polk, Rabun, Stephens, Towns, Union, Walker, Walton, White and Whitfield
Atlanta	<p>Southeastrans <i>(formerly Southeastrans)</i></p> <p>404-209-4000</p> <p><i>Note: For Georgia Families 360°</i></p> <p>1-866-991-6701</p>	Fulton, DeKalb and Gwinnett
Central	<p>ModivCare <i>(formerly LogistiCare)</i></p> <p><i>Toll free</i> 1-888-224-7981</p>	Baldwin, Bibb, Bleckley, Butts, Carroll, Clayton, Coweta, Dodge, Fayette, Heard, Henry, Jasper, Jones, Lamar, Laurens, Meriwether, Monroe, Newton, Pike, Putnam, Rockdale, Spalding, Telfair, Troup, Twiggs and Wilkinson
East	<p>ModivCare <i>(formerly LogistiCare)</i></p> <p><i>Toll free</i> 1-888-224-7988</p> <p><i>Note: For Crisis Stabilization Units and Psychiatric Residential Treatment Facilities</i></p> <p>1-800-486-7642 Ext. 461 or 436</p>	Appling, Bacon, Brantley, Bryan, Bulloch, Burke, Camden, Candler, Charlton, Chatham, Clarke, Columbia, Effingham, Elbert, Emanuel, Evans, Glascock, Glynn, Greene, Hancock, Hart, Jeff Davis, Jefferson, Jenkins, Johnson, Liberty, Lincoln, Long, Madison, McDuffie, McIntosh, Montgomery, Oconee, Oglethorpe, Pierce, Richmond, Screven, Taliaferro, Tattnall, Toombs, Treutlen, Ware, Warren, Washington, Wayne, Wheeler and Wilkes
Southwest	<p>ModivCare <i>(formerly LogistiCare)</i></p> <p><i>Toll free</i> 1-888-224-7985</p>	Atkinson, Baker, Ben Hill, Berrien, Brooks, Calhoun, Chattahoochee, Clay, Clinch, Coffee, Colquitt, Cook, Crawford, Crisp, Decatur, Dooly, Dougherty, Early, Echols, Grady, Harris, Houston, Irwin, Lanier, Lee, Lowndes, Macon, Marion, Miller, Mitchell, Muscogee, Peach, Pulaski, Quitman, Randolph, Schley, Seminole, Stewart, Sumter, Talbot, Taylor, Terrell, Thomas, Tift, Turner, Upson, Webster, Wilcox and Worth

APPENDIX F

Childhood Obesity – Weight Assessment and Counseling for Nutrition and Physical Activity for Children/Adolescents

Rev. 10/13
Rev. 07/14
Rev. 07/16

Body Mass Index (BMI)

What is BMI?

Body Mass Index (BMI) is used as a screening tool to identify possible weight problems for children and adolescents. The Centers for Disease Control and Prevention (CDC) and the American Academy of Pediatrics (AAP) recommend the use of BMI to screen for overweight and obesity in children and adolescents aged 2 through 19 years.

BMI is a number calculated from a child’s weight and height. BMI does not measure body fat directly, but it is a reliable indicator of body fatness for most children and adolescents.

For children and adolescents, BMI is age- and sex-specific and is often referred to as BMI-for-age. A child’s weight status is determined using an age- and sex-specific percentile for BMI rather than the BMI categories used for adults because children’s body composition varies as they age and varies between boys and girls.

What is a BMI percentile?

After the BMI is calculated for children and adolescents, the BMI number is plotted on the BMI-for-age growth charts (for either girls or boys) to obtain a percentile ranking. Percentiles are the most commonly used indicator to assess the size and growth patterns of individual children in the United States. The percentile indicates the relative position of the child’s BMI number among children of the same sex and age. The CDC’s [growth charts](#) show the weight status categories used with children and adolescents.

WEIGHT STATUS CATEGORY	PERCENTILE RANGE
Underweight	Less than the 5 th percentile
Healthy Weight	5 th percentile to less than the 85 th percentile
Overweight	85 th to less than the 95 th percentile
Obese	Equal to or greater than the 95 th percentile

These percentiles are based on the CDC’s [growth charts](#) which are available at <http://www.cdc.gov/growthcharts/>

How to calculate BMI using a handheld calculator?

Rev. 10/13

The BMI can be calculated using either English or metric units.

With the metric system, the formula for BMI is weight in kilograms divided by height in meters squared. Since height is commonly measured in centimeters, an alternate calculation formula, dividing the weight in kilograms by the height in centimeters squared, and then multiplying the result by 10,000, can be used.

$$\text{Formula: weight (kg) / [height (m)]}^2$$

When using English measurements, ounces (oz.) and fractions must be changed to decimal values. Then, calculate BMI by dividing weight in pounds (lbs.) by height in inches (in) squared and multiplying by a conversion factor of 703. Plot the calculated BMI to obtain the BMI percentile.

Rev. 07/14

$$\text{Formula: weight (lb.) / [height (in)]}^2 \times 703$$

How to interpret BMI?

Calculating the BMI-for-age for children of different ages and sexes may yield the same numeric result, but the result may fall at a different percentile for each child for one or both of the following reasons:

- The normal BMI-related changes that take place as children age and as growth occurs (ex. The amount of body fat changes with age).
- The normal BMI-related differences between sexes (ex. The amount of body fat differs between girls and boys).

Documentation for HEDIS Compliance – Weight Assessment

Documentation in the medical record for BMI must include the following:

- Height
- Weight
- BMI percentile

Documentation must include height, weight and BMI percentile during the measurement year. The height, weight and BMI percentile must be from the same data source. Either of the following meets criteria for BMI percentile: BMI percentile documented as a value (e.g., 85th percentile) OR BMI percentile plotted on an age-growth chart. Only evidence of the BMI percentile or BMI percentile plotted on an age-growth chart meets HEDIS criteria.

Rev. 04/14
Rev. 07/14
Rev. 04/15
Rev. 10/15
Rev. 01/16
Rev. 01/18
Rev. 04/18

ICD-10 Codes to Identify BMI

Body Mass Index (BMI) – Pediatrics		
BMI Code	BMI Percentile	Range
Z68.51	< 5 th percentile for age	Underweight
Z68.52	5 th to < 85 th percentile for age	Normal/ Healthy Weight Range
Z68.53	85 th to < 95 th percentile for age	Overweight
Z68.54	≥ 95 th percentile for age	Obese

FFS EPSDT providers are encouraged to report BMI diagnosis codes with the EPSDT preventive health codes. Do not point the preventive health visit code to the BMI diagnosis code because this will cause the FFS claim to deny payment.

Rev. 10/13
Rev. 01/14
Rev. 10/14
Rev. 07/16
Rev. 04/23

Counseling for Nutrition

Guidelines

The [Dietary Guidelines for Americans, 2020-2025 and Online Materials](https://www.dietaryguidelines.gov/resources/2020-2025-dietary-guidelines-online-materials) (<https://www.dietaryguidelines.gov/resources/2020-2025-dietary-guidelines-online-materials>) comprise core principles to help people, ages 2 years and older, develop healthy lifestyles based on individual needs, likes, and dislikes related to both eating and physical activity. The Dietary Guidelines recommend that children and adolescents consume a healthy eating pattern that accounts for all foods and beverages within an appropriate calorie level. A healthy eating pattern includes a variety of vegetables from all of the subgroups, fruits, grains, fat-free or low-fat dairy, a variety of protein foods and oils. A healthy eating pattern limits saturated fats and *trans* fats, added sugars, and sodium.

Documentation for HEDIS compliance – Counseling for Nutrition

Documentation (in the medical record) of counseling for nutrition must include a statement indicating the date and at least one of the following:

- Discussion of current nutrition behaviors (e.g., eating habits, dieting behaviors).
- Checklist indicating nutrition was addressed.
- Counseling or referral for nutrition education.
- Member received educational materials on nutrition during a face-to-face visit.
- Anticipatory guidance for nutrition.
- Weight or obesity counseling.

ICD-10 Code to Identify Counseling for Nutrition

- Z71.3 – nutritional counseling

FFS EPSDT providers are encouraged to report the nutrition diagnosis code with the EPSDT preventive health codes. Do not point the preventive health visit code to the nutrition and physical activity counseling diagnosis code because this will cause the FFS claim to deny payment.

Counseling for Physical Activity

Guidelines

The [Physical Activity Guidelines for Americans](https://health.gov/sites/default/files/2019-09/Physical_Activity_Guidelines_2nd_edition.pdf) (https://health.gov/sites/default/files/2019-09/Physical_Activity_Guidelines_2nd_edition.pdf) describe the types and amounts of physical activity that offer substantial health benefits for children and adolescents (ages 6 to 17) and adults. The *Physical Activity Guidelines for Americans* complement the *Dietary Guidelines for Americans*, and together the documents provide guidance on the importance of being physically active and eating healthy foods to promote health and reduce the risk of chronic diseases. The Physical Activity Guidelines recommend that children and adolescents have 60 minutes (1 hour) or more of physical activity each day. Physical activity includes aerobic/endurance activities (to increase cardiorespiratory fitness), muscle-strengthening (resistance training which builds strong muscles), and bone-strengthening (weight-bearing or weight-loading activities which promote bone growth and strength).

Rev. 10/13
Rev. 01/14
Rev. 10/14
Rev. 07/16
Rev. 04/23

Documentation for HEDIS Compliance – Counseling for Physical Activity

Documentation (in the medical record) of counseling for physical activity must include a statement indicating the date and at least one of the following:

- Discussion of current physical activity behaviors (e.g., exercise routine, participation in sports activities, exam for sports participation).
- Checklist indicating physical activity was addressed.
- Counseling or referral for physical activity.
- Member received educational materials on physical activity during a face-to-face visit.
- Anticipatory guidance specific to the child’s physical activity.
- Weight or obesity counseling.

The following notations or examples of documentation are not compliant with HEDIS requirements:

Rev. 10/16
Rev. 01/18

- **BMI**
 - No BMI percentile documented in medical record or plotted on age-growth chart.
 - Notation of BMI value only.
 - Notation of height and weight only.
- **Nutrition**
 - No counseling/education on nutrition and diet.
 - Counseling/education before or after the HEDIS measurement year
 - Notation of “health education” or “anticipatory guidance” without specific mention of nutrition.
 - A physical exam finding or observation alone (e.g., well-nourished) is not compliant because it does not indicate counseling for nutrition.
 - Documentation related to a member’s “appetite” does not meet criteria
- **Physical Activity**
 - No counseling/education on physical activity.
 - Notation of “cleared for gym class” along without documentation of a discussion.
 - Counseling/education before or after the HEDIS measurement year.
 - Notation of “health education” or “anticipatory guidance” without specific mention of physical activity.
 - Notation of anticipatory guidance related solely to safety (e.g., wears helmet or water safety) without specific mention of physical activity recommendations.
 - Notation solely related to screen time (computer or television) without specific mention of physical activity.

Services may be rendered during a visit other than a well-child visit. These services count if the specified documentation is present, regardless of the primary intent of the visit. **Services specific to the assessment or treatment of an acute or chronic condition do not count toward the “Counseling for Nutrition” and “Counseling for Physical Activity” measures.**

APPENDIX G

EPSDT Program Required Equipment Form

- Scale for Weighing Infants **present**
- Scale for Weighing Children and Adolescents **present**
- Measuring Board or Device for measuring Length or Height in the recumbent position for Infants and Children up to the age of two (2) **present**
- Measuring Board or Device for measuring Height in the vertical position for children who are over two (2) years old **present**
- Blood Pressure apparatus with infant, child, and adult cuffs **present**
- Screening audiometer **present**
- Centrifuge or other device for measuring hematocrit or hemoglobin may be **present**
- Eye charts appropriate for age of the child **present**
- Developmental/Behavioral, Alcohol/Substance Abuse and Depression screening tools and supplies **present** (*The required developmental screenings at ages 9 months, 18 months, and 30 months **must** be accomplished using one or more of the recommended standardized developmental screening tools specified in Section 902.2*)
- Vaccines and immunization administration supplies **present**
- Lab supplies for appropriate lab tests/screenings **present**
- Ophthalmoscope and Otoscope **present**

The information supplied in this document is true, accurate and complete and is hereby released to the Georgia Department of Community Health, Division of Medicaid, for purpose of enrolling in the EPSDT Program. I understand that falsification, omission or misrepresentation of any information in this enrollment document will result in a denial of enrollment, the closure of current enrollment, and the denial of future enrollment request, and may be punishable by criminal, civil or other administrative actions. I understand that my completion of this form certifies that I have the necessary equipment as listed in Part II- Policies and Procedures Manual for the EPSDT Program.

Provider Name

Date

Provider Title

Provider/Confirmation Number

APPENDIX H

Georgia Families

Rev. 01/11
 Rev. 10/12
 Rev. 04/14
 Rev. 07/17
 Rev. 10/18
 Rev. 07/21

Georgia Families® (GF) is a statewide program designed to deliver health care services to members of Medicaid, PeachCare for Kids®, and Planning for Healthy Babies® (P4HB) recipients. The program is a partnership between the Department of Community Health (DCH) and private care management organizations (CMOs). By providing a choice of health plans, Georgia Families allows members to select a health care plan that fits their needs.

It is important to note that GF is a full-risk program; this means that the three CMOs licensed in Georgia to participate in GF are responsible and accept full financial risk for providing and authorizing covered services. This also means a greater focus on case and disease management with an emphasis on preventative care to improve individual health outcomes.

The three licensed CMOs:

 Amerigroup Community Care 1-800-454-3730 www.amerigroup.com	 Peach State Health Plan 866-874-0633 www.pshpgeorgia.com	 CareSource 1-855-202-1058 www.caresource.com
---	---	--

Children, parent/caretaker with children, pregnant women and women with breast or cervical cancer on Medicaid, as well as children enrolled in PeachCare for Kids® are eligible to participate in Georgia Families. Additionally, Planning for Healthy Babies® (P4HB) recipients receive services through Georgia Families® (GF). Children in foster care or receiving adoption assistance and certain youths committed to juvenile justice are enrolled in Georgia Families 360°.

Eligibility Categories for Georgia Families:

Included Populations	Excluded Populations
Parent/Caretaker with Children	Aged, Blind and Disabled
Transitional Medicaid	Nursing home
Pregnant Women (Right from the Start Medicaid – RSM)	Long-term care (Waivers, SOURCE)
Children (Right from the Start Medicaid – RSM)	Federally Recognized Indian Tribe
Children (newborn)	Georgia Pediatric Program (GAPP)
Women Eligible Due to Breast and Cervical Cancer	Hospice
PeachCare for Kids®	Children’s Medical Services program
Parent/Caretaker with Children	Medicare Eligible
Children under 19	Supplemental Security Income (SSI) Medicaid

Included Populations	Excluded Populations
Women’s Health Medicaid (WHM)	Medically Needy
Refugees	Recipients enrolled under group health plans
Planning for Healthy Babies®	Individuals enrolled in a Community Based Alternatives for Youths (CBAY)
Resource Mothers Outreach	

Medicaid and PeachCare for Kids® members will continue to be eligible for the same services they receive through traditional Medicaid and state Value Added Benefits. Members will not have to pay more than they paid for Medicaid co-payments or PeachCare for Kids® premiums. With a focus on health and wellness, the CMOs will provide members with health education and prevention programs giving them the tools needed to live healthier lives. Providers participating in Georgia Families will have the added assistance of the CMOs to educate members about accessing care, referrals to specialists, member benefits, and health and wellness education. **All three CMOs are State-wide.**

The Department of Community Health has contracted with three CMOs to provide these services:

- Amerigroup Community Care
- CareSource
- Peach State Health Plan

Members can contact Georgia Families for assistance to determine which program best fits their family’s needs. If members do not select a plan, Georgia Families will select a health plan for them.

Members can visit the Georgia Families Web site at www.georgia-families.com or call 1-800-GA-ENROLL (1-888-423-6765) to speak to a representative who can give them information about the CMOs and the health care providers.

The following categories of eligibility are included and excluded under Georgia Families:

Included Categories of Eligibility (COE):

COE	DESCRIPTION
104	LIM – Adult
105	LIM – Child
118	LIM – 1st Yr Trans Med Ast Adult
119	LIM – 1st Yr Trans Med Ast Child
122	CS Adult 4 Month Extended
123	CS Child 4 Month Extended
135	Newborn Child
170	RSM Pregnant Women
171	RSM Child
180	P4HB Inter Pregnancy Care
181	P4HB Family Planning Only
182	P4HB ROMC - LIM

Included Categories of Eligibility (COE) continued:	
COE	DESCRIPTION
183	P4HB ROMC - ABD
194	RSM Expansion Pregnant Women
195	RSM Expansion Child < 1 Yr
196	RSM Expn Child w/DOB <= 10/1/83
197	RSM Preg Women Income < 185 FPL
245	Women's Health Medicaid
471	RSM Child
506	Refugee (DMP) – Adult
507	Refugee (DMP) – Child
508	Post Ref Extended Med – Adult
509	Post Ref Extended Med – Child
510	Refugee MAO – Adult
511	Refugee MAO – Child
571	Refugee RSM - Child
595	Refugee RSM Exp. Child < 1
596	Refugee RSM Exp Child DOB <= 10/01/83
790	Peachcare < 150% FPL
791	Peachcare 150 – 200% FPL
792	Peachcare 201 – 235% FPL
793	Peachcare > 235% FPL
835	Newborn
836	Newborn (DFACS)
871	RSM (DHACS)
876	RSM Pregnant Women (DHACS)
894	RSM Exp Pregnant Women (DHACS)
895	RSM Exp Child < 1 (DHACS)
897	RSM Pregnant Women Income > 185% FPL (DHACS)
898	RSM Child < 1 Mother has Aid = 897 (DHACS)
918	LIM Adult
919	LIM Child
920	Refugee Adult
921	Refugee Child

Excluded Categories of Eligibility (COE):

COE	DESCRIPTION
124	Standard Filing Unit – Adult
125	Standard Filing Unit – Child
131	Child Welfare Foster Care
132	State Funded Adoption Assistance
147	Family Medically Needy Spend down
148	Pregnant Women Medical Needy Spend down
172	RSM 150% Expansion
180	Interconceptional Waiver
210	Nursing Home – Aged
211	Nursing Home – Blind
212	Nursing Home – Disabled
215	30 Day Hospital – Aged
216	30 Day Hospital – Blind
217	30 Day Hospital – Disabled
218	Protected Med/1972 Cola - Aged
219	Protected Med/1972 Cola – Blind
220	Protected Med/1972 Cola - Disabled
221	Disabled Widower 1984 Cola - Aged
222	Disabled Widower 1984 Cola – Blind
223	Disabled Widower 1984 Cola – Disabled
224	Pickle - Aged
225	Pickle – Blind
226	Pickle – Disabled
227	Disabled Adult Child - Aged

Excluded Categories of Eligibility (COE) continued:

COE	DESCRIPTION
229	Disabled Adult Child – Disabled
230	Disabled Widower Age 50-59 – Aged
231	Disabled Widower Age 50-59 – Blind
232	Disabled Widower Age 50-59 – Disabled
233	Widower Age 60-64 – Aged
234	Widower Age 60-64 – Blind
235	Widower Age 60-64 – Disabled
236	3 Mo. Prior Medicaid – Aged
237	3 Mo. Prior Medicaid – Blind
238	3 Mo. Prior Medicaid – Disabled
239	Abd Med. Needy Defacto – Aged
240	Abd Med. Needy Defacto – Blind
241	Abd Med. Needy Defacto – Disabled
242	Abd Med Spend down – Aged
243	Abd Med Spend down – Blind
244	Abd Med Spend down – Disabled
246	Ticket to Work
247	Disabled Child – 1996
250	Deeming Waiver
251	Independent Waiver
252	Mental Retardation Waiver
253	Laurens Co. Waiver
254	HIV Waiver
255	Cystic Fibrosis Waiver

Excluded Categories of Eligibility (COE) continued:

COE	DESCRIPTION
259	Community Care Waiver
280	Hospice – Aged
281	Hospice – Blind
282	Hospice – Disabled
283	LTC Med. Needy Defacto – Aged
284	LTC Med. Needy Defacto –Blind
285	LTC Med. Needy Defacto – Disabled
286	LTC Med. Needy Spend down – Aged
287	LTC Med. Needy Spend down – Blind
288	LTC Med. Needy Spend down – Disabled
289	Institutional Hospice – Aged
290	Institutional Hospice – Blind
291	Institutional Hospice – Disabled
301	SSI – Aged
302	SSI – Blind
303	SSI – Disabled
304	SSI Appeal – Aged
305	SSI Appeal – Blind
306	SSI Appeal – Disabled
307	SSI Work Continuance – Aged
309	SSI Work Continuance – Disabled
308	SSI Work Continuance – Blind
315	SSI Zebley Child
321	SSI E02 Month – Aged

Excluded Categories of Eligibility (COE) continued:

COE	DESCRIPTION
322	SSI E02 Month – Blind
323	SSI E02 Month – Disabled
387	SSI Trans. Medicaid – Aged
388	SSI Trans. Medicaid – Blind
389	SSI Trans. Medicaid – Disabled
410	Nursing Home – Aged
411	Nursing Home – Blind
412	Nursing Home – Disabled
424	Pickle – Aged
425	Pickle – Blind
426	Pickle – Disabled
427	Disabled Adult Child – Aged
428	Disabled Adult Child – Blind
429	Disabled Adult Child – Disabled
445	N07 Child
446	Widower – Aged
447	Widower – Blind
448	Widower – Disabled
460	Qualified Medicare Beneficiary
466	Spec. Low Inc. Medicare Beneficiary
575	Refugee Med. Needy Spend down
660	Qualified Medicare Beneficiary
661	Spec. Low Income Medicare Beneficiary
662	Q11 Beneficiary

Excluded Categories of Eligibility (COE) continued:	
COE	DESCRIPTION
663	Q12 Beneficiary
664	Qua. Working Disabled Individual
815	Aged Inmate
817	Disabled Inmate
870	Emergency Alien – Adult
873	Emergency Alien – Child
874	Pregnant Adult Inmate
915	Aged MAO
916	Blind MAO
917	Disabled MAO
983	Aged Medically Needy
984	Blind Medically Needy
985	Disabled Medically Needy

HEALTH CARE PROVIDERS

For information regarding the participating health plans (enrollment, rates, and procedures), please call the numbers listed below.

Prior to providing services, you should contact the member’s health plan to verify eligibility, PCP assignment and covered benefits. You should also contact the health plan to check prior authorizations and submit claims.

Amerigroup Community Care	CareSource	Peach State Health Plan
800-454-3730 (general information) www.amerigroup.com	1-855-202-1058 www.careSource.com/GeorgiaMedicaid	866-874-0633 (general information) 866-874-0633 (claims) 800-704-1483 (medical management) www.pshpgeorgia.com

Registering immunizations with GRITS:

If you are a Vaccine for Children (VFC) provider, please continue to use the GRITS (Georgia Immunization Registry) system for all children, including those in Medicaid and PeachCare for Kids®, fee-for-service, and managed care.

Important tips for the provider to know/do when a member comes in:

Understanding the process for verifying eligibility is now more important than ever. You will need to determine if the patient is eligible for Medicaid/PeachCare for Kids® benefits and if they are enrolled in a Georgia Families health plan. Each plan sets its own medical management and referral processes. Members will have a new identification card and primary care provider assignment.

You may also contact GAINWELL TECHNOLOGIES at 1-800-766-4456 (statewide) or www.mmis.georgia.gov for information on a member's health plan.

Use of the Medicaid Management Information System (MMIS) web portal:

The call center and web portal will be able to provide you information about a member's Medicaid eligibility and health plan enrollment. GAINWELL TECHNOLOGIES will **not** be able to assist you with benefits, claims processing or prior approvals for members assigned to a Georgia Families health plan. You will need to contact the member's plan directly for this information.

Participating in a Georgia Families' health plan:

Each health plan will assign provider numbers, which will be different from the provider's Medicaid provider number and the numbers assigned by other health plans.

Billing the health plans for services provided:

For members who are in Georgia Families, you should file claims with the member's health plan.

If a claim is submitted to GAINWELL TECHNOLOGIES in error:

GAINWELL TECHNOLOGIES will deny the claim with a specific denial code. Prior to receiving this denial, you may go ahead and submit the claim to the member's health plan.

Credentialing

Effective August 1, 2015, Georgia's Department of Community Health (DCH) implemented a NCQA certified Centralized Credentialing Verification Process utilizing a Credentialing Verification Organization (CVO). This functionality has been added to the Georgia Medicaid Management Information System (GAMMIS) website (www.MMIS.georgia.gov) and has streamlined the time frame that it takes for a provider to be fully credentialed.

Credentialing and recredentialing services is provided for Medicaid providers enrolled in Georgia Families and/or the Georgia Families 360° program.

This streamlined process results in administrative simplification thereby preventing inconsistencies, as well as the need for a provider to be credentialed or recredentialled multiple times.

The CVO's one-source application process:

- Saves time
- Increases efficiency
- Eliminates duplication of data needed for multiple CMOs
- Shortens the time period for providers to receive credentialing and recredentialing decisions

The CVO will perform primary source verification, check federal and state databases, obtain information from Medicare's Provider Enrollment Chain Ownership System (PECOS), check required medical malpractice insurance, confirm Drug Enforcement Agency (DEA) numbers, etc. A Credentialing Committee will render a decision regarding the provider's credentialing status. Applications that contain all required credentialing and recredentialing materials at the time of submission will receive a decision within 45 calendar days. Incomplete applications that do not contain all required credentialing documents will be returned to the provider with a request to supplement all missing materials. Incomplete applications may result in a delayed credentialing or recredentialing decision. The credentialing decision is provided to the CMOs. GAINWELL TECHNOLOGIES provider reps will provide training and assistance as needed. Providers may contact GAINWELL TECHNOLOGIES for assistance with credentialing and recredentialing by dialing 1-800-766-4456.

Assignment of separate provider numbers by all of the health plans:

Each health plan will assign provider numbers, which will be different from the provider's Medicaid provider number and the numbers assigned by other health plans.

Billing the health plans for services provided:

For members who are in Georgia Families, you should file claims with the member's health plan.

If a claim is submitted to GAINWELL TECHNOLOGIES in error:

GAINWELL TECHNOLOGIES will deny the claim with a specific denial code. Prior to receiving this denial, you may go ahead and submit the claim to the member's health plan.

Receiving payment:

Claims should be submitted to the member's health plan. Each health plan has its own claims processing, and you should consult the health plan about their payment procedures.

Health plans payment of clean claims:

Each health plan (and subcontractors) has its own claims processing and payment cycles. The claims processing and payment timeframes are as follows:

Amerigroup Community Care	CareSource	Peach State Health Plan
<p>Amerigroup runs claims cycles twice each week (on Monday and Thursday) for clean claims that have been adjudicated.</p> <p>Monday Claims run: Checks mailed on Tuesday. Providers enrolled in ERA/EFT receive the ACH on Thursday.</p> <p>Thursday Claims run: Checks mailed on Wednesday. Providers enrolled in ERA/EFT receive the ACH on Tuesday.</p> <p>Dental: Checks are mailed weekly on Thursday for clean claims.</p> <p>Vision: Checks are mailed weekly on Wednesday for clean claims (beginning June 7th)</p> <p>Pharmacy: Checks are mailed to pharmacies weekly on Friday (except when a holiday falls on Friday, then mailed the next business day)</p>	<p>CareSource runs claims cycles twice each week on Saturdays and Tuesdays for <u>clean</u> claims that have been adjudicated.</p> <p><u>Pharmacy:</u> Payment cycles for pharmacies is weekly on Wednesdays.</p>	<p>Peach State has two weekly claims payment cycles per week that produces payments for clean claims to providers on Monday and Wednesday.</p> <p>For further information, please refer to the Peach State website, or the Peach State provider manual.</p>

How often can a patient change his/her PCP?

Amerigroup Community Care	CareSource	Peach State Health Plan
Anytime	Members can change their PCP one (1) time per month. However, members can change their PCP at any time under extenuating circumstances such as: <ul style="list-style-type: none"> • Member requests to be assigned to a family member’s PCP • PCP does not provide the covered services a member seeks due to moral or religious objections • PCP moves, retires, etc. 	Within the first 90 days of a member’s enrollment, he/she can change PCP monthly. If the member has been with the plan for 90 days or longer, the member can change PCPs once every six months. There are a few exclusions that apply and would warrant an immediate PCP change.

Once the patient requests a PCP change, how long it takes for the new PCP to be assigned:

Amerigroup Community Care	CareSource	Peach State Health Plan
Next business day	PCP selections are updated in CareSource’s systems daily.	PCP changes made before the 24 th day of the month and are effective for the current month. PCP changes made after the 24 th day of the month are effective for the first of the following month.

PHARMACY

Georgia Families does provide pharmacy benefits to members. Check with the member’s health plan about who to call to find out more about enrolling to provide pharmacy benefits, including information about their plans reimbursement rates, specific benefits that are available, including prior approval requirements.

To request information about contracting with the health plans, you can call the CMOs provider enrollment services.

Amerigroup Community Care	CareSource	Peach State Health Plan
800-454-3730 https://providers.amerigroup.com/pages/ga-2012.aspx	844-441-8024 https://cvs.az1.qualtrics.com/jfe/form/SV_cvyY0ohqT2VXYod	866-874-0633 www.pshpgeorgia.com

All providers must be enrolled as a Medicaid provider to be eligible to contract with a health plan to provide services to Georgia Families members.

The CMO Pharmacy Benefit Managers (PBM) and the Bin Numbers, Processor Control Numbers and Group Numbers are:

Health Plan	PBM	BIN #	PCN #	GROUP #	Helpdesk
Amerigroup Community Care	IngenioRx	020107	HL	WKJA	1-833-235-2031
CareSource	Express Scripts (ESI)	003858	MA	RXINN01	1-800-416-3630
Peach State Health Plan	CVS	004336	MCAIDADV	RX5439	1-844-297-0513

If a patient does not have an identification card:

Providers can check the enrollment status of Medicaid and PeachCare for Kids® members through GAINWELL TECHNOLOGIES by calling 1-800-766-4456 or going to the web portal at www.mmis.georgia.gov. GAINWELL TECHNOLOGIES will let you know if the member is eligible for services and the health plan they are enrolled in. You can contact the member’s health plan to get the member’s identification number.

Use of the member’s Medicaid or PeachCare for Kids® identification number to file a pharmacy claim:

Amerigroup Community Care	CareSource	Peach State Health Plan
No, you will need the member’s health plan ID number	Yes, you may also use the health plan ID number.	Yes

Health plans preferred drug list, prior authorization criteria, benefit design, and reimbursement rates:

Each health plan sets their own procedures, including preferred drug list, prior authorization criteria, benefit design, and reimbursement rates.

Will Medicaid cover prescriptions for members that the health plans do not?

No, Medicaid will not provide a “wrap-around” benefit for medications not covered or approved by the health plan. Each health plan will set its own processes for determining medical necessity and appeals.

Who to call to request a PA:

Amerigroup Community Care	CareSource	Peach State Health Plan
1 (800) 454-3730	1 (855) 202-1058 1 (866) 930-0019 (fax)	1 (866) 399-0929

APPENDIX I



Information for Providers Serving Medicaid Members in the Georgia Families 360SM Program

Rev. 04/14
Rev. 07/18

Georgia Families 360SM, the state's managed care program for children, youth, and young adults in Foster Care, children and youth receiving Adoption Assistance, as well as select youth in the juvenile justice system, launched Monday, March 3, 2014. Amerigroup Community Care is the single Care Management Organization (CMO) that will be managing this population.

Amerigroup is responsible, through its provider network, for coordinating all DFCS, DJJ required assessments and medically necessary services for children, youth and young adults who are eligible to participate in the Georgia Families 360SM Program. Amerigroup will coordinate all medical/dental/trauma assessments for youth upon entry into foster care or juvenile justice (and as required periodically).

Georgia Families 360SM Every member in Georgia Families 360SM is assigned a Care Coordinator who works closely with them to ensure access to care and ensure that appropriate, timely, and trauma informed care is provided for acute conditions as well as ongoing preventive care. This ensures that all medical, dental, and behavioral health issues are addressed. Members also have a medical and dental home to promote consistency and continuity of care. The medical and dental homes coordinate care and serve as a place where the child is known over time by providers who can provide holistic care. DFCS, DJJ, foster parents, adoptive parents and other caregivers are involved in the ongoing health care plans to ensure that the physical and behavioral health needs of these populations are met.

Electronic Health Records (EHRs) are being used to enhance effective delivery of care. The EHRs can be accessed by Amerigroup, physicians in the Amerigroup provider network, and DCH sister agencies, including the DFCS, regardless of where the child lives, even if the child experiences multiple placements. Ombudsman and advocacy staff are in place at both DCH and Amerigroup to support caregivers and members, assisting them in navigating the health care system. Additionally, medication management programs are in place to focus on appropriate monitoring of the use of psychotropic medications, to include ADD/ADHD as well as other behavioral health prescribed medications.

Providers can obtain additional information by contacting the Provider Service Line at 1-800-454-3730 or by contacting their Provider Relations representative.

To learn more about DCH and its dedication to A Healthy Georgia, visit www.dch.georgia.gov

APPENDIX J

Preventive Oral Health: Fluoride Varnish

Fluoride varnish acts to retard, arrest, and reverse the caries process. The teeth absorb the fluoride varnish, strengthening the enamel and helping prevent cavities. It is not a substitute for fluoridated water or toothpaste.

Rev. 07/10
Rev. 07/11
Rev. 04/15
Rev. 10/15
Rev. 01/16
Rev. 04/16
Rev. 10/16
Rev. 10/19
Rev. 04/24

Once teeth are present, the application of fluoride varnish is required and may be applied every 3-6 months in the primary care or dental office for children between the ages of 6 months and 5 years.

[Recommendation: Prevention of Dental Caries in Children Younger Than 5 Years: Screening and Interventions | United States Preventive Services Taskforce \(uspreventiveservicestaskforce.org\)](https://uspreventiveservicestaskforce.org)

Indications for fluoride use are noted in the 2020 AAP Clinical Report “[Fluoride Use in Caries Prevention in the Primary Care Setting | Pediatrics | American Academy of Pediatrics \(aap.org\)](https://publications.aap.org/pediatrics/article/146/6/e2020034637/33536/fluoride-use-in-caries-prevention-in-the-primary) <https://publications.aap.org/pediatrics/article/146/6/e2020034637/33536/fluoride-use-in-caries-prevention-in-the-primary>

Documentation: Evidence that fluoride varnish was applied once between the ages of 6 months and 5 years OR evidence that the provider addressed the fluoride varnish requirement and/or its importance with the parent.

Effective April 1, 2024, Health Check providers are allowed reimbursement for the application of topical fluoride varnish. Providers should report cpt code 99188 with the EP modifier and POS 99.

99188 - Current Procedural Terminology (CPT) Code
D1206 - Current Dental Terminology (CDT) Code

Effective January 1, 2015, the application of topical fluoride varnish by a physician or other qualified health care professional may be billed with the new CPT code 99188. This applies to providers enrolled in and filing claims under GA Medicaid programs 430, 431, and 740. Only providers enrolled in and filing claims under GA Medicaid programs 430, 431, 450, and 740 may bill Code D1206 Fluoride Varnish (eff. 1/1/2010).

Note:

- Dentists: under category of service 450
- Physicians: under category of service 430
- Physician Assistants (PA): under category of service 431
- Nurse Practitioners: under category of service 740

Providers may not bill for an Evaluation and Management (E/M) visit in addition to billing for the application of fluoride varnish, if the sole purpose of the visit was to apply the fluoride varnish. In this instance, the provider may bill for the fluoride varnish code only.

For more information including the payment rate for this service, please see the Part II Policies and Procedures Manual for Dental Services.

RESOURCES (not mandatory to use):

Rev. 07/10
Rev. 07/11
Rev. 04/15
Rev. 10/15
Rev. 01/16
Rev. 04/16
Rev. 10/16
Rev. 10/19

Smiles for Life [Oral Health Risk Assessment Tool](#):

Caries risk assessments:

[ADA Caries Risk Assessment Form \(Age 0-6\)](#)

[ADA Caries Risk Assessment Form \(Age >6\)](#)

Smiles for Life Online trainings:

[Child Oral Health](#) (Course 2)

[Caries Risk Assessment, Fluoride Varnish and Counseling \(Course 6\)](#) (Course 6)

Oral Health Professional Websites: American Dental Association (ADA): <http://www.ada.org>

American Academy of Pediatric Dentistry (AAPD): <http://www.aapd.org>

Parent Handouts: 'For The Dental Patient' by the ADA freely available for download and photocopy at <http://www.ada.org/993.aspx>

*Patients at risk for caries include those with: insufficient sources of dietary fluoride; high carbohydrate diets; caretakers who transmit decay-causing bacteria to children via their saliva; areas of tooth decalcification; reduced salivary flow; and poor oral hygiene. AAP training course also includes “children from low socioeconomic and ethnocultural groups.”

APPENDIX K

EPSDT HIPAA Referral Code Examples

Rev. 10/14

1. Child has come in for an EPSDT Interperiodic Hearing Screen and the provider finds that the child has an ear infection. The provider treats the child for the ear infection at the time of the interperiodic visit and requests a follow up appointment with him in two weeks. What EPSDT referral code should be documented?
 - A. EPSDT Referral Code: **S2**

2. Child has come in for an EPSDT Screen and has experienced complications with diabetes since birth. The provider treats the child for the diabetes complications at the time of the preventive health visit and does not request a follow up appointment. What EPSDT referral code should be documented?
 - A. EPSDT Referral Code: **NU**

3. Child has come in for an EPSDT Screen and during the screen, the mother informs the provider that the child has behavior problems. The provider refers the child for further diagnostic testing within two weeks with a Diagnostic and Behavioral Center. What EPSDT referral code should be documented?
 - A. EPSDT Referral Code: **ST**

4. Child has come in for EPSDT Screen and the provider finds that the child has some developmental problems. The provider refers the child for further diagnostic testing with a Developmental and Behavioral Center. Mom refuses the Developmental and Behavioral appointment. What EPSDT referral code should be documented?
 - A. EPSDT Referral Code: **AV**

APPENDIX L

NEW CMS 1500 CLAIM FORM (version 02/12) - SAMPLE

Effective May 1, 2015, paper claims are no longer accepted by Gainwell Technologies. As part of the Georgia Paperless Initiative, providers are required to submit CMS 1500 claims electronically over the GAMMIS web portal. For more information regarding the Paperless Initiative, please access the web portal and review all related Banner Messages.



NEW CMS 1500 CLAIM FORM (version 02/12)
Effective April 1, 2014

HEALTH INSURANCE CLAIM FORM

APPROVED BY NATIONAL UNIFORM CLAIM COMMITTEE (NUCC) ©2012

<input type="checkbox"/> MEDICARE <input type="checkbox"/> MEDICAID <input type="checkbox"/> TRICARE <input type="checkbox"/> CHAMPVA (Member/Spouse) <input type="checkbox"/> GROUP HEALTH PLAN (Self/Spouse/Child/Other) <input type="checkbox"/> FICA (Not Listed) (Self/Spouse) <input type="checkbox"/> OTHER (Self/Spouse)										<input type="checkbox"/> PPO <input type="checkbox"/> HMO													
1. PATIENT'S NAME (Last Name, First Name, Middle Initial)				3. PATIENT'S BIRTH DATE (MM/DD/YY) SEX (M/F)		4. INSURED'S ID NUMBER (For Programs in Item 1)				14. INSURED'S ID NUMBER (For Programs in Item 1)													
5. PATIENT'S ADDRESS (No. Street) CITY STATE ZIP CODE TELEPHONE (Include Area Code)				6. PATIENT RELATIONSHIP TO INSURED (Self/Spouse/Child/Other)		7. INSURED'S ADDRESS (No. Street) CITY STATE ZIP CODE TELEPHONE (Include Area Code)				15. INSURED'S POLICY GROUP OR FICA NUMBER													
8. OTHER INSURED'S NAME (Last Name, First Name, Middle Initial)				10. THIS PATIENT'S CONDITION RELATED TO:		9. INSURED'S DATE OF BIRTH (MM/DD/YY) SEX (M/F)				16. OTHER CLAIM ID (designated by NUCC)													
9. OTHER INSURED'S POLICY OR GROUP NUMBER				7. EMPLOYMENT (Current or Previous) YES/NO		10. INSURED'S DATE OF DEATH (MM/DD/YY) SEX (M/F)				17. INSURANCE PLAN NAME OR PROGRAM NAME													
11. RESERVED FOR NUCC USE				8. AUTO ACCIDENT? YES/NO		11. OTHER CLAIM ID (designated by NUCC)				18. IS THERE ANOTHER HEALTH BENEFIT PLAN? YES/NO													
12. RESERVED FOR NUCC USE				9. OTHER ACCIDENT? YES/NO		12. INSURANCE PLAN NAME OR PROGRAM NAME				19. INSURED'S OR AUTHORIZED PERSON'S SIGNATURE & authorize payment of medical benefits to the undersigned physician or supplier for services described below.													
13. INSURANCE PLAN NAME OR PROGRAM NAME				10. CLAIM CODES (Designated by NUCC)		13. IS THERE ANOTHER HEALTH BENEFIT PLAN? YES/NO				20. PATIENT'S OR AUTHORIZED PERSON'S SIGNATURE & authorize payment of medical benefits to the undersigned physician or supplier for services described below.													
<p>12. PATIENT'S OR AUTHORIZED PERSON'S SIGNATURE & authorize the release of any medical or other information necessary to process this claim. It also request payment of governmental benefits either to itself or to the party also accept(s) assignment below.</p> <p>SIGNED: _____ DATE: _____</p>																							
14. DATE OF CURRENT CLAIM'S DUE DATE OF PAYMENT (MM/DD/YY)				15. OTHER BILL (MM/DD/YY)				16. DATE SYSTEM DISABLED BY WORK IN CURRENT OCCUPATION (FROM/TO) (MM/DD/YY)															
17. NAME OF REFERRING PROVIDER OR OTHER SOURCE (Y/N) (Y/N)				18. HOSPITALIZATION DATE(S) RELATED TO CURRENT SERVICES (FROM/TO) (MM/DD/YY)				19. OUTSIDE LAB? CHARGES? YES/NO															
19. ADDITIONAL CLAIM INFORMATION (Designated by NUCC)																							
20. PROVIDER OR NATURE OF ILLNESS OR INJURY (Please tick in service line below) (I/O Ref.)																							
<table border="1"> <tr> <td>A. _____</td> <td>B. _____</td> <td>C. _____</td> <td>D. _____</td> <td>E. _____</td> <td>F. _____</td> <td>G. _____</td> <td>H. _____</td> <td>I. _____</td> <td>J. _____</td> <td>K. _____</td> <td>L. _____</td> </tr> </table>												A. _____	B. _____	C. _____	D. _____	E. _____	F. _____	G. _____	H. _____	I. _____	J. _____	K. _____	L. _____
A. _____	B. _____	C. _____	D. _____	E. _____	F. _____	G. _____	H. _____	I. _____	J. _____	K. _____	L. _____												
21. A. DATES OF SERVICE (MM/DD/YY - MM/DD/YY) B. PLACE OF SERVICE (FAC) C. PROCEDURES, SERVICES, OR SUPPLIES (ICD-9-CM, ICD-9-CM-PCS, ICD-9-CM-PCS) D. CHARGES POINTERS E. CHARGES F. CODES (ICD-9-CM, ICD-9-CM-PCS) G. ICD-9-CM-PCS H. ICD-9-CM-PCS I. ICD-9-CM-PCS J. ICD-9-CM-PCS K. ICD-9-CM-PCS L. ICD-9-CM-PCS M. ICD-9-CM-PCS N. ICD-9-CM-PCS O. ICD-9-CM-PCS P. ICD-9-CM-PCS Q. ICD-9-CM-PCS R. ICD-9-CM-PCS S. ICD-9-CM-PCS T. ICD-9-CM-PCS U. ICD-9-CM-PCS V. ICD-9-CM-PCS W. ICD-9-CM-PCS X. ICD-9-CM-PCS Y. ICD-9-CM-PCS Z. ICD-9-CM-PCS																							
22. FEDERAL TAX ID NUMBER SSN EIN 23. PATIENT'S ACCOUNT NO. 24. ACCIDENT & OCCASION? YES/NO 25. TOTAL CHARGE 26. AMOUNT PAID 27. PAID FOR NUCC USE																							
28. SIGNATURE OF PHYSICIAN OR SUPPLIER INCLUDING DEGREE OR CREDENTIALS (I certify that the statements on the reverse apply to this bill and are made a part thereof)				29. SERVICE FACILITY LOCATION INFORMATION				30. BILLING PROVIDER INFO & PII															
SIGNED: _____ DATE: _____				31. NPI				32. NPI															

New CMS 1500 Field Locator Instructions

The following table outlines the **revised changes** on the above CMS 1500 claim form version 02/12:

FLD Location	NEW Change
Header	Replaced 1500 rectangular symbol with black and white two-dimensional QR Code (Quick Response Code)
Header	Added "(NUCC)" after "APPROVED BY NATIONAL UNIFORM CLAIM COMMITTEE."
Header	Replaced "08/05" with "02/12"
Item Number 1	Changed "TRICARE CHAMPUS" to "TRICARE" and changed "(Sponsor's SSN)" to "(ID#/DoD#)."
Item Number 1	Changed "(SSN or ID)" to "(ID#)" under "GROUP HEALTH PLAN"
Item Number 1	Changed "(SSN)" to "(ID#)" under "FECA BLK LUNG."
Item Number 1	Changed "(ID)" to "(ID#)" under "OTHER."
Item Number 8	Deleted "PATIENT STATUS" and content of field. Changed title to " RESERVED FOR NUCC USE. "
Item Number 9b	Deleted "OTHER INSURED's DATE OF BIRTH, SEX." Changed title to " RESERVED FOR NUCC USE. "
Item Number 9c	Deleted "EMPLOYER'S NAME OR SCHOOL." Changed title to " RESERVED FOR NUCC USE. "
Item Number 10d	Changed title from "RESERVED FOR LOCAL USE" to "CLAIM CODES (Designated by NUCC)." Field 10d is being changed to receive Worker's Compensation codes or Condition codes approved by NUCC. FOR DCH/Gainwell Technologies: FLD 10d on the OLD Form CMS 1500 Claim (08/05) will no longer support receiving the Medicare provider ID.
Item Number 11b	Deleted "EMPLOYER'S NAME OR SCHOOL." Changed title to "OTHER CLAIM ID (Designated by NUCC)". Added dotted line in the left-hand side of the field to accommodate a 2-byte qualifier
Item Number 11d	Changed "If yes, return to and complete Item 9 a-d" to "If yes, complete items 9, 9a, and 9d." (Is there another Health Benefit Plan?)
Item Number 14	Changed title to "DATE OF CURRENT ILLNESS, INJURY, OR PREGNANCY (LMP)." Removed the arrow and text in the right-hand side of the field. Added "QUAL." with a dotted line to accommodate a 3-byte qualifier. FOR DCH/Gainwell Technologies: Use Qualifiers: 431 (onset of current illness); 484 (LMP); or 453 (Estimated Delivery Date).
Item Number 15	Changed title from 'IF PATIENT HAS HAD SAME OR SIMILAR ILLNESS. GIVE FIRST DATE' to "OTHER DATE." Added "QUALIFIER." with two dotted lines to accommodate a 3-byte qualifier: 454 (Initial Treatment); 304 (Latest Visit or Consultation); 453 (Acute Manifestation of a

FLD Location	NEW Change
	Chronic Condition); 439 (Accident); 455 (Last X-ray); 471 (Prescription); 090 (Report Start [Assumed Care Date]); 091 (Report End [Relinquished Care Date]); 444 (First Visit or Consultation).
Item Number 17	Added a dotted line in the left-hand side of the field to accommodate a 2-byte qualifier – Used by Medicare for identifiers for provider roles: Ordering, Referring and Supervising. FOR DCH/Gainwell Technologies: Use the following Ordering Provider, Referring, Supervising Qualifiers (effective 4/01/2014): Ordering = DK; Referring = DN or Supervising = DQ.
Item Number 19	Changed title from “ RESERVED FOR LOCAL USE ” to “ADDITIONAL CLAIM INFORMATION (Designated by NUCC).” FOR DCH/Gainwell Technologies: Remove the Health Check logic from field 19 and add it in field 24H.
Item Number 21	Changed instruction after title (Diagnosis or Nature of Illness or Injury) from “(Relate Items 1, 2, 3 or 4 to Item 24E by Line)” to “Relate A-L to service line below (24E).”
Item Number 21	Removed arrow pointing to 24E (Diagnosis Pointer).
Item Number 21	Added “ICD Indicator.” and two dotted lines in the upper right-hand corner of the field to accommodate a 1-byte indicator. Use the highest level of code specificity in FLD Locator 21. Diagnosis Code ICD Indicator - new logic to validate acceptable values (0, 9). ICD-9 diagnoses (CM) codes = value 9; or ICD -10 diagnoses (CM) codes = value 0. (Do not bill ICD 10 code sets before October 1, 2015.)
Item Number 21	Added 8 additional lines for diagnosis codes. Evenly space the diagnosis code lines within the field.
Item Number 21	Changed labels of the diagnosis code lines to alpha characters (A-L).
Item Number 21	Removed the period within the diagnosis code lines
Item Number 22	Changed title from “MEDICAID RESUBMISSION” to “RESUBMISSION.” The submission codes are: 7 (Replacement of prior claim) 8 (Void/cancel of prior claim)
Item Numbers 24A – 24 G (Supplemental Information)	The supplemental information is to be placed in the shaded section of 24A through 24G as defined in each Item Number. FOR DCH/Gainwell Technologies: Item numbers 24A & 24G are used to capture Hemophilia drug units. 24H (EPSDT/Family Planning).
Item Number 30	Deleted “BALANCED DUE.” Changed title to “ RESERVED FOR NUCC USE. ”

FLD Location	NEW Change
Footer	Changed “APPROVED OMB-0938-0999 FORM CMS-1500 (08/05)” to “APPROVED OMB-0938-1197 FORM 1500 (02/12).”

Completion of the Health Insurance Claim Form for EPSDT Services Billed by Fee-for-Service Providers

Review these helpful tips for completing the Health Insurance Claim Form (CMS-1500) for EPSDT Services. See Appendix K for EPSDT HIPAA Referral Code Examples.

Item 9 Other Insured’s Name

Leave blank. EPSDT preventive health screenings are exempt from third party liability. Even if the member has other insurance, you may file Medicaid first for preventive health services.

Item 21 Diagnosis

Enter the applicable ICD indicator to identify which version of ICD codes is being reported. Enter the indicator between the vertical, dotted lines in the upper right-hand portion of the field.

Enter the codes to identify the patient’s diagnosis and /or condition. List no more than **twelve (12)** diagnosis codes. Relate lines A-L to the lines of service in 24E by the letter of the line.

Item 24A Dates of Service (DOS)

The “From” and “To” DOS will always be the same. Since there is only one DOS, enter the date under “From.” Leave “To” blank or re-enter “From” date.

Item 24B Place of Service (POS)

Enter POS code 99 for all preventive health services and interperiodic visits.

Item 24C EMG (emergency)

Leave blank for “No”.

Item 24D HCPCS Code and Modifier

Enter procedure code and the EP modifier, plus any additional modifiers as applicable.

Item 24E Diagnosis Pointer

Enter the diagnosis code reference letter (pointer) to relate to the DOS and the procedures performed to the primary diagnosis. When multiple services are performed, the primary reference letter for each service should be listed first, other applicable services should follow. Do not use commas between letters.

Item 24H HIPAA EPSDT Referral Codes

Rev. 04/14
Rev. 10/14

- If EPSDT screening resulted in an EPSDT referral, enter the appropriate referral code:
 - ✓ Document AV, S2, or ST in the shaded area of box 24H
- If EPSDT screening did not result in an EPSDT referral:
 - ✓ Document NU in the shaded area of box 24H
- A “Y” for Yes or “N” for No is **not** entered with the referral code in the shaded area or in the unshaded area of box 24H.

Rev. 01/13
 Rev. 07/13
 Rev. 01/14
 Rev. 10/14
 Rev. 10/15

CMS-1500 Health Claim Form – SAMPLE

**SAMPLE EPSDT CLAIM
 9-month preventive visit**

HEALTH INSURANCE CLAIM FORM
 APPROVED BY NATIONAL UNIFORM CLAIMS COMMITTEE (NUCC) 02/12

1. MEDICARE MEDICAID TRICARE CHAMPVA OTHER HEALTH PLAN FICA (LAW) OTHER (DM) (Member ID) (Member ID) (DOB) (DOB) (DOB) (DOB)

2. PATIENT'S NAME (Last Name, First Name, Middle Initial)
Doe, Mary J.

3. PATIENT'S BIRTH DATE SEX
12 25 2014 M F

4. INSURED'S NAME (Last Name, First Name, Middle Initial)
MEDICAID NUMBER

5. PATIENT'S ADDRESS (No. Street)
123 Any Street

6. PATIENT RELATIONSHIP TO INSURED
 Self Spouse Child Other

7. INSURED'S ADDRESS (No. Street)
 CITY STATE ZIP CODE TELEPHONE (Include Area Code)

8. RESERVED FOR NUCC USE

9. OTHER INSURED'S NAME (Last Name, First Name, Middle Initial)
Leave blank

10. IS PATIENT'S CONDITION RELATED TO
 a. EMPLOYMENT? (Current or Previous) YES NO
 b. AUTO ACCIDENT? YES NO PLACE (State)
 c. OTHER ACCIDENT? YES NO

11. INSURED'S POLICY GROUP OR FICA NUMBER

12. INSURED'S DATE OF BIRTH SEX
 MM DD YY M F

13. OTHER CLAIMS (If designated by NUCC)

14. INSURANCE PLAN NAME OR PROGRAM NAME

15. CLAIM CODES (Designated by NUCC)

16. IS THERE ANOTHER HEALTH BENEFIT PLAN?
 YES NO If yes, complete items 5, 6, and 10.

17. INSURER'S OR AUTHORIZED PERSON'S SIGNATURE. I authorize payment of medical benefits to the undersigned physician or supplier for services described herein.
 SIGNATURE DATE SIGNATURE

18. DATES PATIENT UNABLE TO WORK IN CURRENT OCCUPATION
 FROM MM DD YY TO MM DD YY

19. HOSPITALIZATION DATES RELATED TO CURRENT EPISODES
 FROM MM DD YY TO MM DD YY

20. OUTSIDE LAB? \$ CHARGES
 YES NO

21. PRIOR AUTHORIZATION NUMBER

22. PROCEDURE OR NATURE OF ILLNESS OR INJURY (Refer patient to services listed below) (ICD-9-CM) (ICD-10)
Z00.121 or Z00.129 Z23

1	A. DATE(S) OF SERVICE		B. PLACE OF SERVICE	C. ICD-9-CM	D. PROCEDURES, SERVICES, OR SUPPLIES (Explain Unusual Circumstances)	E. DIAGNOSES (ICD-9-CM)	F. CHARGES	G. DRUGS OR INFUSIONS	H. PAYOR	I. BILLING CODES	J. BILLING PROVIDER(S)
	From MM DD YY	To MM DD YY									
1	10 01 15	10 01 15	99 N	99391 EP 25	A				ST		
2	10 01 15	10 01 15	99 N	96110 EP	A						
3	10 01 15	10 01 15	99 N	90460 EP	A			2			
4	10 01 15	10 01 15	99 N	90744 EP	B						
5	10 01 15	10 01 15	99 N	90713 EP	B						
6											

23. FEDERAL TAX ID NUMBER SSN (EIN) 24. PATIENT'S ACCOUNT NO. 25. ACCEPT ASSIGNMENT? (If patient is cash) YES NO 26. TOTAL CHARGE \$ 27. AMOUNT PAID \$ 28. Rev'd for NUCC Use

29. SIGNATURE OF PHYSICIAN OR SUPPLIER INCLUDING DEGREE(S) OR CREDENTIALS (I certify that the statements on the reverse apply to this bill and are made a part thereof.) 30. SERVICE FACILITY LOCATION INFORMATION 31. BILLING PROVIDER INFO & PAYOR ()

SIGNED DATE

NUCC Instruction Manual available at: www.nucc.org PLEASE PRINT OR TYPE APPROVED OMB 0988-1197 FORM 1500 (02-12)

CMS-1500 Health Claim Form – SAMPLE

Rev. 01/13
 Rev. 07/13
 Rev. 01/14
 Rev. 10/14
 Rev. 10/15

SAMPLE EPSDT CLAIM
 Age: 10 months
 9-month Catch-Up preventive visit

HEALTH INSURANCE CLAIM FORM
 APPROVED BY NATIONAL UNIFORM CLAIM COMMITTEE (NUCC) 10/12

1. MEDICAL: MEDICAID <input checked="" type="checkbox"/> TRICARE <input type="checkbox"/> CHAMPVA <input type="checkbox"/> GROUP HEALTH PLAN <input type="checkbox"/> FECA <input type="checkbox"/> OTHER <input type="checkbox"/>		14. INSURED'S ID NUMBER (For Program in Item 1)	
2. PATIENT'S NAME (Last Name, First Name, Middle Initial) Doe, Mary J.		15. INSURED'S NAME (Last Name, First Name, Middle Initial)	
3. PATIENT'S BIRTH DATE: 11 25 2014		16. MEDICAID NUMBER	
5. PATIENT'S ADDRESS (No. Street) 123 Any Street		7. INSURED'S ADDRESS (No. Street)	
6. PATIENT RELATIONSHIP TO INSURED: <input checked="" type="checkbox"/> Self <input type="checkbox"/> Spouse <input type="checkbox"/> Child <input type="checkbox"/> Other		8. INSURED'S ADDRESS (City, State, ZIP Code)	
9. RESERVED FOR NUCC USE		9. RESERVED FOR NUCC USE	
10. IS PATIENT'S CONDITION RELATED TO:		11. INSURED'S POLICY GROUP OR FECA NUMBER	
10.1 EMPLOYMENT? (Current or Previous) <input type="checkbox"/> YES <input type="checkbox"/> NO		12. INSURED'S DATE OF BIRTH: MM DD YY	
10.2 AUTO ACCIDENT? <input type="checkbox"/> YES <input type="checkbox"/> NO		13. OTHER CLAIM # (Designated by NUCC)	
10.3 OTHER ACCIDENT? <input type="checkbox"/> YES <input type="checkbox"/> NO		14. INSURANCE PLAN NAME OR PROGRAM NAME	
10.4 CLAIM CODES (Designated by NUCC)		15. IS THERE ANOTHER HEALTH BENEFIT PLAN? <input type="checkbox"/> YES <input type="checkbox"/> NO	
READ BACK OF FORM BEFORE COMPLETING & SIGNING THIS FORM. 12. PATIENT'S OR AUTHORIZED REPRESENTATIVE'S SIGNATURE & Address or telephone number of the insured or other person necessary to process this claim. Labor receipt payment of government benefits either to itself or to the party who received assignment below.			
16. DATE OF CURRENT ILLNESS, INJURY, OR PREGNANCY (MM DD YY): 10 01 2015		17. OTHER DATE (MM DD YY)	
18. NAME OF REFERRING PROVIDER OR OTHER SOURCE		19. HOSPITALIZATION DATES (RELATED TO CURRENT SERVICES) (MM DD YY)	
20. ADDITIONAL CLAIM INFORMATION (Designated by NUCC)		21. OUTSIDE LAB? <input type="checkbox"/> YES <input type="checkbox"/> NO	
22. DIAGNOSIS OR NATURE OF ILLNESS OR INJURY (Refer to service line below) (ICD-9-CM): Z00.121 or Z00.129, Z23		23. PRIOR AUTHORIZATION NUMBER	
24. A. DATE(S) OF SERVICE: From MM DD YY To MM DD YY		25. B. PLACE OF SERVICE: EMS	
24. C. PROCEDURE(S), SERVICE(S), OR SUPPLY(S) (Specify Unusual Circumstances) (CPT/HCPCS): EP 25 HA		24. D. DIAGNOSIS PORTER	
24. E. \$ CHARGE		24. F. U. \$ CHARGE	
24. G. \$ CHARGE		24. H. \$ CHARGE	
24. I. \$ CHARGE		24. J. \$ CHARGE	
24. K. \$ CHARGE		24. L. \$ CHARGE	
24. M. \$ CHARGE		24. N. \$ CHARGE	
24. O. \$ CHARGE		24. P. \$ CHARGE	
24. Q. \$ CHARGE		24. R. \$ CHARGE	
24. S. \$ CHARGE		24. T. \$ CHARGE	
24. U. \$ CHARGE		24. V. \$ CHARGE	
24. W. \$ CHARGE		24. X. \$ CHARGE	
24. Y. \$ CHARGE		24. Z. \$ CHARGE	
25. FEDERAL TAX ID NUMBER		26. PATIENT'S ACCOUNT NO.	
27. ACCEPT ASSIGNMENT? <input type="checkbox"/> YES <input type="checkbox"/> NO		28. TOTAL CHARGE	
29. SIGNATURE OF PHYSICIAN OR SUPPLIER INCLUDING DEGREE(S) OR CREDENTIALS		29. AMOUNT PAID	
30. SERVICE FACILITY LOCATION INFORMATION		30. Paid by NUCC Use	
31. BILLING PROVIDER INFO & PII #			

NUCC Instruction Manual available at: www.nucc.org PLEASE PRINT OR TYPE APPROVED OMB 0938-1107 FORM 1500 (07-12)

CMS-1500 Health Claim Form – SAMPLE

Rev. 01/13
 Rev. 07/13
 Rev. 01/14
 Rev. 10/14
 Rev. 10/15

SAMPLE EPSDT CLAIM
12-month preventive visit

HEALTH INSURANCE CLAIM FORM

APPROVED BY NATIONAL UNIFORM CLAIM COMMITTEE (NUCC) 02/12

1. MEDICARE <input type="checkbox"/> MEDICAID <input checked="" type="checkbox"/> TRICARE <input type="checkbox"/> CHAMPVA <input type="checkbox"/> GROUP HEALTH PLAN <input type="checkbox"/> FECA (SCLUNG) <input type="checkbox"/> OTHER <input type="checkbox"/>										1a. INSURED'S ID NUMBER (If or Program in Item 1) MEDICAID NUMBER	
2. PATIENT'S NAME (Last Name, First Name, Middle Initial) DOE, John C.				3. PATIENT'S BIRTH DATE 09 30 2014		4. INSURED'S NAME (Last Name, First Name, Middle Initial)		5. PATIENT'S ADDRESS (No., Street) 123 Any Street		6. PATIENT RELATIONSHIP TO INSURED Self <input type="checkbox"/> Spouse <input type="checkbox"/> Child <input type="checkbox"/> Other <input type="checkbox"/>	
7. INSURED'S ADDRESS (No., Street)		8. RESERVED FOR NUCC USE		9. RESERVED FOR NUCC USE		10. IS PATIENT'S CONDITION RELATED TO: a. EMPLOYMENT? (Current or Previous) <input type="checkbox"/> YES <input type="checkbox"/> NO		11. INSURED'S POLICY GROUP OR FECA NUMBER		12. INSURED'S DATE OF BIRTH MM DD YY	
13. OTHER CLAIM ID (Designated by NUCC)		14. INSURANCE PLAN NAME OR PROGRAM NAME		15. IS THERE ANOTHER HEALTH BENEFIT PLAN? <input type="checkbox"/> YES <input type="checkbox"/> NO		16. IS THERE ANOTHER HEALTH BENEFIT PLAN? If yes, complete items 8, 9a, and 9c.		17. INSURER'S OR AUTHORIZED PERSON'S SIGNATURE (Authenticate payment of medical benefits to the undersigned physician or supplier for services described below.)		18. SIGNATURE OF PHYSICIAN OR SUPPLIER	
19. ADDITIONAL CLAIM INFORMATION (Designated by NUCC)		20. OUTSIDE LAB? <input type="checkbox"/> YES <input type="checkbox"/> NO		21. RE SUBMISSION CODE		22. ORIGINAL REF. NO.		23. PRIOR AUTHORIZATION NUMBER		24. A. DATE(S) OF SERVICE From To B. PLACE OF SERVICE C. PROCEDURE(S), SERVICE(S), OR SUPPLIES (Explain Unusual Circumstances) D. MAGNIFIER E. CHARGES F. DAYS OR UNITS G. FREQ. (Per Unit) H. ICD-9-CM I. QUAL. J. RENDERING PROVIDER ID. #	
25. FEDERAL TAX ID NUMBER		26. PATIENT'S ACCOUNT NO.		27. ACCEPT ASSIGNMENT? (For paid, clearly, see back)		28. TOTAL CHARGE \$		29. AMOUNT PAID \$		30. Reserved for NUCC Use	
31. SIGNATURE OF PHYSICIAN OR SUPPLIER INCLUDING DEGREE(S) OR CREDENTIALS (I certify that the statements on the reverse apply to this bill and are made a part thereof.)				32. SERVICE FACILITY LOCATION INFORMATION				33. BILLING PROVIDER INFO & IP# ()			
SIGNED		DATE		a.		b.		c.		d.	

Rev. 01/13
 Rev. 07/13
 Rev. 01/14
 Rev. 07/14
 Rev. 10/14
 Rev. 10/15

CMS-1500 Health Claim Form – SAMPLE

SAMPLE EPSDT CLAIM
Preventive visit with
Immunization Administration Codes
and EPSDT Referral Code

HEALTH INSURANCE CLAIM FORM
 APPROVED BY NATIONAL UNIFORM CLAIM COMMITTEE (NUCC) 03/12

1. MEDICARE MEDICAID TRICARE CHAMPVA GROUP HEALTH PLAN FECA OTHER
 (Medicare) (Medicaid) (TRICARE) (Member ID) (SM) (LAW) (LAW)

2. PATIENT'S NAME (Last Name, First Name, Middle Initial)
DOE, Jane

3. PATIENT'S BIRTH DATE **09 29 2004** SEX F M

4. INSURED'S ID NUMBER (For Program # Item 1)
MEDICAID NUMBER

5. PATIENT'S ADDRESS (No. Street)
123 Any Street

6. PATIENT RELATIONSHIP TO INSURED
 Self Spouse Child Other

7. INSURED'S ADDRESS (No. Street)

8. RESERVED FOR NUCC USE

9. CITY **Anytown** STATE **GA**

10. ZIP CODE **00000-0000** TELEPHONE (Include Area Code) **(404) 123-4567**

11. INSURED'S POLICY GROUP OR FECA NUMBER

12. PATIENT'S CONDITION RELATED TO
 a. EMPLOYMENT (Current or Previous) YES NO
 b. AUTO ACCIDENT? YES NO PLACE (State)
 c. OTHER ACCIDENT? YES NO

13. INSURED'S DATE OF BIRTH **MM DD YY** SEX M F

14. OTHER CLAIM ID (Assigned by NUCC)

15. INSURANCE PLAN NAME OR PROGRAM NAME

16. IS THERE ANOTHER HEALTH BENEFIT PLAN?
 YES NO If yes, complete items 8, 9, and 10.

17. INSURER'S OR AUTHORIZED PERSON'S SIGNATURE. I authorize the release of any medical or other information necessary to process this claim. This request for payment and benefits shall be void if not signed by the party who accepts assignment below.

18. SIGNATURE OF PHYSICIAN OR SUPPLIER

19. DATE

20. DATE OF CURRENT BENEFIT PERIOD OR PREGNANCY (MM/DD/YYYY) QUAL

21. NAME OF REFERRING PHYSICIAN OR OTHER SOURCE

22. HOSPITALIZATION DATES RELATED TO CURRENT SERVICES (MM/DD/YYYY)

23. ADDITIONAL CLAIM INFORMATION (Designated by NUCC)

24. OUTSIDE LAB? YES NO \$ CHARGES

25. DIAGNOSIS OR NATURE OF ILLNESS OR INJURY (Refer back to services (see below) (ICD-9-CM) (ICD-10)

26. PRIOR AUTHORIZATION NUMBER

1	A. DATE(S) OF SERVICE		B. ICD-9-CM	C. PROCEDURE, SERVICE, OR SUPPLY (Explain Unusual Circumstances)	D. DIAGNOSIS (ICD-9-CM)	E. CHARGES	F. DAYS OF INPATIENT	G. H. I. J. K. L. M. N. O. P. Q. R. S. T. U. V. W. X. Y. Z.
	From	To						
1	10 01 15	10 01 15	99	N 99393 EP 25	A			NU
2	10 01 15	10 01 15	99	N 90460 EP	A		2	NPI
3	10 01 15	10 01 15	99	N 90715 EP	B			NPI
4	10 01 15	10 01 15	99	N 90650 EP	B			NPI
5	10 01 15	10 01 15	99	N 90472 EP	A		1	NPI
6	10 01 15	10 01 15	99	N 90649 EP	B			NPI

27. FEDERAL TAX ID NUMBER SSN EIN

28. PATIENT'S ACCOUNT NO.

29. ACCEPT ASSIGNMENT? YES NO

30. TOTAL CHARGE \$

31. AMOUNT PAID \$

32. Paid by NUCC Use

33. SIGNATURE OF PHYSICIAN OR SUPPLIER (INCLUDING DEGREE OR CREDENTIALS) I certify that the statements on this invoice apply to me and are made a part thereof.

34. SERVICE FACILITY LOCATION INFORMATION

35. BILLING PROVIDER INFO ()

SIGNED DATE

NUCC Instruction Manual available at: www.nucc.org PLEASE PRINT OR TYPE APPROVED OMB 0938-1107 FORM 1500 (03-12)

CMS-1500 Health Claim Form – SAMPLE

Rev. 01/14
Rev. 10/14
Rev. 10/15

SAMPLE EPSDT CLAIM
19 year old – Preventive
with Immunization Administration Codes
and EPSDT Referral Code

HEALTH INSURANCE CLAIM FORM

APPROVED BY NATIONAL UNIFORM CLAIM COMMITTEE (NUCC) 02/12

1. MEDICARE <input type="checkbox"/> (Medicare) MEDICAID <input checked="" type="checkbox"/> (Medicaid) TRICARE <input type="checkbox"/> (TRICARE) CHAMPVA <input type="checkbox"/> (ChambVA) GROUP HEALTH PLAN <input type="checkbox"/> (GRP) FECA (BUL LUNG) <input type="checkbox"/> (BUL) OTHER <input type="checkbox"/> (OTH)		14. INSURED'S ID. NUMBER (If on Program in Item 1) MEDICAID NUMBER	
2. PATIENT'S NAME (Last Name, First Name, Middle Initial) DOE, James		3. PATIENT'S BIRTH DATE 09 30 1996 <input checked="" type="checkbox"/> M <input type="checkbox"/> F	
5. PATIENT'S ADDRESS (No., Street) 123 Any Street		6. PATIENT RELATIONSHIP TO INSURED Self <input type="checkbox"/> Spouse <input type="checkbox"/> Child <input type="checkbox"/> Other <input type="checkbox"/>	
CITY Anytown STATE GA		7. INSURED'S ADDRESS (No., Street)	
ZIP CODE 00000-0000 TELEPHONE (Include Area Code) (404) 123-4567		8. RESERVED FOR NUCC USE	
9. OTHER INSURED'S NAME (Last Name, First Name, Middle Initial) Leave blank		10. IS PATIENT'S CONDITION RELATED TO: a. EMPLOYMENT? (Current or Previous) <input type="checkbox"/> YES <input type="checkbox"/> NO b. AUTO ACCIDENT? <input type="checkbox"/> YES <input type="checkbox"/> NO c. OTHER ACCIDENT? <input type="checkbox"/> YES <input type="checkbox"/> NO	
a. OTHER INSURED'S POLICY OR GROUP NUMBER		11. INSURED'S POLICY GROUP OR FECA NUMBER	
b. RESERVED FOR NUCC USE		12. INSURED'S DATE OF BIRTH MM DD YY <input type="checkbox"/> M <input type="checkbox"/> F	
c. RESERVED FOR NUCC USE		13. OTHER CLAIM ID. (Designated by NUCC)	
d. INSURANCE PLAN NAME OR PROGRAM NAME		14. IS THERE ANOTHER HEALTH BENEFIT PLAN? <input type="checkbox"/> YES <input type="checkbox"/> NO <i>If yes, complete items 9, 10, and 11.</i>	
READ BACK OF FORM BEFORE COMPLETING & SIGNING THIS FORM. 12. PATIENT'S OR AUTHORIZED PERSON'S SIGNATURE (I authorize the release of any medical or other information necessary to process this claim. I also request payment of government benefits other to myself or to the party who accepted assignment below.)			
SIGN'D		DATE	
14. DATE OF CURRENT ILLNESS, INJURY, or PREGNANCY (MM/DD/YY) 10 01 2015 QUAL		15. OTHER DATE (MM/DD/YY)	
17. NAME OF AFFIRMING PROVIDER OR OTHER SOURCE		18. HOSPITALIZATION DATES RELATED TO CURRENT SERVICES FROM MM/DD/YY TO MM/DD/YY	
19. ADDITIONAL CLAIM INFORMATION (Designated by NUCC)		20. OUTSIDE LAB? <input type="checkbox"/> YES <input type="checkbox"/> NO \$ CHARGES	
21. DIAGNOSIS OR NATURE OF ILLNESS OR INJURY Refer to service line below (IHL) ICD Ind. Z00.00 or Z00.01 Z23 C. D. E. F. G. H. I. J. K. L.		22. RESUBMISSION CODE ORIGINAL REF. NO.	
24. A. DATE(S) OF SERVICE From MM/DD/YY To MM/DD/YY B. PLACE OF SERVICE CPT/HCPCS MODIFIER E. DIAGNOSIS POINTER F. \$ CHARGES G. DAYS OR UNITS H. FREQ. I. R. QUAL J. BILLING PROVIDER ID. #		23. PRIOR AUTHORIZATION NUMBER	
1 10 01 15 10 01 15 99 N 99395 EP 25 A		AV	
2 10 01 15 10 01 15 99 N 90473 EP A 1		NPI	
3 10 01 15 10 01 15 99 N 90672 EP B		NPI	
4 10 01 15 10 01 15 99 N 90472 EP A 1		NPI	
5 10 01 15 10 01 15 99 N 90649 EP B		NPI	
6		NPI	
25. FEDERAL TAX ID. NUMBER SSN EIN		26. PATIENT'S ACCOUNT NO.	
27. ACCEPT ASSIGNMENT? <input type="checkbox"/> YES <input type="checkbox"/> NO		28. TOTAL CHARGE \$	
29. SIGNATURE OF PHYSICIAN OR SUPPLIER INCLUDING DEGREE(S) OR CREDENTIAL(S) I certify that the statements on the reverse apply to this bill and are made a part hereof.		29. AMOUNT PAID \$	
30. SERVICE FACILITY LOCATION INFORMATION		30. Revid for NUCC Use	
31. BILLING PROVIDER INFO & PPH #			
SIGN'D DATE		SIGN'D DATE	

APPENDIX M

Resources for Children in Georgia

Georgia Public Health Programs

Rev. 01/09
Rev. 07/11
Rev. 07/14
Rev. 07/16
Rev. 07/23

Programs for Children with Disabilities or Special Health Care Needs:

Babies Can't Wait Program (Birth – 3 years)

200 Piedmont Avenue, SE

Atlanta, GA 30334

<http://dph.georgia.gov/Babies-Cant-Wait>

404-657-2850

888-651-8224

Children's Medical Services (Birth – 21 years)

200 Piedmont Avenue, SE

Atlanta, GA 30334

<http://dph.georgia.gov/CMS>

404-657-2850

Children 1st Program

200 Piedmont Avenue, SE

Atlanta, GA 30334

<http://dph.georgia.gov/children1st>

404-657-2850

Women, Infants, and Children (WIC)

200 Piedmont Avenue, SE

Atlanta, GA 30334

<https://dph.georgia.gov/WIC>

1-800-228-9173

Rev. 01/08
Rev. 07/11
Rev. 04/12
Rev. 07/14
Rev. 07/16
Rev. 02/23

Georgia Families

For members in Medicaid or PeachCare for Kids®

Most Medicaid and PeachCare for Kids members must enroll in the Georgia Families managed care program and choose a health plan and a provider.

https://www.georgia-families.com/GASelfService/en_US/home.htm

1-888-GA-ENROLL (1-888-423-6765)

PeachCare for Kids®

CHIP Program

(PeachCare for Kids offers free to low cost health insurance, inclusive of the EPSDT benefit, to uninsured, eligible children living in Georgia)

P.O. Box 786

Alma, GA 31510

www.PeachCare.org

1-877-GA-PEACH (1-877-427-3224)

Vaccines for Children (VFC) Program

GA Department of Public Health Immunization Program

200 Piedmont Avenue, SE

Atlanta, GA 30334

<https://dph.georgia.gov/vaccines-children-program>

(800) 848-3868

(404) 657-5013/ 5015

DPH-Immunization@dph.ga.gov

Georgia Department of Education (GaDOE)

Ask DOE Manager
2054 Twin Towers East
205 Jesse Hill Jr. Drive SE
Atlanta, GA 30334
(404) 656-2800
(800) 311-3627 (GA)
(404) 651-8737 (fax)
askdoe@doe.k12.ga.us

Special Education

Division for Special Education Services and
Supports
Georgia Department of Education
1870 Twin Towers East
Atlanta, GA 30334-9048
(404) 656-3963
Web: www.doe.k12.ga.us/

Programs for Children with Disabilities: Ages 3 through 7

Young Children/619 Coordinator
Division for Special Education Services and Supports
Georgia Department of Education
1870 Twin Towers East
Atlanta, GA 30334-5060
(404) 657-9965
Web: www.doe.k12.ga.us

Division of Family & Children Services (DFCS)

<http://dfcs.dhs.georgia.gov>
1.800.georgia (1.800.436.7442)
678.georgia (678.436.7442) – Atlanta area

DFCS Office of Constituent Services

(404) 657-3433

- [Child Welfare Online Contact Form](#)

complete online contact form for issues related to Adoptions, Child Protective Services, Foster Care or any other Child Welfare issue.

Rev. 07/10
Rev. 07/14
Rev. 07/16

Child Protective Services / Child Abuse & Neglect

1-855-GACHILD / 1-855-422-4453

(404) 657-3400

Medicaid

(877) 423-4746

Food Stamps

(877) 423-4746

Energy Assistance

(877) 423-4746

Temporary Assistance for Needy Families

(877) 423-4746

Department of Behavioral Health and Developmental Disabilities (DBHDD)

200 Piedmont Avenue, SE

Atlanta, GA 30334

404-657-2252

<http://dbhdd.georgia.gov>

Other Resources:

Rev. 07/14

Rev. 10/14

Parent-To-Parent of Georgia

Parent to Parent of Georgia offers a variety of services to Georgia residents ages birth to 26 years and their families impacted by disabilities or special healthcare needs.

3070 Presidential Parkway, Suite 130

Atlanta, GA 30340

(770) 451-5484

(800) 229-2038

Web: <http://p2pga.org>

Healthy Mothers, Healthy Babies Powerline

Source for healthcare referrals and information

2300 Henderson Mill Road

Suite 410

Atlanta, GA 30345

(770) 451-0020

(770) 451-2466

(800) 300-9003

(800) 822-2539

thecoalition@hmbbga.org

www.hmbbga.org

APPENDIX N

General Claims Submission Policy for Ordering, Prescribing, or Referring (OPR) Providers

Rev. 07/13

Rev. 04/14

The Affordable Care Act (ACA) requires physicians and other eligible practitioners who order, prescribe and refer items or services for Medicaid beneficiaries to be enrolled in the GA Medicaid Program. As a result, CMS expanded the claim editing requirements in Section 1833(q) of the Social Security Act and the providers' definitions in sections 1861-r and 1842(b)(18)C. Therefore, claims for services that are ordered, prescribed or referred must indicate who the ordering, prescribing or referring (OPR) practitioner is. The Department will utilize an enrolled OPR provider identification number for this purpose. Any OPR physicians or other eligible practitioners who are NOT already enrolled in Medicaid as participating (i.e. billing) must enroll separately as OPR. The National Provider Identifier (NPI) of the OPR provider must be included on the claim submitted by the participating, i.e. rendering provider. If the NPI of the OPR provider noted on the Georgia Medicaid claim is associated with a provider who is not enrolled in the Georgia Medicaid program, the claim will not be paid.

Effective 4/1/2014, DCH will begin editing claims submitted through the web, EDI, and on CMS 1500 forms for the presence of an ordering, prescribing or referring provider as required by program policy. The edit will be informational until 6/1/2014. Effective 6/1/2014, the ordering, prescribing or referring provider will become a mandatory field and claims that do not contain the information as required by policy will begin to deny.

For the New CMS-1500 claim form:

Enter qualifiers to indicate if the claim has an ordering, prescribing or referring to the left of the dotted line in box 17 (Ordering = DK; Referring = DN or Supervising = DQ).

For claims entered via the web:

Claims headers were updated to accept ordering or referring Provider ID and name for Dental and Institutional claims and the referring provider's name for Professional claims. The claim detail was updated to accept an ordering or referring provider ID and name. Utilize the "ordering" provider field for claims that require a prescribing physician.

For claims transmitted via EDI:

The 837 D, I and P companion guides were updated to specifically point out the provider loops that capture the rendering, ordering, prescribing, referring and service facility provider information that is now used to transmit OPR information.

APPENDIX O

Screening Tools

The CRAFFT Screening Interview

Begin: "I'm going to ask you a few questions that I ask all my patients. Please be honest. I will keep your answers confidential."

Part A

During the PAST 12 MONTHS, did you:	No	Yes
1. Drink any <u>alcohol</u> (more than a few sips)? (Do not count sips of alcohol taken during family or religious events.)	<input type="checkbox"/>	<input type="checkbox"/>
2. Smoke any <u>marijuana</u> or <u>hashish</u> ?	<input type="checkbox"/>	<input type="checkbox"/>
3. Use <u>anything else</u> to <u>get high</u> ? ("anything else" includes illegal drugs, over the counter and prescription drugs, and things that you sniff or "huff")	<input type="checkbox"/>	<input type="checkbox"/>

For clinic use only: Did the patient answer "yes" to any questions in Part A?

No

Yes

↓
Ask CAR question only, then stop

↓
Ask all 6 CRAFFT questions

Part B

	No	Yes
1. Have you ever ridden in a <u>CAR</u> driven by someone (including yourself) who was "high" or had been using alcohol or drugs?	<input type="checkbox"/>	<input type="checkbox"/>
2. Do you ever use alcohol or drugs to <u>RELAX</u> , feel better about yourself, or fit in?	<input type="checkbox"/>	<input type="checkbox"/>
3. Do you ever use alcohol or drugs while you are by yourself, or <u>ALONE</u> ?	<input type="checkbox"/>	<input type="checkbox"/>
4. Do you ever <u>FORGET</u> things you did while using alcohol or drugs?	<input type="checkbox"/>	<input type="checkbox"/>
5. Do your <u>FAMILY</u> or <u>FRIENDS</u> ever tell you that you should cut down on your drinking or drug use?	<input type="checkbox"/>	<input type="checkbox"/>
6. Have you ever gotten into <u>TROUBLE</u> while you were using alcohol or drugs?	<input type="checkbox"/>	<input type="checkbox"/>

CONFIDENTIALITY NOTICE:

The information recorded on this page may be protected by special federal confidentiality rules (42 CFR Part 2), which prohibit disclosure of this information unless authorized by specific written consent. A general authorization for release of medical information is NOT sufficient for this purpose.

© CHILDREN'S HOSPITAL BOSTON, 2009. ALL RIGHTS RESERVED.

Reproduced with permission from the Center for Adolescent Substance Abuse Research, CeASAR, Children's Hospital Boston. (www.ceasar.org)

APPENDIX O

Rev. 10/15

Screening Tools

Patient Health Questionnaire-2 (PHQ-2)

PHQ-2

Over the past two weeks, how often have you been bothered by any of the following problems?

Little interest or pleasure in doing things.
0 = Not at all
1 = Several days
2 = More than half the days
3 = Nearly every day

Feeling down, depressed, or hopeless.
0 = Not at all
1 = Several days
2 = More than half the days
3 = Nearly every day

Total point score: _____

Score interpretation:

<i>PHQ-2 score</i>	<i>Probability of major depressive disorder (%)</i>	<i>Probability of any depressive disorder (%)</i>
1	15.4	36.9
2	21.1	48.3
3	38.4	75.0
4	45.5	81.2
5	56.4	84.6
6	78.6	92.9

Figure 1. Patient Health Questionnaire-2 (PHQ-2). This questionnaire is used as the initial screening test for major depressive episode.

Information from Kroenke K, Spitzer RL, Williams JB. The Patient Health Questionnaire-2: validity of a two-item depression screener. *Med Care* 2003; 41:1284-92.

Thibault JM, Prasaad Steiner, RW. Efficient identification of adults with depression and dementia. *American Family Physician*, Vol. 70/No. 6 (September 15, 2004)

APPENDIX O

Screening Tools

Patient Health Questionnaire-9 (PHQ-9)

PATIENT HEALTH QUESTIONNAIRE-9 (PHQ-9)

Over the last 2 weeks, how often have you been bothered by any of the following problems?
(Use "✓" to indicate your answer)

	Not at all	Several days	More than half the days	Nearly every day
1. Little interest or pleasure in doing things	0	1	2	3
2. Feeling down, depressed, or hopeless	0	1	2	3
3. Trouble falling or staying asleep, or sleeping too much	0	1	2	3
4. Feeling tired or having little energy	0	1	2	3
5. Poor appetite or overeating	0	1	2	3
6. Feeling bad about yourself — or that you are a failure or have let yourself or your family down	0	1	2	3
7. Trouble concentrating on things, such as reading the newspaper or watching television	0	1	2	3
8. Moving or speaking so slowly that other people could have noticed? Or the opposite — being so fidgety or restless that you have been moving around a lot more than usual	0	1	2	3
9. Thoughts that you would be better off dead or of hurting yourself in some way	0	1	2	3

For office codes: 0 + + +
=Total Score:

If you checked off any problems, how difficult have these problems made it for you to do your work, take care of things at home, or get along with other people?

Not difficult at all	Somewhat difficult	Very difficult	Extremely difficult
<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>

Developed by Drs. Robert L. Spitzer, Janet B.W. Williams, Kurt Kroenke and colleagues, with an educational grant from Pfizer Inc. No permission required to reproduce, translate, display or distribute.

APPENDIX P

2016 Policy Revisions Record

Rev. 01/17

Part II Policies and Procedures Manual for EPSDT (Health Check) Services

REVISION DATE	SECTION & PAGE	REVISION DESCRIPTION	REVISION TYPE A = Added D = Deleted M = Modified	CITATION (Revision required by Regulation, Legislation, etc.)
01/01/16	902	Revised AAP BF Periodicity Schedule	M	AAP Bright Futures (BF)
01/01/16	1003.3	Depression Screenings and Alcohol/SA Risk Assessments (96127) – added	A	N/A
01/01/16	1003.16	Incontinence Products - added	A	N/A
01/01/16	Appendix C-2	Vaccine product codes: 90634 - Hepatitis A – removed 90655 – Influenza - added	D A	N/A
01/01/16	Appendix R	<ul style="list-style-type: none"> Deleted BF Periodicity Schedule Added upcoming 4/16 EPSDT Updates 	D A	N/A
04/01/16	Introduction iii	Added brief emotional/behavioral assessments to services reimbursed under Health Check	M	N/A
04/01/16	902 p. IX-1-3	Revised AAP BF Periodicity Schedule (copyright 2016, updated 10/15)	A	AAP BF
04/01/16	902.1 #7, #25, #26 p. IX-4-7	Updated BF Footnotes: <ul style="list-style-type: none"> Visual Acuity, Oral Health Fluoride Varnish 	M A	AAP BF
04/01/16	902.2 D.1., p. IX-9 E.4 & 5, p. IX-12-13 G.1.b., p. IX-14-15 G.8., p. IX-18 I., p. IX-19	Updated Screening Components <ul style="list-style-type: none"> Vision- routine screening at age 18 changed to risk assessment. Clarifications to Alcohol/Substance Abuse Risk Assessment & Depression Screening Critical Congenital Heart Disease (CCHD) Screening – updated Cervical Dysplasia / Pap Test – updated Fluoride Varnish –added 	M M M M A	AAP BF N/A AAP BF AAP BF N/A
04/01/16	904 Table A p. IX-23 & 23	99384 & 99394 – ICD-10 Z00.00 & Z00.01 for ages 15-17 years	M	ICD-10 coding guidelines
04/01/16	Immunization Schedules IX-30 - 32	2016 Immunizations Schedules posted	M	CDC
04/01/16	1003.3 p. X-2 & 3	Brief Emotional / Behavioral Assessment (96127) – updated	M	N/A

APPENDIX P

2016 Policy Revisions Record

REVISION DATE	SECTION & PAGE	REVISION DESCRIPTION	REVISION TYPE A = Added D = Deleted M = Modified	CITATION (Revision required by Regulation, Legislation, etc.)
04/01/16	1003. p. 5-7 #13 #17 #19 #20	<ul style="list-style-type: none"> • School-based Telemedicine Services for LEAs (Q3014) – updated • Fluoride Varnish – added • NCCI-MUE limits – added • Other Reimbursement Rates - added 	M A A A	N/A
04/01/16	Appendix C-2	Minor updates to vaccine procedure code descriptions: 90620, 90621, 90632, 90633, 90644, 90647, 90648, 90649, 90650, 90651, 90655, 90656, 90657, 90658, 90670, 90672, 90680, 90681, 90685, 90686, 90687, 90688, 90696, 90698, 90714, 90716, 90732, 90734, 90744, 90746, 90474 90670 – clarified age restriction (6 weeks to 17 years) 90733 - added for ages 19-20 years	M M A	2016 HCPCS N/A N/A
04/01/16	Appendix H	updates to CMO PBM, BIN# and PCN for AMG & WC	M	N/A
04/01/16	Appendix J	Fluoride varnish - revised	M	N/A
04/01/16	Appendix R	Removed upcoming 4/2016 EPSDT Updates	D	N/A
07/01/16	902.2 p. IX-7	Bright Futures helpful materials – updated hyperlink	M	N/A
07/01/16	902.2, D.2. p. IX-10	EHDI Program - updated hyperlink and info	M	N/A
07/01/16	902.2, E.3. 902.2, E.4. 902.2, E.5. p. IX-12 & 13	MCHAT tool - updated hyperlink CRAFFT tool - added hyperlink PHQ-2 tool - updated hyperlink	M A M	N/A
07/01/16	902.2, G.1.a. & b. 902.2, G.2 p. IX-14 & 15	Georgia NBS Program - added hyperlink and minor info GRITS - added hyperlink	A	N/A
07/01/16	Tables A, C p. IX-22, 27	Minor revision for clarification related to modifiers	M	N/A
07/01/16	Appendix A p. A-1- 4	Minor revisions Posted the HB76 FY2016 PCP Increased Rates <i>(Implementation of rates is pending. Refer to Banner Message dated 6-17-2016)</i>	M A	N/A Legislation - HB76

APPENDIX P

2016 Policy Revisions Record

Part II Policies and Procedures Manual for EPSDT (Health Check) Services

REVISION DATE	SECTION & PAGE	REVISION DESCRIPTION	REVISION TYPE A = Added D = Deleted M = Modified	CITATION (Revision required by Regulation, Legislation, etc.)
07/01/16	Appendix F p. F-2 - 4	Counseling for Nutrition and Physical Activity – added info related to HEDIS requirements	A	HEDIS
07/01/16	Appendix M	Updated Resources	M	N/A
10/01/16	902.2 H., p. IX-19 902.2 I., p. IX-19 1003. #17. p. X-6 Appendix J p. J-1	Added clarifications to Oral Health section. Revised the statement to read: “The AAP recommends the establishment of a dental home six months after the first tooth erupts or by 12 months of age (whichever comes first). For clarification, deleted the statement “ <i>One application of fluoride varnish is required for children between the ages of 6 months and 5 years.</i> ” Revised to read, “Once teeth are present, the application of fluoride varnish is required and may be applied every 3-6 months in the primary care or dental office for children between the ages of 6 months and 5 years.”	M	AAP Bright Futures (BF)
10/01/16	Table A-1 p. IX-24 Table B-2 p. IX-26 Table C-2 p. IX-28 1003. #20. p. X-7 Appendix C-1 Table C-1b p. C-1	Posted the HB 751 FY 2017 PCP rate increases for the preventive visits 99381, 99391-99395 Posted the HB 751 FY 2017 PCP rate increases for the catch-up preventive visits 99381, 99391, 99392 Posted the HB 751 FY 2017 PCP rate increases for the interperiodic visits 99202-99203, 99212-99214 Posted the HB 751 FY 2017 PCP rate increase for the office consultation code 99244 Posted the HB 751 FY 2017 PCP rate increase for the vaccine administration codes 90460, 90471, 90472	A	Legislation – HB 751

APPENDIX P

2016 Policy Revisions Record

Part II Policies and Procedures Manual for EPSDT (Health Check) Services

REVISION DATE	SECTION & PAGE	REVISION DESCRIPTION	REVISION TYPE A = Added D = Deleted M = Modified	CITATION (Revision required by Regulation, Legislation, etc.)
10/01/16	Appendix C-2 p. C-7	90630 (<i>influenza virus vaccine, quadrivalent, intradermal</i>) - added for ages 19-20 years	A	ACIP
10/01/16	Appendix F p. F-4	Under ICD-10 Codes to Identify Counseling for Physical Activity, ICD-10 diagnosis code Z71.89 (other specified counseling) was deleted.	M	HEDIS
10/01/16	Appendix H p. H-2	Updated the Georgia Families Regions – “Counties” Column revised	M	Policy
10/01/16	Appendix H p. H-11	Updated with the new PBM information for Peach State Health Plan	M	Policy

APPENDIX P

2017 Policy Revisions Record

Part II Policies and Procedures Manual for EPSDT (Health Check) Services

Rev. 01/18

REVISION DATE	SECTION & PAGE	REVISION DESCRIPTION	REVISIO TYPE A = Added D = Deleted M = Modified	CITATION (Revision required by Regulation, Legislation, etc.)
01/01/17	902.2 H. p. IX-19	Clarifications to Oral Health: <ul style="list-style-type: none"> The AAP recommends both the establishment of a dental home and the first dental exam no later than 12 months of age. An oral health risk assessment tool has been developed by the AAP/Bright Futures. 	M	AAP Bright Futures (BF)
01/01/17	Appendix C-2	The 2017 HCPCS influenza vaccine code description changes will not go into effect until the updates are completed in GAMMIS.	M	HCPCS
01/01/17		The “Mapping of EPSDT Preventive Health ICD-9 Codes to ICD-10 Codes” table was removed from Appendices. Accordingly, all references to the table were removed throughout the manual.	D	N/A
01/01/17	Appendix Q	The “2016 Policy Revisions Record” was added as Appendix Q.	A	N/A
04/01/17	902 p. IX-1	Notification – updated AAP 2017 BF Periodicity Schedule (effective date July 1, 2017)	M	AAP Bright Futures (BF)
04/01/17	p. IX-30-32	2017 Immunization Schedules posted	M	ACIP
04/01/17	Appendix C-2	90670 – age restriction removed. Refer to the ACIP Immunization Schedule for recommended age	M	N/A
04/01/17	Appendix H	Georgia Families – updated appendix	M	N/A
04/01/17	Appendix O	Performance Measures - minor updates	M	N/A
04/01/17	Appendix R	2017 BF Periodicity Schedule – displays updated schedule and summary of changes	A	AAP Bright Futures (BF)

APPENDIX P

2017 Policy Revisions Record

Part II Policies and Procedures Manual for EPSDT (Health Check) Services

REVISION DATE	SECTION & PAGE	REVISION DESCRIPTION	REVISIO TYPE A = Added D = Deleted M = Modified	CITATION (Revision required by Regulation, Legislation, etc.)
07/01/17	Entire Manual	Vendor name change - HPE updated to reflect new name –DXC Technology (DXC)	M	N/A
07/01/17	<p>902-902.1 p. IX-1 – IX-7</p> <p>902.2 p. IX-11 – IX-22</p> <p>903 IX-23</p>	<p>Updated 2017 BF Periodicity Schedule – adoption of new schedule effective July 1, 2017 with revised footnotes. Added new screening BF requirements, and deleted/ revised current screening requirements.</p> <p>Revised the minimum standards for screening components in accordance with the updated 2017 BF Periodicity Schedule. Changes to:</p> <ul style="list-style-type: none"> • Hearing • Tobacco, Alcohol, or Drug Use Assessment • Depression Screening • Maternal Depression Screening • Newborn Blood • Newborn Bilirubin • Dyslipidemia Screening • STIs • HIV • Oral Health <p>Screening tools requirements revised to reflect updated 2017 BF Periodicity Schedule.</p>	M	AAP Bright Futures (BF)
07/01/17	902.2, G.4. p. IX-18, 907.2, C. p. IX-40, Appendix A, p. A-5	Blood Lead Test – provided clarification that “All venous sample lead screening tests conducted using any Magellan Diagnostic lead testing system should be laboratory analyzed by a properly accredited laboratory.”	M	FDA, CDC, GA. Dept. of Public Health (DPH)
07/01/17	1003, #3, p. X-2	Effective July 1, 2017, DCH will allow separate reimbursement for the Autism screening (96110 EP, UA)	A	DCH

APPENDIX P

2017 Policy Revisions Record

Part II Policies and Procedures Manual for EPSDT (Health Check) Services

REVISION DATE	SECTION & PAGE	REVISION DESCRIPTION	REVISIO TYPE A = Added D = Deleted M = Modified	CITATION (Revision required by Regulation, Legislation, etc.)
07/01/17	1003, p. X-2 – X-3, p. X-7	Billing Requirements revisions: <ul style="list-style-type: none"> • added Autism screening code, modifiers and rate, • revised age ranges for brief emotional/behavioral assessments 	M	DCH/ AAP Bright Futures (BF)
10/01/17	1003, #2 & #3, p. X-2	2) Developmental Screenings – included ICD-10 diagnosis codes 3) Autism Screenings – included ICD-10 diagnosis codes	M	DCH
10/01/17	Appendix C-2	Updates to influenza vaccines <ol style="list-style-type: none"> 1) 90672 - removed 2) 90674 – added 3) 90682 – added (19-20 years) 	A, D	DCH

APPENDIX P

2018 Policy Revisions Record

Part II Policies and Procedures Manual for EPSDT (Health Check) Services

Rev. 01/19

REVISION DATE	SECTION & PAGE	REVISION DESCRIPTION	REVISIO TYPE A = Added D = Deleted M = Modified	CITATION (Revision required by Regulation, Legislation, etc.)
10/1/2018	902.2.E.4. p. IX-14	➤ Tobacco, Alcohol, or Drug Use Assessment: Added clarification to Documentation - Screening tool must be standardized and scorable.	A	DCH
10/1/2018	902.2.E.6. p. IX-15	➤ Maternal Depression Screening: Added clarification to Documentation - Screening tool must be standardized and scorable.	A	DCH
10/1/2018	1003.7 p. X-3	➤ Autism Screenings: Updated ICD-10 diagnosis code Z13.4 to Z13.41	M	DCH
10/1/2018	1003.9. p. X-4	➤ Patient-Focused Health Risk Assessment - (96160) updated the guidance in order to allow the 59 modifier to bypass the NCCI PTP edit.	M	DCH
10/1/2018	1003.10. p. X-4-5	➤ Caregiver-Focused Health Risk Assessment - (96161) updated the guidance in order to allow the 59 modifier to bypass the NCCI PTP edit.	M	DCH
10/1/2018	1003.23. p. X-9	➤ NCCI PTP Edits –added guidance on NCCI PTP edits	A	DCH
10/1/2018	1003.23. p. X-9	➤ Q3014 – revised to include EP GT modifiers	M	DCH
10/1/2018	Appendix C-2 p. C-5	Updated vaccine codes (ages birth through 18 years) in accordance with current VFC supply ➤ Removed 90644, 90649, 90650 ➤ Removed trivalent influenza vaccines 90655, 90656, 90657, 90658	D	DCH
10/1/2018	Appendix C-2 p. C-7	Updated vaccine codes (ages 19 years through 20 years) in accordance with current VFC supply ➤ Removed HP bivalent vaccine 90650 ➤ Removed trivalent influenza vaccines 90658 ➤ Removed 90733	D	DCH
10/1/2018	Appendix H	Updated Georgia Families Appendix	M	DCH

APPENDIX P

2018 Policy Revisions Record

Part II Policies and Procedures Manual for EPSDT (Health Check) Services

REVISION DATE	SECTION & PAGE	REVISION DESCRIPTION	REVISIO TYPE A = Added D = Deleted M = Modified	CITATION (Revision required by Regulation, Legislation, etc.)
07/01/18	1003, p. X-3-4	<ul style="list-style-type: none"> ➤ Revised billing guidance for brief emotional/behavioral assessments (96127) ➤ New billing guidance for patient focused health risk assessments (96160) ➤ New billing guidance for maternal depression screenings (96161) 	M A A	DCH
07/01/18	Appendix I p. I-1	➤ Georgia Families 360SM Appendix-Revisions to info for providers serving Medicaid members in the Georgia Families 360SM Program	M	DCH
04/01/18	905, p. IX-37-41	2018 Immunization Schedules posted	A	ACIP
04/01/18	<p>Appendix C-1, Table C-2, p. C-5</p> <p>904, Table C-1, p. IX-32 904, Table C-2, p. IX-33</p> <p>1003, #21, p. X-8</p> <p>904, Table A-1 p. IX-27 904, Table B-2, p. IX-30</p>	<p>Posted the HB 44 FY 2018 PCP rate increase for the following codes:</p> <p>vaccine administration codes</p> <ul style="list-style-type: none"> ➤ 90473 – FFS ➤ 90474 – FFS, PCK <p>interperiodic visit codes</p> <ul style="list-style-type: none"> ➤ 99201 – FFS ➤ 99211 – FFS, PCK <p>office consultation codes</p> <ul style="list-style-type: none"> ➤ 99241, 99242, 99243, 99245 <p>behavior change smoking codes</p> <ul style="list-style-type: none"> ➤ 99406, 99407 <p>preventive visit codes 99382, 99383, 99384, 99385</p>	A	Legislation – HB 44

APPENDIX P

2018 Policy Revisions Record

Part II Policies and Procedures Manual for EPSDT (Health Check) Services

REVISION DATE	SECTION & PAGE	REVISION DESCRIPTION	REVISIO TYPE A = Added D = Deleted M = Modified	CITATION (Revision required by Regulation, Legislation, etc.)
04/01/18	Appendix C-2 p. C-7 p. C-9	Added new 2018 HCPCS code 90756 – Influenza virus vaccine, quadrivalent (ccIV4), derived from cell cultures, subunit, antibiotic free, 0.5mL dosage, for intramuscular use	A	HCPCS
04/01/18	Appendix F	Clarifications to Weight Assessment - BMI	M	N/A
01/01/18	#6 p. IX-14 & 15	Maternal Depression – added clarification to recommended tools	M	AAP USPSTF
01/01/18	Table C p. IX-32	Updated the hyperlink for the DPH Form 3300 - Certificate of Vision, Hearing, Dental and Nutrition Screening	M	DPH
01/01/18	Appendix C p. C-5 p. C-7	Revised descriptions for CPT codes 90620 90621 90651	M	HCPCS 2018
01/01/18	Appendix F p. F-2 p. F-4	1) Documentation for BMI – added the following clarification: “Only evidence of the BMI percentile or BMI percentile plotted on an age-growth chart meet HEDIS criteria.” 2) Under the section containing the notations or examples of documentation that are not compliant with HEDIS requirements for Nutrition , added the following: ○ Documentation related to a member’s “appetite” does not meet criteria.	M	HEDIS 2018

APPENDIX P

2019 Policy Revisions Record

Part II Policies and Procedures Manual for EPSDT (Health Check) Services

Rev. 01/20

REVISION DATE	SECTION & PAGE	REVISION DESCRIPTION	REVISION TYPE A = Added D = Deleted M = Modified	CITATION (Revision required by Regulation, Legislation, etc.)
10/2019	902.2, E.4., p. IX-14	4. Tobacco, Alcohol, or Drug Use Assessment – updated hyperlink to access CRAFFT screening tool	M	DCH
10/2019	902.2, G.5., p. IX-20	5. Tuberculin Risk Assessment and Test – updated hyperlinks to access info o DPH TB and forms	M	DCH/DPH
10/2019	908, p. IX-44	Oral Health and Dental Services - Added guidance - Periodicity of Examination, Preventive Dental Services, Anticipatory Guidance/Counseling, and Oral Treatment for Infants, Children, and Adolescents	A	DCH
10/2019	908, p. IX-45	Oral Health and Dental Services - Replaced the Recommendations for Pediatric Oral Health Assessment, Preventive Services, and Anticipatory Guidance/Counseling	M	DCH
10/2019	Appendix A, p. A-2	“Lead Screening Requirements and Medical Management Recommendations for Children” – Updated table to reflect blood lead level of 5 µg/dL or greater requires further testing and monitoring.	M	DPH
10/2019	Appendix A, p. A-3	D) Lead Poisoning Education – minor rewording revision	M	DPH
10/2019	Appendix A, p. A-4	Blood Lead Test – minor rewording revision	M	DPH
10/2019	Appendix A, p. A-5	Lab Submission – lab submission guidance section added	A	DPH

APPENDIX P

2019 Policy Revisions Record

Part II Policies and Procedures Manual for EPSDT (Health Check) Services

REVISION DATE	SECTION & PAGE	REVISION DESCRIPTION	REVISIO TYPE A = Added D = Deleted M = Modified	CITATION (Revision required by Regulation, Legislation, etc.)
10/2019	Appendix A, p. A-5, A-6	Reporting Guidelines - added clarification to the guidance - provided hyperlink for current Blood Lead Test Reporting Log provided by GHHLPPP - deleted previous copy of Blood Lead Test Reporting Log SendSS ‘Registration and Login Manual for Uploading Lead Report Files’ - - Updated hyperlink to access SendSS	M	DPH
10/2019	Appendix A, p. A-15 - A-18 Appendix A, p. A-19	Georgia Childhood Lead Poisoning Prevention Program Case Management Guidelines - updated guidelines to reflect blood lead level of 5 µg/dL or greater requires further testing and monitoring. - updated Regional Lead Coordinator (RLC) title to Regional Healthy Homes Coordinator (RHHC) - removed the specific number of coordinators. Georgia Public Health Laboratory (GPHL) - updated locations, removed Albany location	M	DPH
10/2019	Appendix C-4 p. C-10, C-11	Blood Lead Level Testing Procedure Codes - Clarifications to billing guidance	M	DCH
10/2019	Appendix J, p. J-1, J-2	Preventive Oral Health: Fluoride Varnish - - revised hyperlinks -	M	ADA

APPENDIX P

2019 Policy Revisions Record

Part II Policies and Procedures Manual for EPSDT (Health Check) Services

REVISION DATE	SECTION & PAGE	REVISION DESCRIPTION	REVISION TYPE A = Added D = Deleted M = Modified	CITATION (Revision required by Regulation, Legislation, etc.)
07/2019	902, p. IX-1 – IX-7	AAP Bright Futures Periodicity Schedule, updated – Effective July 1, 2019, DCH will adopt the updated 2019 AAP Bright Futures Periodicity Schedule. Copyright © 2019 by the American Academy of Pediatrics, updated March 2019	M	AAP/ BF
07/2019	Appendix C-1 p. C-3	<p>ICD 10 diagnosis code Z23.0 was added as a reportable diagnosis with the applicable vaccine administration code.</p> <ul style="list-style-type: none"> • May report diagnosis code Z00.121 or Z00.129 or Z23.0 with each of the vaccine administration codes ONLY when vaccines are administered during EPSDT preventive health visits for members through age 17 years. • May report diagnosis code Z00.00 or Z00.01 or Z23.0 with the applicable vaccine administration code ONLY when vaccines are administered during EPSDT preventive health visits for members age 15 years through 20 years 	A	DCH
07/2019	Appendix C-2 p. C-6 p. C-8	<p>90734 – revised long description.</p> <p><u>Revised description:</u> Meningococcal conjugate vaccine, serogroups A, C, W, Y, quadrivalent, diphtheria toxoid carrier, (MenACWY-D) or CRM197 carrier (MenACWY-CRM), for intramuscular use</p>	M	<p>American Medical Association (AMA) – Jan. 1, 2019 released to AMA website Eff. July 1, 2019 Updated 12/19/18</p> <p>CPT – Publication CPT® 2020</p>

APPENDIX P

2019 Policy Revisions Record

Part II Policies and Procedures Manual for EPSDT (Health Check) Services

REVISION DATE	SECTION & PAGE	REVISION DESCRIPTION	REVISIO TYPE A = Added D = Deleted M = Modified	CITATION (Revision required by Regulation, Legislation, etc.)
04/2019	902, p. IX-1	Notification – effective July 1, 2019 DCH will adopt the updated 2019 AAP Bright Futures Periodicity Schedule	M	AAP/ BF
04/2019	905, p. IX-37-41	2019 CDC Immunization Schedules posted	A	CDC/ACIP
04/2019	1003, p. X-3-6 #6-10	Clarifications – (1) the appropriate ICD-10 diagnosis code may be either preventive or non-preventive; (2) the examples of the ICD-10 diagnosis codes listed are preventive	M	DCH
04/2019	Appendix H p. H-15	Pharmacy – updated Peach State Health Plan PBM US Script to PBM Envolve Pharmacy Solutions	M	DCH
04/2019	Appendix Q	2019 AAP Bright Futures Periodicity Schedule - summary of changes	A	AAP/ BF
04/2019	Appendix T	2019 CDC Immunization Schedules – summary of changes	A	CDC/ACIP
01/2019	Appendix C-2	90713 – added Poliovirus (IPV) added for high risk members, ages 19-20 yrs	A	DCH
01/2019	Appendix C-2	90630 – removed Influenza virus vaccine, quadrivalent (IIV4), no preservative, for intradermal use	D	CDC/DCH
01/2019	Appendix C-2	90672 - added Influenza virus vaccine (FluMist), quadrivalent, live (LAIV4), for intranasal use (≥2 years)	A	CDC
01/2019	Appendix	Performance Measures Appendix– removed	D	DCH

APPENDIX P

2020 Policy Revisions Record

Part II Policies and Procedures Manual for EPSDT (Health Check) Services

Rev. 01/21

REVISION DATE	SECTION & PAGE	REVISION DESCRIPTION	REVISIO TYPE A = Added D = Deleted M = Modified	CITATION (Revision required by Regulation, Legislation, etc.)
10/2020	Appendix C-2, p. C-7	CPT 90656 influenza virus vaccine, trivalent (IIV3) – deleted from vaccines for ages 19-20 years	D	DCH/ CDC/ CMS
07/2020	Appendix S	Clarifications that rates are for Medicaid-eligible and PeachCare for Kids-eligible members.	M	DCH
04/2020	905.4, p. IX-36	New 11 th Grade Immunization Requirement – booster dose of meningococcal conjugate vaccine required for new entrants or transfers into a GA school in the eleventh grade for 2020-21 academic school year	A	DPH
04/2020	905, p. IX-37-41	2020 CDC Immunization Schedules posted	A	CDC/ACIP
04/2020	Appendix A	Risk Factors Assessment Questionnaire revised	M	DPH
04/2020	Appendix R	2020 Immunization Schedules Changes & Guidance	A	CDC/ACIP
04/2020	Appendix S	EPSDT Health Check Reimbursement Rates	A	DCH
01/2020	1003, p. X-7 & 8 # 19	Incontinence Supplies: <i>Minimim age revised to 2 years.</i> Incontinence supplies are covered for children ages 2 through 21 years of age who have an underlying medical condition that prevents control of the bowels or bladder.	M	CMS
01/2020	Appendix A, p. A-1	Clarification made to requirements for Blood Lead Risk Assessment – Current Language: the questionnaire should be administered at 6, 9, and 18 months, and 3, 4, 5, 6 years of age.	M	DCH
01/2020	Appendix C-1 p. C-3	Correction: ICD 10 diagnosis code Z23 was inadvertently listed as Z23.0	M	DCH

APPENDIX P

2021 Policy Revisions Record

Part II Policies and Procedures Manual for EPSDT (Health Check) Services

REVISION DATE	SECTION & PAGE	REVISION DESCRIPTION	REVISIO TYPE A = Added D = Deleted M = Modified	CITATION (Revision required by Regulation, Legislation, etc.)
10/2021	904, Table A p. IX-26, IX-30 1003, #12, p. X-6 Appx C-4, p.C-10	83655 – clarifications to ICD-10 diagnosis code (i.e., Z13.88, Z00.121, Z00.129, Z77.011) to signify blood lead level screening.	M	DCH
10/2021	1003, #15, p. X-7	Meningococcal conjugate - DPH providers should use POS code 03 when billing vaccine administration fee for meningococcal conjugate vaccine (90734) administered during school-based vaccine clinics.	M	DCH
10/2021	Appendix C-2, p. C-6, p. C-5, C-7	New vaccines: -Vaxelis vaccine (90697) eff 7-1-2021 -MenQuadfi (90619) eff 4-1-2021	A	DCH/ DPH
07/2021	902, p. IX-1, 4-5	Effective July 1, 2021 – adopted the updated AAP 2021 Bright Futures Periodicity Schedule. Following footnotes revised: • #11: Developmental – updated • #12: Autism Spectrum Disorder - updated • #31: Hepatitis C Virus Infection added – Screening for Hep C virus infection has been added to occur at least once between the ages of 18 and 79 years (to be consistent with recommendations of the USPSTF and CDC)	M	AAP
07/2021	902.2, E.2., p. IX-13-14	Developmental Screening – reporting criteria updated in accordance with the Core Set of Children’s Health Care Quality Measures for Medicaid and CHIP (Child Core Set) Technical Specs & Resource Manual for FFY2021 Reporting March 2021	M	CMS
07/2021	Appendix A, p. A1-A3	Guidelines in Lead Screening • updates to Recommended Medical and Case Mgmt Actions- Blood Lead Levels	M	DPH (GHHLPPP)
07/2021	Appendix H	Georgia Families - Updated to remove occurrences of WellCare from the appendix	M	DCH

APPENDIX P

2021 Policy Revisions Record

Part II Policies and Procedures Manual for EPSDT (Health Check) Services

REVISION DATE	SECTION & PAGE	REVISION DESCRIPTION	REVISIO TYPE A = Added D = Deleted M = Modified	CITATION (Revision required by Regulation, Legislation, etc.)
04/2021	902, IX-1, 4-5	<p>Effective July 1, 2021, the Georgia Division of Medical Assistance Plans will adopt the AAP 2021 Bright Futures “Recommendations for Pediatric Health Care” Periodicity Schedule as the periodicity schedule for EPSDT visits and services. The schedule is available at https://www.aap.org/en-us/Documents/periodicity_schedule.pdf The updated 2021 schedule consists of revisions to footnote 11 (developmental) and footnote 12 (autism spectrum disorder). Footnote 31 (hepatitis C virus infection) was added. A copy of the summary of changes is displayed in Appendix Q.</p> <p>AAP 2021 Bright Futures “Recommendations for Pediatric Health Care” Periodicity Schedule- updated</p> <ul style="list-style-type: none"> • Footnote 11: Developmental – updated • Footnote 12: Autism Spectrum Disorder - updated • Footnote 31: Hepatitis C Virus Infection added – Screening for hepatitis C virus infection has been added to occur at least once between the ages of 18 and 79 years (to be consistent with recommendations of the USPSTF and CDC) 	M	AAP
04/2021	905, p IX 39-43	2021 CDC Immunization Schedules posted	M	CDC/ACIP
04/2021	Appendix E	Non-Emergency Medical Transportation (NEMT) – updated	M	DCH
04/2021	Appendix Q	AAP Bright Futures Periodicity Schedule summary of changes: 2021 changes added	M	AAP
04/2021	Appendix R	2021 CDC Immunization Schedule – summary of changes added	M	CDC/ACIP

APPENDIX P

2021 Policy Revisions Record

Part II Policies and Procedures Manual for EPSDT (Health Check) Services

REVISION DATE	SECTION & PAGE	REVISION DESCRIPTION	REVISIO TYPE A = Added D = Deleted M = Modified	CITATION (Revision required by Regulation, Legislation, etc.)
01/2021	Entire Manual	Vendor name change - DXC Technology (DXC) updated to reflect new name – Gainwell Technologies	M	DCH
01/2021	Tables C, C-1, C-2 p. IX-32 & 33 Appendix C-1, p. C-3 Appendix S, p. S-2	CPT code 99201 deleted effective 1/1/2021	D	CMS

APPENDIX P

2022 Policy Revisions Record

Part II Policies and Procedures Manual for EPSDT (Health Check) Services

REVISION DATE	SECTION & PAGE	REVISION DESCRIPTION	REVISIO TYPE A = Added D = Deleted M = Modified	CITATION (Revision required by Regulation, Legislation, etc.)
10/2022	902, p. IX-1	Notification – Effective January 1, 2023, DCH will adopt the AAP 2022 BF Periodicity Schedule. The schedule is available at https://www.aap.org/en-us/Documents/periodicity_schedule.pdf	M	AAP
10/2022	Appendix C-5, p. C-11	For listing of Covid-19 Vaccines & Administration Codes, see the ‘ <i>Provider’s Administered Drug List – Covid-19 Vaccine Administration Codes</i> ’ fee schedule located on GAMMIS.	A	DCH
10/2022	Appendix Q, p. Q-1 – Q-4	AAP 2022 BF Periodicity Schedule posted	A	AAP
07/2022		July 2022 - No policy updates		
04/2022	905, IX-38 – IX-42	2022 CDC Immunization Schedules posted	M	CDC/ACIP*
04/2022	1003, #24 p. X-10	99401 Preventive Medicine Counseling added	A	DCH
01/2022	Page iv	Added title ‘Autism Spectrum Disorder (ASD) Services Manual’ to listing of provider policy manuals relevant to the EPSDT benefit.	A	DCH
01/2022	902.2, E., #3 p. IX-15 1003, #7 p. X-3	Added the following language: “For guidelines related to Autism Spectrum Disorder, including a list of acceptable Autism Spectrum Disorder Assessment Tools, refer to the Autism Spectrum Disorder policy manual Section 801.”	A	DCH

APPENDIX P

2023 Policy Revisions Record

Part II Policies and Procedures Manual for EPSDT (Health Check) Services

REVISION DATE	SECTION & PAGE	REVISION DESCRIPTION	REVISION TYPE A = Added D = Deleted M = Modified	CITATION (Revision required by Regulation, Legislation, etc.)
10/2023	Appendix C-2	Added the following vaccine: - 90759 HepB vaccine (Hevbrio) ages 19-20 years eff. 08/01/2023	A	DCH/ DPH
07/2023	903, p. IX-24	Minor revisions to language for clarifications.	M	DCH
07/2023	908, p. IX-45 – IX-47	Posted updated AAPD Dental Periodicity Schedule.	M	AAPD
07/2023	910, p. IX-48	Minor revisions to language for clarifications.	M	DCH
07/2023	1003, #15, p. X-7	Policy was updated to allow all applicable vaccines to be administered during school-based vaccine clinics. Providers should report POS 03.	M	DCH
07/2023	#17, p. X-8 #24, p. X-10	Removed 99241. CMS termination 12/31/22	D	CMS
07/2023	Appendix E	Posted updated ‘NEMT Fact Sheet and NEMT Regions & Counties Served’	M	DCH
07/2023	p. A-4, A-11, M-1&2, M-4	Address updates for DPH and DBHDD	M	DPH DBHDD
04/2023	902, p. IX-1	Effective April 1, 2023, DCH adopted the AAP 2023 BF Periodicity Schedule.	M	AAP
04/2023	902.2 E.2. p. IX-13	Updated reference for Child Core Set Technical Specifications and Resource Manual for FFY23	M	CMS
04/2023	902.2 E.6. p. IX-15	Updated AAP recommendation for depression screening – added the Patient Health Questionnaire (PHQ)-9 Modified for Teens (PHQ-A)	A	AAP
04/2023	905, p. IX-38 – IX-42	2023 CDC Immunization Schedules posted	M	CDC/ACIP

APPENDIX P

2023 Policy Revisions Record

Part II Policies and Procedures Manual for EPSDT (Health Check) Services

REVISION DATE	SECTION & PAGE	REVISION DESCRIPTION	REVISION TYPE A = Added D = Deleted M = Modified	CITATION (Revision required by Regulation, Legislation, etc.)
04/2023	907.4 p. IX-44	Lead Case Management - Updated environmental lead risk assessment with a confirmed blood lead level of 3.5 ug/dL	M	DPH GHHLPPP
04/2023	Appendix A	Updated Guidelines in Screening and Reporting Elevated Blood Lead Levels - Blood lead levels lowered to 3.5 ug/dl.	M	DPH GHHLPPP
04/2023	Appendix C-2	Added the following vaccines: - 90671 (PCV15) eff. 12/01/2022 - 90677 (PCV20) eff. 12/01/2022 - 90739 (HepB) eff. 01/01/2023	A	DCH/ DPH Immunization- VFC
04/2023	Appendix F F-4	Updated reference for: - Counseling for Nutrition Guidelines - Counseling for Physical Activity	M	DGA- US Dept of HHS
04/2023	Appendix O	Patient Health Questionnaire-9 (PHQ-9) – added	A	AAP
04/2023	Appendix Q	AAP 2023 Bright Futures Periodicity Schedule – summary of changes added	A	AAP
04/2023	Appendix R	Deleted 2021 CDC Immunization Schedules – summary of changes	D	DCH
04/2023	Appendix S	Updated PCP Rate Increases tables	M	DCH
02/2023	Appendix M	Corrected address for PeachCare for Kids. <i>P.O. Box 786 Alma, GA. 31510</i>	M	Gainwell DCH
01/2023	902, p. IX-1	Effective January 1, 2023, DCH adopted the AAP 2022 BF Periodicity Schedule.	M	AAP

APPENDIX Q

2023 Recommendations for Preventive Pediatric Health Care *(Copyright © 2023 by the AAP, updated April 2023)*

The Division will adopt the AAP 2023 Bright Futures “Recommendations for Pediatric Health Care” Periodicity Schedule effective date April 1, 2023.

The schedule is available at [Preventive Care/Periodicity Schedule \(aap.org\)](https://www.aap.org/periodicityschedule)
<https://www.aap.org/periodicityschedule>

This schedule reflects changes approved in December 2022 and published in April 2023.

Changes Made in December 2022

HIV

- The HIV screening recommendation has been updated to extend the upper age limit from 18 to 21 years (to account for the range in which the screening can take place) to align with recommendations of the US Preventive Services Task Force and AAP policy (“Adolescents and Young Adults: The Pediatrician’s Role in HIV Testing and Pre- and Postexposure HIV Prophylaxis”).
- Footnote 30 has been updated to read as follows: “Screen adolescents for HIV at least once between the ages of 15 and 21, making every effort to preserve confidentiality of the adolescent, as per ‘Human Immunodeficiency Virus (HIV) Infection: Screening’ (<https://www.uspreventiveservicestaskforce.org/uspstf/recommendation/human-immunodeficiency-virus-hiv-infection-screening>); after initial screening, youth at increased risk of HIV infection should be retested annually or more frequently, as per ‘Adolescents and Young Adults: The Pediatrician’s Role in HIV Testing and Pre- and Postexposure HIV Prophylaxis’ (<https://doi.org/10.1542/peds.2021-055207>).”

APPENDIX Q

2022 Recommendations for Preventive Pediatric Health Care (Copyright © 2022 by the AAP, updated July 2022)

The Division will adopt the AAP 2022 Bright Futures “Recommendations for Pediatric Health Care” Periodicity Schedule effective date January 1, 2023.

The schedule is available at https://www.aap.org/en-us/Documents/periodicity_schedule.pdf

The 2022 Bright Futures/AAP Periodicity Schedule is Now Available

The 2022 *Recommendations for Preventive Pediatric Health Care* [policy statement](#) (also known as the Periodicity Schedule) was published in the July 2022 issue of *Pediatrics* (released online on June 21, 2022). The Bright Futures/AAP Periodicity Schedule is updated annually. Be sure to check the date in the top right corner to ensure that you are referencing the most updated version. Here is a summary of the changes to the 2022 [Bright Futures/ AAP Periodicity Schedule](#):

Hepatitis B Virus Infection

- Assessing risk for HBV infection has been added to occur from newborn to 21 years (to account for the range in which the risk assessment can take place) to be consistent with recommendations of the USPSTF and the 2021–2024 edition of the AAP *Red Book: Report of the Committee on Infectious Diseases*.
- Footnote 31 has been added to read as follows: “Perform a risk assessment for hepatitis B virus (HBV) infection according to recommendations per the USPSTF (<https://www.uspreventiveservicestaskforce.org/uspstf/recommendation/hepatitis-b-virus-infection-screening>) and in the 2021– 2024 edition of the AAP *Red Book: Report of the Committee on Infectious Diseases*, making every effort to preserve confidentiality of the patient.”

Sudden Cardiac Arrest and Sudden Cardiac Death

- Assessing risk for sudden cardiac arrest and sudden cardiac death has been added to occur from 11 to 21 years (to account for the range in which the risk assessment can take place) to be consistent with AAP policy (“Sudden Death in the Young: Information for the Primary Care Provider”).
- Footnote 33 has been added to read as follows: “Perform a risk assessment, as appropriate, per ‘Sudden Death in the Young: Information for the Primary Care Provider’ (<https://doi.org/10.1542/peds.2021-052044>).”

Depression and Suicide Risk

- Screening for suicide risk has been added to the existing depression screening recommendation to be consistent with the GLAD-PC and AAP policy.
- Footnote 16 has been updated to read as follows: “Screen adolescents for depression and suicide risk, making every effort to preserve confidentiality of the adolescent. See ‘Guidelines for Adolescent Depression in Primary Care (GLAD-PC): Part I. Practice Preparation, Identification, Assessment, and Initial Management’ (<https://doi.org/10.1542/peds.2017-4081>), ‘Mental Health Competencies for Pediatric Practice’ (<https://doi.org/10.1542/peds.2019-2757>), ‘Suicide and Suicide Attempts in Adolescents’ (<https://doi.org/10.1542/peds.2016-1420>), and ‘The 21st Century Cures Act & Adolescent Confidentiality’ ([https://www.adolescenthealth.org/Advocacy/Advocacy-Activities/2019-\(1\)/NASPAG-SAHM-Statement.aspx](https://www.adolescenthealth.org/Advocacy/Advocacy-Activities/2019-(1)/NASPAG-SAHM-Statement.aspx)).”

APPENDIX Q

Behavioral/Social/Emotional

- The Psychosocial/Behavioral Assessment recommendation has been updated to Behavioral/Social/Emotional Screening (annually from newborn to 21 years) to align with AAP policy, the American College of Obstetricians and Gynecologists (Women’s Preventive Services Initiative) recommendations, and the American Academy of Child & Adolescent Psychiatry guidelines.
- Footnote 14 has been updated to read as follows: “Screen for behavioral and social-emotional problems per ‘Promoting Optimal Development: Screening for Behavioral and Emotional Problems’ (<https://doi.org/10.1542/peds.2014-3716>), ‘Mental Health Competencies for Pediatric Practice’ (<https://doi.org/10.1542/peds.2019-2757>), ‘Clinical Practice Guideline for the Assessment and Treatment of Children and Adolescents With Anxiety Disorders’ (<https://pubmed.ncbi.nlm.nih.gov/32439401>), and ‘Screening for Anxiety in Adolescent and Adult Women: A Recommendation From the Women’s Preventive Services Initiative’ (<https://pubmed.ncbi.nlm.nih.gov/32510990/>). The screening should be family centered and may include asking about caregiver emotional and mental health concerns and social determinants of health, racism, poverty, and relational health. See ‘Poverty and Child Health in the United States’ (<https://doi.org/10.1542/peds.2016-0339>), ‘The Impact of Racism on Child and Adolescent Health’ (<https://doi.org/10.1542/peds.2019-1765>), and ‘Preventing Childhood Toxic Stress: Partnering With Families and Communities to Promote Relational Health’ (<https://doi.org/10.1542/peds.2021-052582>).”

Fluoride Varnish

- Footnote 37 has been updated to read as follows: “The USPSTF recommends that primary care clinicians apply fluoride varnish to the primary teeth of all infants and children starting at the age of primary tooth eruption (<https://www.uspreventiveservicestaskforce.org/uspstf/recommendation/prevention-of-dental-caries-in-children-younger-than-age-5-years-screening-and-interventions1>). Once teeth are present, apply fluoride varnish to all children every 3 to 6 months in the primary care or dental office based on caries risk. Indications for fluoride use are noted in ‘Fluoride Use in Caries Prevention in the Primary Care Setting’ (<https://doi.org/10.1542/peds.2020-034637>).”

Fluoride Supplementation

- Footnote 38 has been updated to read as follows: “If primary water source is deficient in fluoride, consider oral fluoride supplementation. See ‘Fluoride Use in Caries Prevention in the Primary Care Setting’ (<https://doi.org/10.1542/peds.2020-034637>).”

2022 Coding for Pediatric Preventive Care Booklet is Available

In addition to the 2022 Bright Futures/AAP Periodicity Schedule, the 2022 [Coding for Pediatric Preventive Care](#) booklet has been revised and is available online and in print. This complementary resource contains comprehensive listings of codes and services that coincide with the 2022 Bright Futures/AAP Periodicity Schedule.

APPENDIX Q

2021 Recommendations for Preventive Pediatric Health Care

(Copyright © 2021 by the AAP, updated March 2021)

The Division will adopt the AAP 2021 Bright Futures “Recommendations for Pediatric Health Care” Periodicity Schedule effective date July 1, 2021.

The schedule is available at https://www.aap.org/en-us/Documents/periodicity_schedule.pdf

This schedule reflects changes approved in November 2020 and published in March 2021. For updates and a list of previous changes made, visit www.aap.org/periodicschedule.

PUBLISHED IN MARCH 2021, CHANGES MADE IN NOVEMBER 2020

DEVELOPMENTAL

- Footnote 11 has been updated to read as follows: “Screening should occur per ‘Promoting Optimal Development: Identifying infant and Young Children With Developmental Disorders Through Developmental Surveillance and Screening’ (<https://pediatrics.aappublications.org/content/145/1/e20193449>).”

AUTISM SPECTRUM DISORDER

- Footnote 12 has been updated to read as follows: “Screening should occur per ‘Identification, Evaluation, and Management of Children With Autism Spectrum Disorder’ (<https://pediatrics.aappublications.org/content/145/1/e20193447>).”

HEPATITIS C VIRUS INFECTION

- Screening for hepatitis C virus infection has been added to occur at least once between the ages of 18 and 79 years (to be consistent with recommendations of the USPSTF and CDC).
- Footnote 31 has been added to read as follows: “All individuals should be screened for hepatitis C virus (HCV) infection according to the USPSTF (<https://www.uspreventiveserivcestraskforce.org/uspstf/recommendation/hepatitis-c-screening>) and Centers for Disease Control and Prevention (CDC) recommendations (<https://www.cdc.gov/mmwr/volumes/69/rr/rr6902a1.htm>) at least once between use, should be tested for HCV infection and reassessed annually.”
- Footnotes 31 through 35 have been renumbered as footnotes 32 through 36.

APPENDIX Q

2019 Recommendations for Preventive Pediatric Health Care *(Copyright © 2019 by the AAP, updated March 2019)*

The Division will adopt the AAP 2019 Bright Futures “Recommendations for Pediatric Health Care” Periodicity Schedule effective date July 1, 2019.

The schedule is available at https://www.aap.org/en-us/Documents/periodicity_schedule.pdf

The changes implemented in the 2019 schedule include the following footnote revisions. The following is a summary of changes:

BLOOD PRESSURE

- Footnote 6 has been updated to read as follows: "Screening should occur per 'Clinical Practice Guideline for Screening and Management of High Blood Pressure in Children and Adolescents.' Blood pressure measurement in infants and children with specific risk conditions should be performed at visits before age 3 years."

ANEMIA

- Footnote 24 has been updated to read as follows: "Perform risk assessment or screening, as appropriate, per recommendations in the current edition of the AAP *Pediatric Nutrition: Policy of the American Academy of Pediatrics* (Iron chapter)."

LEAD

- Footnote 25 has been updated to read as follows: "For children at risk of lead exposure, see 'Prevention of Childhood Lead Toxicity' and 'Low Level Lead Exposure Harms Children: A Renewed Call for Primary Prevention'."

APPENDIX Q

2017 Recommendations for Preventive Pediatric Health Care *(Copyright © 2017 by the AAP, updated February 2017)*

The Division adopted the AAP 2017 Bright Futures “Recommendations for Pediatric Health Care” Periodicity Schedule effective date July 1, 2017.

The schedule is available at https://www.aap.org/en-us/documents/periodicity_schedule.pdf

The changes implemented in the 2017 schedule include numerous footnote revisions. The following is a summary of changes:

HEARING

- Timing and follow-up of the screening recommendations for hearing during the infancy visits have been delineated. Adolescent risk assessment has changed to screening once during each time period.
- Footnote 8 has been updated to read as follows: “Confirm initial screen was completed, verify results, and follow up, as appropriate. Newborns should be screened, per ‘Year 2007 Position Statement: Principles and Guidelines for Early Hearing Detection and Intervention Programs’ <http://pediatrics.aappublications.org/content/120/4/898.full>”
- Footnote 9 has been added to read as follows: “Verify results as soon as possible, and follow up, as appropriate.”
- Footnote 10 has been added to read as follows: “Screen with audiometry including 6,000 and 8,000 Hz high frequencies once between 11 and 14 years, once between 15 and 17 years, and once between 18 and 21 years. See ‘The Sensitivity of Adolescent Hearing Screens Significantly Improves by Adding High Frequencies’ [http://www.jahonline.org/article/S1054-139X\(16\)00048-3/fulltext](http://www.jahonline.org/article/S1054-139X(16)00048-3/fulltext)”

PSYCHOSOCIAL/BEHAVIORAL ASSESSMENT

- Footnote 13 has been added to read as follows: “This assessment should be family centered and may include an assessment of child social-emotional health, caregiver depression, and social determinants of health. See ‘Promoting Optimal Development: Screening for Behavioral and Emotional Problems’ (<http://pediatrics.aappublications.org/content/135/2/384>) and ‘Poverty and Child Health in the United States’ (<http://pediatrics.aappublications.org/content/137/4/e20160339>).”

APPENDIX Q

TOBACCO, ALCOHOL, OR DRUG USE ASSESSMENT

- The header was updated to be consistent with recommendations

DEPRESSION SCREENING

- Adolescent depression screening begins routinely at 12 years of age (to be consistent with recommendations of the US Preventive Services Task Force [USPSTF]).

MATERNAL DEPRESSION SCREENING

- Screening for maternal depression at 1-, 2-, 4-, and 6-month visits has been added.
- Footnote 16 was added to read as follows: “Screening should occur per ‘Incorporating Recognition and Management of Perinatal and Postpartum Depression Into Pediatric Practice’ (<http://pediatrics.aappublications.org/content/126/5/1032>).”

NEWBORN BLOOD

- Timing and follow-up of the newborn blood screening recommendations have been delineated.
- Footnote 19 has been updated to read as follows: “Confirm initial screen was accomplished, verify results, and follow up, as appropriate. The Recommended Uniform Newborn Screening Panel (<https://www.hrsa.gov/advisory-committees/heritable-disorders/rusp/index.html>), as determined by The Secretary’s Advisory Committee on Heritable Disorders in Newborns and Children, and state newborn screening laws/regulations (<http://genes-r-us.uthscsa.edu/sites/genes-r-us/files/nbsdisorders.pdf>) establish the criteria for and coverage of newborn screening procedures and programs.”
- Footnote 20 has been added to read as follows: “Verify results as soon as possible, and follow up, as appropriate.”

NEWBORN BILIRUBIN

- Screening for bilirubin concentration at the newborn visit has been added.
- Footnote 21 has been added to read as follows: “Confirm initial screening was accomplished, verify results, and follow up, as appropriate. See ‘Hyperbilirubinemia in the Newborn Infant ≥ 35 Weeks’ Gestation: An Update with Clarifications’ (<http://pediatrics.aappublications.org/content/124/4/1193>).”

APPENDIX Q

DYSLIPIDEMIA

- Screening for dyslipidemia has been updated to occur once between 9 and 11 years of age, and once between 17 and 21 years of age (to be consistent with guidelines of the National Heart, Lung, and Blood Institute).

SEXUALLY TRANSMITTED INFECTIONS

- Footnote 29 has been updated to read as follows: “Adolescents should be screened for sexually transmitted infections (STIs) per recommendations in the current edition of the AAP Red Book: Report of the Committee on Infectious Diseases.”

HIV

- A subheading has been added for the HIV universal recommendation to avoid confusion with STIs selective screening recommendation.
- Screening for HIV has been updated to occur once between 15 and 18 years of age (to be consistent with recommendations of the USPSTF).
- Footnote 30 has been added to read as follows: “Adolescents should be screened for HIV according to the USPSTF recommendations (<http://www.uspreventiveservicestaskforce.org/uspstf/uspshivi.htm>) once between the ages of 15 and 18, making every effort to preserve confidentiality of the adolescent. Those at increased risk of HIV infection, including those who are sexually active, participate in injection drug use, or are being tested for other STIs, should be tested for HIV and reassessed annually.”

ORAL HEALTH

- Assessing for a dental home has been updated to occur at the 12-month and 18-month through 6-year visits. A subheading has been added for fluoride supplementation, with a recommendation from the 6-month through 12-month and 18-month through 16-year visits.
- Footnote 32 has been updated to read as follows: “Assess whether the child has a dental home. If no dental home is identified, perform a risk assessment https://www.aap.org/en-us/Documents/oralhealth_RiskAssessmentTool.pdf and refer to a dental home. Recommend brushing with fluoride toothpaste in the proper dosage for age. See ‘Maintaining and Improving the Oral Health of Young Children’ (<http://pediatrics.aappublications.org/content/134/6/1224>)”
- Footnote 33 has been updated to read as follows: “Perform a risk assessment (<http://www2.aap.org/oralhealth/docs/RiskAssessmentTool.pdf>). See ‘Maintaining and Improving the Oral Health of Young Children’ (<http://pediatrics.aappublications.org/content/134/6/1224>).”
- Footnote 35 has been added to read as follows: “If primary water source is deficient in fluoride, consider oral fluoride supplementation. See ‘Fluoride Use in Caries Prevention in the Primary Care Setting’ (<http://pediatrics.aappublications.org/content/134/3/626>).”

Rev. 04/23

APPENDIX R

INTENTIONALLY LEFT BLANK

APPENDIX S

**EPSDT Health Check Program (COS 600)
Reimbursement Rates for
Medicaid-Eligible Members & PeachCare for Kids® (PCK)-Eligible
Members**

Physicians and physician extenders (physician assistants, nurse practitioners) are reimbursed at 100% of the established rates when billing the specified codes and modifiers for Health Check services rendered to Medicaid-eligible members and PCK-eligible members.

Physicians and physician extenders who are eligible for the House Bill (HB) Primary Care Providers (PCP) rate increases are reimbursed 100% of the established rates, when billing the specified codes and modifiers for Health Check services rendered to Medicaid-eligible and PCK-eligible members.

	<u>PCP Rate Increase</u>	<u>Fiscal Year</u>	<u>Effective Date</u>
Rev. 04/20	HB 76	FY 2016	eff. July 1, 2015
Rev. 07/20	HB 751	FY 2017	eff. July 1, 2016
Rev. 04/23	HB 44	FY 2018	eff. July 1, 2017
	HB 793	FY 2021	eff. Aug. 14, 2020
			eff. July 1, 2021

APPENDIX S

**EPSDT Health Check Program (COS 600)
Reimbursement Rates for
Medicaid-Eligible Members & PeachCare for Kids® (PCK)-Eligible Members**

Preventive Visit Codes	HIPAA Modifiers	Rates for Non-Attested Providers	HB 76 (eff. July 1, 2015)	HB 751 (eff. July 1, 2016)	HB 44 (eff. July 1, 2017)	HB 793 (eff. Aug. 14, 2020)	HB 793 (eff. July 1, 2021)
99381	EP EP HA EP 25 EP HA 25	\$67.38 ¹		\$106.68		\$107.75	
99391			\$86.47	\$96.08		\$97.04	\$99.99
99382					\$111.27	\$112.38	
99392			\$92.46	\$102.74		\$103.77	\$106.92
99383 (age 5-7 years)	EP EP 25	\$67.38 ¹			\$116.19	\$117.35	
99393 (age 5-7 years)			\$92.17	\$102.41		\$103.43	\$106.58
99383 (age 8-11 years)	EP EP 25	\$75.38 - private \$55.38 – public health			\$116.19	\$117.35	
99393 (age 8-11 years)			\$92.17	\$102.41		\$103.43	\$106.58
99384	EP EP 25	\$75.38 - private \$55.38 – public health			\$131.62	\$132.94	
99394			\$101.03	\$112.25		\$113.37	\$116.82
99385	EP EP 25	\$75.38 - private \$55.38 – public health			\$127.75	\$129.03	
99395			\$103.24	\$114.71		\$115.86	
¹ This applies to non-attested private and public health providers.							
When reporting the preventive visit codes with EP, EP 25, EP HA, EP HA 25 modifier(s), reimbursement is at 100% for physicians and mid-level providers (physician assistants, nurse practitioners).							

APPENDIX S

**EPSDT Health Check Program (COS 600)
Reimbursement Rates for
Medicaid-Eligible Members & PeachCare for Kids® (PCK)-Eligible Members**

Interperiodic Visit Codes & Modifiers (EP, EP 25)	Rates for Non-Attested Providers		HB 76 (eff. July 1, 2015)	HB 751 (eff. July 1, 2016)	HB 44 (eff. July 1, 2017)	HB 793 (eff. Aug. 14, 2020)	HB 793 (eff. July 1, 2021)
	Medicaid-eligible members	PCK-eligible members					
99201 ⁴ deleted eff. 1-1-2021	\$35.13	\$41.20			\$41.30 ¹	\$41.71 ¹	
99202	\$54.57	\$71.16		\$71.33 ¹		\$72.04 ¹	
99203	\$76.53	\$103.01		\$103.80 ¹		\$104.84 ¹	\$109.35 ¹
99204	\$110.51	\$158.37					\$167.09 ¹
99205	\$137.12	\$197.18				\$202.13 ¹	
99211	\$17.46	\$17.46			\$18.97 ² \$19.79 ³	\$19.16 ² \$19.99 ³	
99212	\$29.67	\$41.54		\$41.63 ¹		\$42.05 ¹	\$46.19 ¹
99213	\$40.70	\$69.11	\$63.14 ²	\$70.15 ¹		\$70.85 ¹	\$76.15 ¹
99214	\$62.71	\$102.49		\$103.72 ¹		\$104.76 ¹	\$110.43 ¹
99215	\$93.46	\$137.88					\$148.33 ¹
¹ This rate increase applies to Medicaid-eligible members and PeachCare for Kids®-eligible members. ² This rate increase does not apply to PeachCare for Kids®-eligible members. ³ This rate increase does not apply to Medicaid-eligible members. ⁴ CPT code 99201 deleted effective 1-1-2021 CPT codes 99204, 99205, 99215 added effective 1-1-2024							
When reporting the interperiodic visit codes with EP, EP 25 modifier(s), reimbursement is at 100% for physicians and mid-level providers (physician assistants, nurse practitioners)							

APPENDIX S

**EPSDT Health Check Program (COS 600)
Reimbursement Rates for
Medicaid-Eligible Members & PeachCare for Kids® (PCK)-Eligible Members**

Immunization Codes & Modifier (EP)	Rates for Non-Attested Providers		HB 751 (eff. July 1, 2016)		HB 44 (eff. July 1, 2017)		HB 793 (eff. Aug. 14, 2020)		HB 793 (eff. July 1, 2021)
	Medicaid-eligible members	PCK-eligible members	Medicaid-eligible members	PCK-eligible members	Medicaid-eligible members	PCK-eligible members	Medicaid-eligible members	PCK-eligible members	Medicaid & PCK-eligible members
90460	\$10.00	\$18.50	\$21.93	\$21.93			\$22.15	\$22.15	
90471	\$10.00	\$18.50	\$23.54	\$23.54			\$23.78 ¹	\$23.78 ¹	
90472	\$10.00	\$18.50	\$11.98	\$18.50			\$12.10	\$18.69	\$12.99
90473	\$10.00	\$18.50			\$23.54	\$23.54	\$23.78	\$23.78	
90474	\$10.00	\$18.50			\$11.98	\$18.50	\$12.10	\$18.69	
When reporting the immunization codes with EP modifier, reimbursement is at 100% for physicians and mid-level providers (physician assistants, nurse practitioners)									
¹ effective 6/30/2020									